myriad WOMEN'S HEALTH

RESULTS RECIPIENT
SEATTLE SPERM BANK
Attn: Jeffrey Olliffe
4915 25th Ave NE Ste 204W
Seattle, WA 98105
Phone: (206) 588-1484
Fax: (206) 466-4696
NPI: 1306838271
Report Date: 01/22/2021

MALE
DONOR 14184
DOB: [redacted]
Ethnicity: East Asian
Sample Type: EDTA Blood
Date of Collection: 01/08/2021
Date Received: 01/13/2021
Date Tested: 01/20/2021
Barcode: 11004512825040
Accession ID: CSLZ33H926YYMF6
Indication: Egg or sperm donor

FEMALE
N/A

Foresight® Carrier Screen

NEGATIVE

ABOUT THIS TEST

The **Myriad Foresight Carrier Screen** utilizes sequencing, maximizing coverage across all DNA regions tested, to help you learn about your chance to have a child with a genetic disease.

RESULTS SUMMARY

Risk Details	DONOR 14184	Partner
Panel Information	Foresight Carrier Screen Universal Panel Fundamental Plus Panel Fundamental Panel **(175 conditions tested)**	N/A
All conditions tested A complete list of all conditions tested can be found on page 5.	⊖ **NEGATIVE** No disease-causing mutations were detected.	N/A

CLINICAL NOTES

- None

NEXT STEPS

- If necessary, patients can discuss residual risks with their physician or a genetic counselor.

Copyright 2021 Myriad Women's Health, Inc. All rights reserved.

Methods and Limitations

DONOR 14184 [Foresight Carrier Screen]: Sequencing with copy number analysis, spinal muscular atrophy, analysis of homologous regions, and alpha thalassemia (HBA1/HBA2) sequencing with targeted copy number analysis (Assay(s): DTS v3.2).

Sequencing with copy number analysis

High-throughput sequencing and read-depth-based copy number analysis are used to analyze the genes listed in the Conditions Tested section of the report. Except where otherwise noted, the region of interest (ROI) comprises the indicated coding regions and 20 non-coding bases flanking each region. In a minority of cases where genomic features (e.g., long homopolymers) compromise calling fidelity, the affected non-coding bases are excluded from the ROI. The ROI is sequenced to a minimum acceptable read depth, and the sequences are compared to a reference genomic sequence (Genome Reference Consortium Human Build 37 [GRCh37]/hg19). On average, 99% of all bases in the ROI are sequenced at a read depth that is greater than the minimum read depth. Sequence variants may not be detected in areas of lower sequence coverage. Insertions and deletions may not be detected as accurately as single-nucleotide variants. Select genes or regions for which pseudogenes or other regions of homology impede reliable variant detection may be assayed using alternate technology, or they may be excluded from the ROI. *CFTR* and *DMD* testing includes analysis for exon-level deletions and duplications with an average sensitivity of ~99%. Only exon-level deletions are assayed for other genes on the panel and such deletions are detected with a sensitivity of ≥75%. Selected founder deletions may be detected at slightly higher sensitivity. Affected exons and/or breakpoints of copy number variants are estimated from junction reads, where available, or using the positions of affected probes. Only exons known to be included in the region affected by a copy number variant are provided in the variant nomenclature. In some cases, the copy number variant may be larger or smaller than indicated. If *GJB2* is tested, large upstream deletions involving the *GJB6* and/or *CRYL1* genes that may affect the expression of *GJB2* are also analyzed.

Spinal muscular atrophy

Targeted copy number analysis via high-throughput sequencing is used to determine the copy number of exon 7 of the *SMN1* gene. Other genetic variants may interfere with this analysis. Some individuals with two copies of *SMN1* are "silent" carriers with both *SMN1* genes on one chromosome and no copies of the gene on the other chromosome. This is more likely in individuals who have two copies of the *SMN1* gene and are positive for the g.27134T>G single-nucleotide polymorphism (SNP) (PMID: 9199562, 23788250, and 28676062), which affects the reported residual risk; Ashkenazi Jewish or Asian patients with this genotype have a high post-test likelihood of being carriers for SMA and are reported as carriers. The g.27134T>G SNP is only reported in individuals who have two copies of *SMN1*.

Analysis of homologous regions

A combination of high-throughput sequencing, read-depth-based copy number analysis, and targeted genotyping is used to determine the number of functional gene copies and/or the presence of selected loss-of-function variants in certain genes that have homology to other genomic regions. The precise breakpoints of large deletions in these genes cannot be determined but are instead estimated from copy number analysis. Pseudogenes may interfere with this analysis, especially when many pseudogene copies are present.

If *CYP21A2* is tested, patients who have one or more additional copies of the *CYP21A2* gene and a pathogenic variant may or may not be a carrier of 21-hydroxylase deficient CAH, depending on the chromosomal location of the variants (phase). Benign *CYP21A2* gene duplications and/or triplications will only be reported in this context. Some individuals with two functional *CYP21A2* gene copies may be "silent" carriers, with two gene copies resulting from a duplication on one chromosome and a gene deletion on the other chromosome. This and other similar rare carrier states, where complementary changes exist between the chromosomes, may not be detected by the assay. Given that the true incidence of non-classic CAH is unknown, the residual carrier and reproductive risk numbers on the report are based only on the published incidence for classic CAH. However, the published prevalence of non-classic CAH is highest in individuals of Ashkenazi Jewish, Hispanic, Italian, and Yugoslav descent. Therefore, the residual and reproductive risks are likely an underestimate for CAH, especially in the aforementioned populations, as they do not account for non-classic CAH.

Copyright 2021 Myriad Women's Health, Inc. All rights reserved.

Alpha thalassemia (HBA1/HBA2) sequencing with targeted copy number analysis

High-throughput sequencing and read-depth-based copy number analysis are used to identify sequence variation and functional gene copies within the region of interest (ROI) of *HBA1* and *HBA2*, which includes the listed exons plus 20 intronic flanking bases. In a minority of cases where genomic features (e.g , long homopolymers) compromise calling fidelity, the affected intronic bases are not included in the ROI. The ROI is sequenced to a minimum acceptable read depth, and the sequences are compared to a reference genomic sequence (Genome Reference Consortium Human Build 37 [GRCh37]/hg19). On average, 99% of all bases in the ROI are sequenced at a read depth that is greater than the minimum read depth. Sequence variants may not be detected in areas of lower sequence coverage. Insertions and deletions may not be detected as accurately as single-nucleotide variants. For large deletions or duplications in these genes, the precise breakpoints cannot be determined but are instead estimated from copy number analysis. This assay has been validated to detect up to two additional copies of each alpha globin gene. In rare instances where assay results suggest greater than two additional copies are present, this will be noted but the specific number of gene copies observed will not be provided.

Extensive sequence homology exists between *HBA1* and *HBA2*. This sequence homology can prevent certain variants from being localized to one gene over the other. In these instances, variant nomenclature will be provided for both genes. If follow-up testing is indicated for patients with the nomenclature provided for both genes, both *HBA1* and *HBA2* should be tested. Some individuals with four functional alpha globin gene copies may be "silent" carriers, with three gene copies resulting from triplication on one chromosome and a single gene deletion on the other chromosome. This and other similar rare carrier states, where complementary changes exist between the chromosomes, may not be detected by the assay.

Interpretation of reported variants

Variants are classified according to internally defined criteria, which are compatible with the ACMG Standards and Guidelines for the Interpretation of Sequence Variants (PMID: 25741868). Variants that are considered disease-causing by Myriad Women's Health (MWH) are reported. In addition, variants that have not previously been established as a recognized cause of disease may be identified. In these cases, only variants classified as 'likely' pathogenic are reported. Likely pathogenic variants are described elsewhere in the report as 'likely to have a negative impact on gene function.' Variant pathogenicity is evaluated and classified by an assessment of the nature of the variant and reviews of reports of allele frequencies in cases and controls, functional studies, variant annotation and effect prediction, segregation studies, and other resources, where available. Exon-level duplications in the *DMD* and *CFTR* genes are assumed to be in tandem and are classified according to their predicted effect on the reading frame. Benign variants, variants of uncertain significance, and variants not directly associated with the specified disease phenotype(s) are not reported.

Limitations

The MWH Foresight Carrier Screen is designed to detect and report germline (constitutional) alterations. Mosaic (somatic) variation may not be detected, and if it is detected, it may not be reported. If more than one variant is detected in a gene, additional studies may be necessary to determine if those variants lie on the same chromosome or different chromosomes (phase). This test is not designed to detect sex-chromosome copy number variations. If present, sex-chromosome abnormalities may significantly reduce test sensitivity for X-linked conditions. Variant interpretation and residual and reproductive risk estimations assume a normal karyotype and may be different for individuals with abnormal karyotypes. The test does not fully address all inherited forms of intellectual disability, birth defects, or heritable diseases. Furthermore, not all forms of genetic variation are detected by this assay (i.e., duplications [except in specified genes], chromosomal rearrangements, structural abnormalities, etc.). Additional testing may be appropriate for some individuals. Pseudogenes and other regions of homology may interfere with this analysis. In an unknown number of cases, other genetic variation may interfere with variant detection. Rare carrier states where complementary changes exist between the chromosomes may not be detected by the assay. Other possible sources of diagnostic error include sample mix-up, trace contamination, bone marrow transplantation, blood transfusions, and technical or analytical errors.

Detection rates are determined using published scientific literature and/or reputable databases, when available, to estimate the fraction of disease alleles, weighted by frequency, that the methodology is predicted to be able or unable to detect. Detection rates are approximate and only account for analytical sensitivity. Certain variants that have been previously described in the literature may not be reported, if there is insufficient evidence for pathogenicity. Detection rates do not account for the disease specific rates of *de novo* variation.

This test was developed, and its performance characteristics determined by, Myriad Women's Health, Inc. It has not been cleared or approved by the US Food and Drug Administration (FDA). The FDA does not require this test to go through premarket review. This test is used for clinical purposes. It should not be regarded as investigational or for research. This laboratory is certified under the Clinical Laboratory Improvement Amendments of 1988 (CLIA) as qualified to perform high-complexity clinical testing. These results are adjunctive to the ordering physician's evaluation. CLIA Number: #05D1102604.

Incidental Findings

Copyright 2021 Myriad Women's Health, Inc. All rights reserved.

Unless otherwise indicated, these results and interpretations are limited to the specific disease panel(s) requested by the ordering healthcare provider. In some cases, standard data analyses may identify genetic findings beyond the region(s) of interest specified by the test, and such findings may not be reported. These findings may include genomic abnormalities with major, minor, or no, clinical significance.

If you have questions or would like more information about any of the test methods or limitations, please contact (888) 268-6795.

Resources

GENOME CONNECT | http://www.genomeconnect.org

Patients can share their reports using research registries such as Genome Connect, an online research registry building a genetics and health knowledge base. Genome Connect provides patients, physicians, and researchers an opportunity to share genetic information to support the study of the impact of genetic variation on health conditions.

SENIOR LABORATORY DIRECTOR



Karla R. Bowles, PhD, FACMG, CGMB

Report content approved by Karla Bowles, PhD, FACMG, CGMB on Jan 22, 2021

Copyright 2021 Myriad Women's Health, Inc. All rights reserved.

Conditions Tested

11-beta-hydroxylase-deficient Congenital Adrenal Hyperplasia - **Gene:** CYP11B1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000497:1-9. **Detection Rate:** East Asian 94%.

6-pyruvoyl-tetrahydropterin Synthase Deficiency - **Gene:** PTS. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000317:1-6. **Detection Rate:** East Asian >99%.

ABCC8-related Familial Hyperinsulinism - **Gene:** ABCC8. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000352:1-39. **Detection Rate:** East Asian >99%.

Adenosine Deaminase Deficiency - **Gene:** ADA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000022:1-12. **Detection Rate:** East Asian >99%.

Alpha Thalassemia, HBA1/HBA2-related - **Genes:** HBA1, HBA2. Autosomal Recessive. Alpha thalassemia (HBA1/HBA2) sequencing with targeted copy number analysis. **Exons:** NM_000517:1-3; NM_000558:1-3. **Variants (19):** -(alpha)20.5, --BRIT, --MEDI, --MEDII, --SEA, --THAI or --FIL, -alpha3.7, -alpha4.2, HBA1+HBA2 deletion, Hb Constant Spring, Poly(A) AATAAA>AATA--, Poly(A) AATAAA>AATAAG, Poly(A) AATAAA>AATGAA, anti3.7, anti4.2, c.*93A>G, c.*94_*95delAA, c.*95A>G, del HS-40. **Detection Rate:** East Asian >99%.

Alpha-mannosidosis - **Gene:** MAN2B1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000528:1-23. **Detection Rate:** East Asian >99%.

Alpha-sarcoglycanopathy - **Gene:** SGCA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000023:1-9. **Detection Rate:** East Asian >99%.

Alstrom Syndrome - **Gene:** ALMS1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_015120:1-23. **Detection Rate:** East Asian >99%.

AMT-related Glycine Encephalopathy - **Gene:** AMT. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000481:1-9. **Detection Rate:** East Asian >99%.

Andermann Syndrome - **Gene:** SLC12A6. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_133647:1-25. **Detection Rate:** East Asian >99%.

Argininemia - **Gene:** ARG1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000045:1-8. **Detection Rate:** East Asian 97%.

Argininosuccinic Aciduria - **Gene:** ASL. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001024943:1-16. **Detection Rate:** East Asian >99%.

Aspartylglucosaminuria - **Gene:** AGA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000027:1-9. **Detection Rate:** East Asian >99%.

Ataxia with Vitamin E Deficiency - **Gene:** TTPA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000370:1-5. **Detection Rate:** East Asian >99%.

Ataxia-telangiectasia - **Gene:** ATM. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000051:2-63. **Detection Rate:** East Asian >99%.

ATP7A-related Disorders - **Gene:** ATP7A. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000052:2-23. **Detection Rate:** East Asian 92%.

Autoimmune Polyglandular Syndrome Type 1 - **Gene:** AIRE. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000383:1-14. **Detection Rate:** East Asian >99%.

Autosomal Recessive Osteopetrosis Type 1 - **Gene:** TCIRG1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_006019:2-20. **Detection Rate:** East Asian >99%.

Autosomal Recessive Polycystic Kidney Disease, PKHD1-related - **Gene:** PKHD1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_138694 2-67. **Detection Rate:** East Asian >99%.

Autosomal Recessive Spastic Ataxia of Charlevoix-Saguenay - **Gene:** SACS. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_014363 2-10. **Detection Rate:** East Asian 99%.

Bardet-Biedl Syndrome, BBS1-related - **Gene:** BBS1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_024649:1-17. **Detection Rate:** East Asian >99%.

Bardet-Biedl Syndrome, BBS10-related - **Gene:** BBS10. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_024685:1-2. **Detection Rate:** East Asian >99%.

Bardet-Biedl Syndrome, BBS12-related - **Gene:** BBS12. Autosomal Recessive. Sequencing with copy number analysis. **Exon:** NM_152618:2. **Detection Rate:** East Asian >99%.

Bardet-Biedl Syndrome, BBS2-related - **Gene:** BBS2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_031885:1-17. **Detection Rate:** East Asian >99%.

BCS1L-related Disorders - **Gene:** BCS1L. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_004328:3-9. **Detection Rate:** East Asian >99%.

Beta-sarcoglycanopathy - **Gene:** SGCB. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000232:1-6. **Detection Rate:** East Asian >99%.

Biotinidase Deficiency - **Gene:** BTD. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000060:1-4. **Detection Rate:** East Asian >99%.

Bloom Syndrome - **Gene:** BLM. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000057:2-22. **Detection Rate:** East Asian >99%.

Calpainopathy - **Gene:** CAPN3. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000070:1-24. **Detection Rate:** East Asian >99%.

Canavan Disease - **Gene:** ASPA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000049:1-6. **Detection Rate:** East Asian 98%.

Carbamoylphosphate Synthetase I Deficiency - **Gene:** CPS1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001875:1-38. **Detection Rate:** East Asian >99%.

Carnitine Palmitoyltransferase IA Deficiency - **Gene:** CPT1A. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001876:2-19. **Detection Rate:** East Asian >99%.

Carnitine Palmitoyltransferase II Deficiency - **Gene:** CPT2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000098:1-5. **Detection Rate:** East Asian >99%.

Cartilage-hair Hypoplasia - **Gene:** RMRP. Autosomal Recessive. Sequencing with copy number analysis. **Exon:** NR_003051:1. **Detection Rate:** East Asian >99%.

Cerebrotendinous Xanthomatosis - **Gene:** CYP27A1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000784:1-9. **Detection Rate:** East Asian >99%.

Citrullinemia Type 1 - **Gene:** ASS1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000050:3-16. **Detection Rate:** East Asian 86%.

CLN3-related Neuronal Ceroid Lipofuscinosis - **Gene:** CLN3. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001042432 2-16. **Detection Rate:** East Asian >99%.

CLN5-related Neuronal Ceroid Lipofuscinosis - **Gene:** CLN5. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_006493:1-4. **Detection Rate:** East Asian >99%.

CLN6-related Neuronal Ceroid Lipofuscinosis - **Gene:** CLN6. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_017882:1-7. **Detection Rate:** East Asian >99%.

CLN8-related Neuronal Ceroid Lipofuscinosis - **Gene:** CLN8. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_018941:2-3. **Detection Rate:** East Asian >99%.

Cohen Syndrome - **Gene:** VPS13B. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_017890:2-62. **Detection Rate:** East Asian 97%.

COL4A3-related Alport Syndrome - **Gene:** COL4A3. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000091:1-52. **Detection Rate:** East Asian 97%.

COL4A4-related Alport Syndrome - **Gene:** COL4A4. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000092:2-48. **Detection Rate:** East Asian 98%.

Copyright 2021 Myriad Women's Health, Inc. All rights reserved.

Combined Pituitary Hormone Deficiency, PROP1-related - **Gene:** PROP1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_006261:1-3. **Detection Rate:** East Asian >99%.

Congenital Adrenal Hyperplasia, CYP21A2-related - **Gene:** CYP21A2. Autosomal Recessive. Analysis of homologous regions. **Variants (13):** CYP21A2 deletion, CYP21A2 duplication, CYP21A2 triplication, G111Vfs*21, I173N, L308Ffs*6, P31L, Q319*, Q319*+CYP21A2dup, R357W, V282L, [I237N;V238E;M240K], c.293-13C>G. **Detection Rate:** East Asian 88%.

Congenital Disorder of Glycosylation Type Ia - **Gene:** PMM2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000303:1-8. **Detection Rate:** East Asian >99%.

Congenital Disorder of Glycosylation Type Ic - **Gene:** ALG6. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_013339:2-15. **Detection Rate:** East Asian >99%.

Congenital Disorder of Glycosylation, MPI-related - **Gene:** MPI. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_002435:1-8. **Detection Rate:** East Asian >99%.

Costeff Optic Atrophy Syndrome - **Gene:** OPA3. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_025136:1-2. **Detection Rate:** East Asian >99%.

Cystic Fibrosis - **Gene:** CFTR. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000492:1-27. IVS8-5T allele analysis is only reported in the presence of the R117H mutation. **Detection Rate:** East Asian >99%.

Cystinosis - **Gene:** CTNS. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_004937:3-12. **Detection Rate:** East Asian >99%.

D-bifunctional Protein Deficiency - **Gene:** HSD17B4. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000414:1-24. **Detection Rate:** East Asian 98%.

Delta-sarcoglycanopathy - **Gene:** SGCD. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000337:2-9. **Detection Rate:** East Asian 99%.

Dihydrolipoamide Dehydrogenase Deficiency - **Gene:** DLD. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000108:1-14. **Detection Rate:** East Asian >99%.

Dysferlinopathy - **Gene:** DYSF. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_003494:1-55. **Detection Rate:** East Asian 98%.

Dystrophinopathy (Including Duchenne/Becker Muscular Dystrophy) - **Gene:** DMD. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_004006:1-79. **Detection Rate:** East Asian >99%.

ERCC6-related Disorders - **Gene:** ERCC6. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000124:2-21. **Detection Rate:** East Asian 99%.

ERCC8-related Disorders - **Gene:** ERCC8. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000082:1-12. **Detection Rate:** East Asian 95%.

EVC-related Ellis-van Creveld Syndrome - **Gene:** EVC. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_153717:1-21. **Detection Rate:** East Asian 96%.

EVC2-related Ellis-van Creveld Syndrome - **Gene:** EVC2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_147127:1-22. **Detection Rate:** East Asian >99%.

Fabry Disease - **Gene:** GLA. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000169:1-7. **Detection Rate:** East Asian 98%.

Familial Dysautonomia - **Gene:** IKBKAP. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_003640:2-37. **Detection Rate:** East Asian >99%.

Familial Mediterranean Fever - **Gene:** MEFV. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000243:1-10. **Detection Rate:** East Asian >99%.

Fanconi Anemia Complementation Group A - **Gene:** FANCA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000135:1-43. **Detection Rate:** East Asian 92%.

Fanconi Anemia, FANCC-related - **Gene:** FANCC. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000136:2-15. **Detection Rate:** East Asian >99%.

FKRP-related Disorders - **Gene:** FKRP. Autosomal Recessive. Sequencing with copy number analysis. **Exon:** NM_024301:4. **Detection Rate:** East Asian >99%.

FKTN-related Disorders - **Gene:** FKTN. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001079802:3-11. **Detection Rate:** East Asian 10%.

Free Sialic Acid Storage Disorders - **Gene:** SLC17A5. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_012434:1-11. **Detection Rate:** East Asian 98%.

Galactokinase Deficiency - **Gene:** GALK1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000154:1-8. **Detection Rate:** East Asian >99%.

Galactosemia - **Gene:** GALT. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000155:1-11. **Detection Rate:** East Asian >99%.

Gamma-sarcoglycanopathy - **Gene:** SGCG. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000231:2-8. **Detection Rate:** East Asian 88%.

Gaucher Disease - **Gene:** GBA. Autosomal Recessive. Analysis of homologous regions. **Variants (10):** D409V, D448H, IVS2+1G>A, L444P, N370S, R463C, R463H, R496H, V394L, p.L29Afs*18. **Detection Rate:** East Asian 60%.

GJB2-related DFNB1 Nonsyndromic Hearing Loss and Deafness - **Gene:** GJB2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_004004:1-2. **Detection Rate:** East Asian >99%.

GLB1-related Disorders - **Gene:** GLB1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000404:1-16. **Detection Rate:** East Asian >99%.

GLDC-related Glycine Encephalopathy - **Gene:** GLDC. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000170:1-25. **Detection Rate:** East Asian 94%.

Glutaric Acidemia, GCDH-related - **Gene:** GCDH. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000159:2-12. **Detection Rate:** East Asian >99%.

Glycogen Storage Disease Type Ia - **Gene:** G6PC. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000151:1-5. **Detection Rate:** East Asian >99%.

Glycogen Storage Disease Type Ib - **Gene:** SLC37A4. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001164277 3-11. **Detection Rate:** East Asian >99%.

Glycogen Storage Disease Type III - **Gene:** AGL. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000642:2-34. **Detection Rate:** East Asian >99%.

GNE Myopathy - **Gene:** GNE. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001128227:1-12. **Detection Rate:** East Asian >99%.

GNPTAB-related Disorders - **Gene:** GNPTAB. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_024312:1-21. **Detection Rate:** East Asian 98%.

HADHA-related Disorders - **Gene:** HADHA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000182:1-20. **Detection Rate:** East Asian >99%.

Hb Beta Chain-related Hemoglobinopathy (Including Beta Thalassemia and Sickle Cell Disease) - **Gene:** HBB. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000518:1-3. **Detection Rate:** East Asian >99%.

Hereditary Fructose Intolerance - **Gene:** ALDOB. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000035:2-9. **Detection Rate:** East Asian >99%.

Herlitz Junctional Epidermolysis Bullosa, LAMB3-related - **Gene:** LAMB3. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000228 2-23. **Detection Rate:** East Asian >99%.

Hexosaminidase A Deficiency (Including Tay-Sachs Disease) - **Gene:** HEXA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000520:1-14. **Detection Rate:** East Asian >99%.

HMG-CoA Lyase Deficiency - **Gene:** HMGCL. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000191:1-9. **Detection Rate:** East Asian 98%.

Holocarboxylase Synthetase Deficiency - **Gene:** HLCS. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000411:4-12. **Detection Rate:** East Asian >99%.

Homocystinuria, CBS-related - **Gene:** CBS. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000071:3-17. **Detection Rate:** East Asian >99%.

Hydrolethalus Syndrome - **Gene:** HYLS1. Autosomal Recessive. Sequencing with copy number analysis. **Exon:** NM_145014:4. **Detection Rate:** East Asian >99%.

Copyright 2021 Myriad Women's Health, Inc. All rights reserved.

Hypophosphatasia - **Gene:** ALPL. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000478:2-12. **Detection Rate:** East Asian >99%.
Isovaleric Acidemia - **Gene:** IVD. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_002225:1-12. **Detection Rate:** East Asian >99%.
Joubert Syndrome 2 - **Gene:** TMEM216. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001173990:1-5. **Detection Rate:** East Asian >99%.
Junctional Epidermolysis Bullosa, LAMA3-related - **Gene:** LAMA3. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000227:1-38. **Detection Rate:** East Asian >99%.
Junctional Epidermolysis Bullosa, LAMC2-related - **Gene:** LAMC2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_005562:1-23. **Detection Rate:** East Asian >99%.
KCNJ11-related Familial Hyperinsulinism - **Gene:** KCNJ11. Autosomal Recessive. Sequencing with copy number analysis. **Exon:** NM_000525:1. **Detection Rate:** East Asian >99%.
Krabbe Disease - **Gene:** GALC. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000153:1-17. **Detection Rate:** East Asian >99%.
LAMA2-related Muscular Dystrophy - **Gene:** LAMA2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000426:1-43,45-65. **Detection Rate:** East Asian >99%.
Leigh Syndrome, French-Canadian Type - **Gene:** LRPPRC. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_133259:1-38. **Detection Rate:** East Asian >99%.
Lipoid Congenital Adrenal Hyperplasia - **Gene:** STAR. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000349:1-7. **Detection Rate:** East Asian >99%.
Lysosomal Acid Lipase Deficiency - **Gene:** LIPA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000235:2-10. **Detection Rate:** East Asian >99%.
Maple Syrup Urine Disease Type Ia - **Gene:** BCKDHA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000709:1-9. **Detection Rate:** East Asian >99%.
Maple Syrup Urine Disease Type Ib - **Gene:** BCKDHB. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_183050:1-10. **Detection Rate:** East Asian >99%.
Maple Syrup Urine Disease Type II - **Gene:** DBT. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001918:1-11. **Detection Rate:** East Asian 96%.
Medium Chain Acyl-CoA Dehydrogenase Deficiency - **Gene:** ACADM. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000016:1-12. **Detection Rate:** East Asian >99%.
Megalencephalic Leukoencephalopathy with Subcortical Cysts - **Gene:** MLC1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_015166 2-12. **Detection Rate:** East Asian >99%.
Metachromatic Leukodystrophy - **Gene:** ARSA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000487:1-8. **Detection Rate:** East Asian >99%.
Methylmalonic Acidemia, cblA Type - **Gene:** MMAA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_172250:2-7. **Detection Rate:** East Asian >99%.
Methylmalonic Acidemia, cblB Type - **Gene:** MMAB. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_052845:1-9. **Detection Rate:** East Asian >99%.
Methylmalonic Aciduria and Homocystinuria, cblC Type - **Gene:** MMACHC. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_015506:1-4. **Detection Rate:** East Asian >99%.
MKS1-related Disorders - **Gene:** MKS1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_017777:1-18. **Detection Rate:** East Asian >99%.
Mucolipidosis III Gamma - **Gene:** GNPTG. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_032520:1-11. **Detection Rate:** East Asian >99%.
Mucolipidosis IV - **Gene:** MCOLN1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_020533:1-14. **Detection Rate:** East Asian >99%.
Mucopolysaccharidosis Type I - **Gene:** IDUA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000203:1-14. **Detection Rate:** East Asian >99%.
Mucopolysaccharidosis Type II - **Gene:** IDS. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000202:1-9. **Detection Rate:** East Asian 88%.
Mucopolysaccharidosis Type IIIA - **Gene:** SGSH. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000199:1-8. **Detection Rate:** East Asian >99%.
Mucopolysaccharidosis Type IIIB - **Gene:** NAGLU. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000263:1-6. **Detection Rate:** East Asian >99%.
Mucopolysaccharidosis Type IIIC - **Gene:** HGSNAT. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_152419:1-18. **Detection Rate:** East Asian >99%.
MUT-related Methylmalonic Acidemia - **Gene:** MUT. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000255:2-13. **Detection Rate:** East Asian >99%.
MYO7A-related Disorders - **Gene:** MYO7A. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000260:2-49. **Detection Rate:** East Asian >99%.
NEB-related Nemaline Myopathy - **Gene:** NEB. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001271208:3-80,117-183. **Detection Rate:** East Asian 92%.
Nephrotic Syndrome, NPHS1-related - **Gene:** NPHS1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_004646:1-29. **Detection Rate:** East Asian >99%.
Nephrotic Syndrome, NPHS2-related - **Gene:** NPHS2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_014625:1-8. **Detection Rate:** East Asian >99%.
Niemann-Pick Disease Type C1 - **Gene:** NPC1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000271:1-25. **Detection Rate:** East Asian >99%.
Niemann-Pick Disease Type C2 - **Gene:** NPC2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_006432:1-5. **Detection Rate:** East Asian >99%.
Niemann-Pick Disease, SMPD1-related - **Gene:** SMPD1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000543:1-6. **Detection Rate:** East Asian >99%.
Nijmegen Breakage Syndrome - **Gene:** NBN. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_002485:1-16. **Detection Rate:** East Asian >99%.
Ornithine Transcarbamylase Deficiency - **Gene:** OTC. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000531:1-10. **Detection Rate:** East Asian 97%.
PCCA-related Propionic Acidemia - **Gene:** PCCA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000282:1-24. **Detection Rate:** East Asian 95%.
PCCB-related Propionic Acidemia - **Gene:** PCCB. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000532:1-15. **Detection Rate:** East Asian >99%.
PCDH15-related Disorders - **Gene:** PCDH15. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_033056:2-33. **Detection Rate:** East Asian 93%.
Pendred Syndrome - **Gene:** SLC26A4. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000441:2-21. **Detection Rate:** East Asian >99%.
Peroxisome Biogenesis Disorder Type 1 - **Gene:** PEX1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000466:1-24. **Detection Rate:** East Asian >99%.
Peroxisome Biogenesis Disorder Type 3 - **Gene:** PEX12. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000286:1-3. **Detection Rate:** East Asian >99%.
Peroxisome Biogenesis Disorder Type 4 - **Gene:** PEX6. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000287:1-17. **Detection Rate:** East Asian 97%.

Copyright 2021 Myriad Women's Health, Inc. All rights reserved.

Peroxisome Biogenesis Disorder Type 5 - **Gene:** PEX2. Autosomal Recessive. Sequencing with copy number analysis. **Exon:** NM_000318:4. **Detection Rate:** East Asian >99%.
Peroxisome Biogenesis Disorder Type 6 - **Gene:** PEX10. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_153818:1-6. **Detection Rate:** East Asian >99%.
Phenylalanine Hydroxylase Deficiency - **Gene:** PAH. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000277:1-13. **Detection Rate:** East Asian >99%.
POMGNT-related Disorders - **Gene:** POMGNT1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_017739:2-22. **Detection Rate:** East Asian 96%.
Pompe Disease - **Gene:** GAA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000152:2-20. **Detection Rate:** East Asian >99%.
PPT1-related Neuronal Ceroid Lipofuscinosis - **Gene:** PPT1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000310:1-9. **Detection Rate:** East Asian >99%.
Primary Carnitine Deficiency - **Gene:** SLC22A5. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_003060:1-10. **Detection Rate:** East Asian >99%.
Primary Hyperoxaluria Type 1 - **Gene:** AGXT. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000030:1-11. **Detection Rate:** East Asian >99%.
Primary Hyperoxaluria Type 2 - **Gene:** GRHPR. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_012203:1-9. **Detection Rate:** East Asian >99%.
Primary Hyperoxaluria Type 3 - **Gene:** HOGA1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_138413:1-7. **Detection Rate:** East Asian >99%.
Pycnodysostosis - **Gene:** CTSK. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000396:2-8. **Detection Rate:** East Asian >99%.
Pyruvate Carboxylase Deficiency - **Gene:** PC. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000920:3-22. **Detection Rate:** East Asian >99%.
Rhizomelic Chondrodysplasia Punctata Type 1 - **Gene:** PEX7. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000288:1-10. **Detection Rate:** East Asian >99%.
RTEL1-related Disorders - **Gene:** RTEL1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_032957:2-35. **Detection Rate:** East Asian >99%.
Sandhoff Disease - **Gene:** HEXB. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000521:1-14. **Detection Rate:** East Asian 99%.
Short-chain Acyl-CoA Dehydrogenase Deficiency - **Gene:** ACADS. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000017:1-10. **Detection Rate:** East Asian >99%.
Sjogren-Larsson Syndrome - **Gene:** ALDH3A2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000382:1-10. **Detection Rate:** East Asian 96%.
SLC26A2-related Disorders - **Gene:** SLC26A2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000112:2-3. **Detection Rate:** East Asian >99%.
Smith-Lemli-Opitz Syndrome - **Gene:** DHCR7. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001360:3-9. **Detection Rate:** East Asian >99%.
Spastic Paraplegia Type 15 - **Gene:** ZFYVE26. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_015346:2-42. **Detection Rate:** East Asian >99%.
Spinal Muscular Atrophy - **Gene:** SMN1. Autosomal Recessive. Spinal muscular atrophy. **Variant (1):** SMN1 copy number. **Detection Rate:** East Asian 93%.
Spondylothoracic Dysostosis - **Gene:** MESP2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001039958:1-2. **Detection Rate:** East Asian >99%.
TGM1-related Autosomal Recessive Congenital Ichthyosis - **Gene:** TGM1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000359 2-15. **Detection Rate:** East Asian >99%.
TPP1-related Neuronal Ceroid Lipofuscinosis - **Gene:** TPP1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000391:1-13. **Detection Rate:** East Asian >99%.
Tyrosine Hydroxylase Deficiency - **Gene:** TH. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_199292:1-14. **Detection Rate:** East Asian >99%.
Tyrosinemia Type I - **Gene:** FAH. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000137:1-14. **Detection Rate:** East Asian >99%.
Tyrosinemia Type II - **Gene:** TAT. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000353:2-12. **Detection Rate:** East Asian >99%.
USH1C-related Disorders - **Gene:** USH1C. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_005709:1-21. **Detection Rate:** East Asian >99%.
USH2A-related Disorders - **Gene:** USH2A. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_206933:2-72. **Detection Rate:** East Asian 94%.
Usher Syndrome Type 3 - **Gene:** CLRN1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_174878:1-3. **Detection Rate:** East Asian >99%.
Very-long-chain Acyl-CoA Dehydrogenase Deficiency - **Gene:** ACADVL. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000018:1-20. **Detection Rate:** East Asian >99%.
Wilson Disease - **Gene:** ATP7B. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000053:1-21. **Detection Rate:** East Asian >99%.
X-linked Adrenoleukodystrophy - **Gene:** ABCD1. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000033:1-6. **Detection Rate:** East Asian 77%.
X-linked Alport Syndrome - **Gene:** COL4A5. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000495:1-51. **Detection Rate:** East Asian 95%.
X-linked Congenital Adrenal Hypoplasia - **Gene:** NR0B1. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000475:1-2. **Detection Rate:** East Asian 99%.
X-linked Juvenile Retinoschisis - **Gene:** RS1. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000330:1-6. **Detection Rate:** East Asian 98%.
X-linked Myotubular Myopathy - **Gene:** MTM1. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000252:2-15. **Detection Rate:** East Asian 99%.
X-linked Severe Combined Immunodeficiency - **Gene:** IL2RG. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000206:1-8. **Detection Rate:** East Asian >99%.
Xeroderma Pigmentosum Group A - **Gene:** XPA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000380:1-6. **Detection Rate:** East Asian >99%.
Xeroderma Pigmentosum Group C - **Gene:** XPC. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_004628:1-16. **Detection Rate:** East Asian 97%.

Copyright 2021 Myriad Women's Health, Inc. All rights reserved.

Risk Calculations

Below are the risk calculations for all conditions tested. Negative results do not rule out the possibility of being a carrier. Residual risk is an estimate of each patient's post-test likelihood of being a carrier, while the reproductive risk represents an estimated likelihood that the patients' future children could inherit each disease. These risks are inherent to all carrier-screening tests, may vary by ethnicity, are predicated on a negative family history, and are present even given a negative test result. Inaccurate reporting of ethnicity may cause errors in risk calculation. In addition, average carrier rates are estimated using incidence or prevalence data from published scientific literature and/or reputable databases, where available, and are incorporated into residual risk calculations for each population/ethnicity. When population-specific data is not available for a condition, average worldwide incidence or prevalence is used. Further, incidence and prevalence data are only collected for the specified phenotypes (which include primarily the classic or severe forms of disease) and may not include alternate or milder disease manifestations associated with the gene. Actual incidence rates, prevalence rates, and carrier rates, and therefore actual residual risks, may be higher or lower than the estimates provided. Carrier rates, incidence/prevalence, and/or residual risks are not provided for some genes with biological or heritable properties that would make these estimates inaccurate. A '†' symbol indicates a positive result. See the full clinical report for interpretation and details. The reproductive risk presented is based on a hypothetical pairing with a partner of the same ethnic group.

Disease	DONOR 14184 Residual Risk	Reproductive Risk
11-beta-hydroxylase-deficient Congenital Adrenal Hyperplasia	1 in 3,300	< 1 in 1,000,000
6-pyruvoyl-tetrahydropterin Synthase Deficiency	1 in 35,000	< 1 in 1,000,000
ABCC8-related Familial Hyperinsulinism	1 in 14,000	< 1 in 1,000,000
Adenosine Deaminase Deficiency	1 in 39,000	< 1 in 1,000,000
Alpha Thalassemia, HBA1/HBA2-related	Alpha globin status: aa/aa.	Not calculated
Alpha-mannosidosis	1 in 35,000	< 1 in 1,000,000
Alpha-sarcoglycanopathy	1 in 34,000	< 1 in 1,000,000
Alstrom Syndrome	< 1 in 50,000	< 1 in 1,000,000
AMT-related Glycine Encephalopathy	1 in 22,000	< 1 in 1,000,000
Andermann Syndrome	< 1 in 50,000	< 1 in 1,000,000
Argininemia	< 1 in 17,000	< 1 in 1,000,000
Argininosuccinic Aciduria	1 in 13,000	< 1 in 1,000,000
Aspartylglucosaminuria	< 1 in 50,000	< 1 in 1,000,000
Ataxia with Vitamin E Deficiency	< 1 in 50,000	< 1 in 1,000,000
Ataxia-telangiectasia	1 in 12,000	< 1 in 1,000,000
ATP7A-related Disorders	< 1 in 1,000,000	1 in 720,000
Autoimmune Polyglandular Syndrome Type 1	1 in 18,000	< 1 in 1,000,000
Autosomal Recessive Osteopetrosis Type 1	1 in 35,000	< 1 in 1,000,000
Autosomal Recessive Polycystic Kidney Disease, PKHD1-related	1 in 8,100	< 1 in 1,000,000
Autosomal Recessive Spastic Ataxia of Charlevoix-Saguenay	< 1 in 44,000	< 1 in 1,000,000
Bardet-Biedl Syndrome, BBS1-related	< 1 in 50,000	< 1 in 1,000,000
Bardet-Biedl Syndrome, BBS10-related	< 1 in 50,000	< 1 in 1,000,000
Bardet-Biedl Syndrome, BBS12-related	< 1 in 50,000	< 1 in 1,000,000
Bardet-Biedl Syndrome, BBS2-related	< 1 in 50,000	< 1 in 1,000,000
BCS1L-related Disorders	< 1 in 50,000	< 1 in 1,000,000
Beta-sarcoglycanopathy	1 in 39,000	< 1 in 1,000,000
Biotinidase Deficiency	1 in 67,000	< 1 in 1,000,000
Bloom Syndrome	< 1 in 50,000	< 1 in 1,000,000
Calpainopathy	1 in 13,000	< 1 in 1,000,000
Canavan Disease	1 in 9,700	< 1 in 1,000,000
Carbamoylphosphate Synthetase I Deficiency	1 in 45,000	< 1 in 1,000,000
Carnitine Palmitoyltransferase IA Deficiency	< 1 in 50,000	< 1 in 1,000,000
Carnitine Palmitoyltransferase II Deficiency	1 in 31,000	< 1 in 1,000,000
Cartilage-hair Hypoplasia	< 1 in 50,000	< 1 in 1,000,000
Cerebrotendinous Xanthomatosis	1 in 11,000	< 1 in 1,000,000
Citrullinemia Type 1	1 in 700	1 in 270,000
CLN3-related Neuronal Ceroid Lipofuscinosis	1 in 13,000	< 1 in 1,000,000
CLN5-related Neuronal Ceroid Lipofuscinosis	< 1 in 50,000	< 1 in 1,000,000
CLN6-related Neuronal Ceroid Lipofuscinosis	< 1 in 50,000	< 1 in 1,000,000
CLN8-related Neuronal Ceroid Lipofuscinosis	< 1 in 50,000	< 1 in 1,000,000
Cohen Syndrome	< 1 in 15,000	< 1 in 1,000,000
COL4A3-related Alport Syndrome	1 in 11,000	< 1 in 1,000,000
COL4A4-related Alport Syndrome	1 in 21,000	< 1 in 1,000,000
Combined Pituitary Hormone Deficiency, PROP1-related	1 in 6,100	< 1 in 1,000,000
Congenital Adrenal Hyperplasia, CYP21A2-related	1 in 550	1 in 140,000

Copyright 2021 Myriad Women's Health, Inc. All rights reserved.

Disease	DONOR 14184 Residual Risk	Reproductive Risk
Congenital Disorder of Glycosylation Type Ia	1 in 16,000	< 1 in 1,000,000
Congenital Disorder of Glycosylation Type Ic	< 1 in 50,000	< 1 in 1,000,000
Congenital Disorder of Glycosylation, MPI-related	< 1 in 50,000	< 1 in 1,000,000
Costeff Optic Atrophy Syndrome	< 1 in 50,000	< 1 in 1,000,000
Cystic Fibrosis	1 in 9,000	< 1 in 1,000,000
Cystinosis	1 in 22,000	< 1 in 1,000,000
D-bifunctional Protein Deficiency	1 in 9,000	< 1 in 1,000,000
Delta-sarcoglycanopathy	< 1 in 40,000	< 1 in 1,000,000
Dihydrolipoamide Dehydrogenase Deficiency	< 1 in 50,000	< 1 in 1,000,000
Dysferlinopathy	1 in 11,000	< 1 in 1,000,000
Dystrophinopathy (Including Duchenne/Becker Muscular Dystrophy)	Not calculated	Not calculated
ERCC6-related Disorders	1 in 26,000	< 1 in 1,000,000
ERCC8-related Disorders	< 1 in 9,800	< 1 in 1,000,000
EVC-related Ellis-van Creveld Syndrome	1 in 7,500	< 1 in 1,000,000
EVC2-related Ellis-van Creveld Syndrome	< 1 in 50,000	< 1 in 1,000,000
Fabry Disease	< 1 in 1,000,000	1 in 80,000
Familial Dysautonomia	< 1 in 50,000	< 1 in 1,000,000
Familial Mediterranean Fever	< 1 in 50,000	< 1 in 1,000,000
Fanconi Anemia Complementation Group A	1 in 3,100	< 1 in 1,000,000
Fanconi Anemia, FANCC-related	< 1 in 50,000	< 1 in 1,000,000
FKRP-related Disorders	< 1 in 50,000	< 1 in 1,000,000
FKTN-related Disorders	1 in 110	1 in 40,000
Free Sialic Acid Storage Disorders	< 1 in 30,000	< 1 in 1,000,000
Galactokinase Deficiency	< 1 in 50,000	< 1 in 1,000,000
Galactosemia	1 in 32,000	< 1 in 1,000,000
Gamma-sarcoglycanopathy	1 in 3,200	< 1 in 1,000,000
Gaucher Disease	1 in 450	1 in 320,000
GJB2-related DFNB1 Nonsyndromic Hearing Loss and Deafness	1 in 3,300	1 in 450,000
GLB1-related Disorders	1 in 19,000	< 1 in 1,000,000
GLDC-related Glycine Encephalopathy	1 in 2,800	< 1 in 1,000,000
Glutaric Acidemia, GCDH-related	1 in 13,000	< 1 in 1,000,000
Glycogen Storage Disease Type Ia	1 in 18,000	< 1 in 1,000,000
Glycogen Storage Disease Type Ib	1 in 35,000	< 1 in 1,000,000
Glycogen Storage Disease Type III	1 in 16,000	< 1 in 1,000,000
GNE Myopathy	< 1 in 50,000	< 1 in 1,000,000
GNPTAB-related Disorders	1 in 17,000	< 1 in 1,000,000
HADHA-related Disorders	1 in 25,000	< 1 in 1,000,000
Hb Beta Chain-related Hemoglobinopathy (Including Beta Thalassemia and Sickle Cell Disease)	1 in 5,000	1 in 990,000
Hereditary Fructose Intolerance	1 in 7,900	< 1 in 1,000,000
Herlitz Junctional Epidermolysis Bullosa, LAMB3-related	< 1 in 50,000	< 1 in 1,000,000
Hexosaminidase A Deficiency (Including Tay-Sachs Disease)	1 in 30,000	< 1 in 1,000,000
HMG-CoA Lyase Deficiency	< 1 in 33,000	< 1 in 1,000,000
Holocarboxylase Synthetase Deficiency	1 in 16,000	< 1 in 1,000,000
Homocystinuria, CBS-related	< 1 in 50,000	< 1 in 1,000,000
Hydrolethalus Syndrome	< 1 in 50,000	< 1 in 1,000,000
Hypophosphatasia	1 in 19,000	< 1 in 1,000,000
Isovaleric Acidemia	1 in 39,000	< 1 in 1,000,000
Joubert Syndrome 2	< 1 in 50,000	< 1 in 1,000,000
Junctional Epidermolysis Bullosa, LAMA3-related	< 1 in 50,000	< 1 in 1,000,000
Junctional Epidermolysis Bullosa, LAMC2-related	< 1 in 50,000	< 1 in 1,000,000
KCNJ11-related Familial Hyperinsulinism	1 in 42,000	< 1 in 1,000,000
Krabbe Disease	1 in 17,000	< 1 in 1,000,000
LAMA2-related Muscular Dystrophy	< 1 in 61,000	< 1 in 1,000,000
Leigh Syndrome, French-Canadian Type	< 1 in 50,000	< 1 in 1,000,000
Lipoid Congenital Adrenal Hyperplasia	1 in 40,000	< 1 in 1,000,000
Lysosomal Acid Lipase Deficiency	1 in 30,000	< 1 in 1,000,000
Maple Syrup Urine Disease Type Ia	1 in 49,000	< 1 in 1,000,000
Maple Syrup Urine Disease Type Ib	1 in 23,000	< 1 in 1,000,000
Maple Syrup Urine Disease Type II	< 1 in 13,000	< 1 in 1,000,000
Medium Chain Acyl-CoA Dehydrogenase Deficiency	1 in 11,000	< 1 in 1,000,000
Megalencephalic Leukoencephalopathy with Subcortical Cysts	< 1 in 50,000	< 1 in 1,000,000

Copyright 2021 Myriad Women's Health, Inc. All rights reserved.

myriad WOMEN'S HEALTH

RESULTS RECIPIENT
SEATTLE SPERM BANK
Attn: Jeffrey Olliffe
NPI: 1306838271
Report Date: 01/22/2021

MALE
DONOR 14184
DOB: [redacted]
Ethnicity: East Asian
Barcode: 11004512825040

FEMALE
N/A

Disease	DONOR 14184 Residual Risk	Reproductive Risk
Metachromatic Leukodystrophy	1 in 16,000	< 1 in 1,000,000
Methylmalonic Acidemia, cblA Type	< 1 in 50,000	< 1 in 1,000,000
Methylmalonic Acidemia, cblB Type	< 1 in 50,000	< 1 in 1,000,000
Methylmalonic Aciduria and Homocystinuria, cblC Type	1 in 33,000	< 1 in 1,000,000
MKS1-related Disorders	< 1 in 50,000	< 1 in 1,000,000
Mucolipidosis III Gamma	< 1 in 50,000	< 1 in 1,000,000
Mucolipidosis IV	< 1 in 50,000	< 1 in 1,000,000
Mucopolysaccharidosis Type I	1 in 16,000	< 1 in 1,000,000
Mucopolysaccharidosis Type II	1 in 390,000	1 in 98,000
Mucopolysaccharidosis Type IIIA	1 in 16,000	< 1 in 1,000,000
Mucopolysaccharidosis Type IIIB	1 in 30,000	< 1 in 1,000,000
Mucopolysaccharidosis Type IIIC	< 1 in 50,000	< 1 in 1,000,000
MUT-related Methylmalonic Acidemia	1 in 11,000	< 1 in 1,000,000
MYO7A-related Disorders	1 in 15,000	< 1 in 1,000,000
NEB-related Nemaline Myopathy	1 in 1,200	1 in 400,000
Nephrotic Syndrome, NPHS1-related	< 1 in 50,000	< 1 in 1,000,000
Nephrotic Syndrome, NPHS2-related	1 in 35,000	< 1 in 1,000,000
Niemann-Pick Disease Type C1	1 in 17,000	< 1 in 1,000,000
Niemann-Pick Disease Type C2	< 1 in 50,000	< 1 in 1,000,000
Niemann-Pick Disease, SMPD1-related	1 in 25,000	< 1 in 1,000,000
Nijmegen Breakage Syndrome	< 1 in 50,000	< 1 in 1,000,000
Ornithine Transcarbamylase Deficiency	< 1 in 1,000,000	1 in 140,000
PCCA-related Propionic Acidemia	1 in 4,200	< 1 in 1,000,000
PCCB-related Propionic Acidemia	1 in 6,500	< 1 in 1,000,000
PCDH15-related Disorders	1 in 3,300	< 1 in 1,000,000
Pendred Syndrome	1 in 6,400	< 1 in 1,000,000
Peroxisome Biogenesis Disorder Type 1	1 in 16,000	< 1 in 1,000,000
Peroxisome Biogenesis Disorder Type 3	< 1 in 50,000	< 1 in 1,000,000
Peroxisome Biogenesis Disorder Type 4	1 in 9,300	< 1 in 1,000,000
Peroxisome Biogenesis Disorder Type 5	< 1 in 71,000	< 1 in 1,000,000
Peroxisome Biogenesis Disorder Type 6	< 1 in 50,000	< 1 in 1,000,000
Phenylalanine Hydroxylase Deficiency	1 in 7,700	< 1 in 1,000,000
POMGNT-related Disorders	< 1 in 12,000	< 1 in 1,000,000
Pompe Disease	1 in 10,000	< 1 in 1,000,000
PPT1-related Neuronal Ceroid Lipofuscinosis	1 in 7,700	< 1 in 1,000,000
Primary Carnitine Deficiency	1 in 10,000	< 1 in 1,000,000
Primary Hyperoxaluria Type 1	1 in 13,000	< 1 in 1,000,000
Primary Hyperoxaluria Type 2	< 1 in 50,000	< 1 in 1,000,000
Primary Hyperoxaluria Type 3	1 in 20,000	< 1 in 1,000,000
Pycnodysostosis	1 in 43,000	< 1 in 1,000,000
Pyruvate Carboxylase Deficiency	1 in 25,000	< 1 in 1,000,000
Rhizomelic Chondrodysplasia Punctata Type 1	1 in 16,000	< 1 in 1,000,000
RTEL1-related Disorders	< 1 in 50,000	< 1 in 1,000,000
Sandhoff Disease	1 in 30,000	< 1 in 1,000,000
Short-chain Acyl-CoA Dehydrogenase Deficiency	1 in 9,700	< 1 in 1,000,000
Sjogren-Larsson Syndrome	< 1 in 12,000	< 1 in 1,000,000
SLC26A2-related Disorders	1 in 16,000	< 1 in 1,000,000
Smith-Lemli-Opitz Syndrome	< 1 in 50,000	< 1 in 1,000,000
Spastic Paraplegia Type 15	< 1 in 50,000	< 1 in 1,000,000
Spinal Muscular Atrophy	Negative for g.27134T>G SNP SMN1: 2 copies 1 in 700	1 in 150,000
Spondylothoracic Dysostosis	< 1 in 50,000	< 1 in 1,000,000
TGM1-related Autosomal Recessive Congenital Ichthyosis	1 in 22,000	< 1 in 1,000,000
TPP1-related Neuronal Ceroid Lipofuscinosis	1 in 30,000	< 1 in 1,000,000
Tyrosine Hydroxylase Deficiency	< 1 in 50,000	< 1 in 1,000,000
Tyrosinemia Type I	1 in 16,000	< 1 in 1,000,000
Tyrosinemia Type II	1 in 25,000	< 1 in 1,000,000
USH1C-related Disorders	1 in 35,000	< 1 in 1,000,000
USH2A-related Disorders	1 in 2,200	< 1 in 1,000,000
Usher Syndrome Type 3	1 in 41,000	< 1 in 1,000,000
Very-long-chain Acyl-CoA Dehydrogenase Deficiency	1 in 12,000	< 1 in 1,000,000

Copyright 2021 Myriad Women's Health, Inc. All rights reserved.

myriad WOMEN'S HEALTH

RESULTS RECIPIENT
SEATTLE SPERM BANK
Attn: Jeffrey Olliffe
NPI: 1306838271
Report Date: 01/22/2021

MALE
DONOR 14184
DOB: [redacted]
Ethnicity: East Asian
Barcode: 11004512825040

FEMALE
N/A

Disease	DONOR 14184 Residual Risk	Reproductive Risk
Wilson Disease	1 in 6,500	< 1 in 1,000,000
X-linked Adrenoleukodystrophy	1 in 170,000	1 in 80,000
X-linked Alport Syndrome	Not calculated	Not calculated
X-linked Congenital Adrenal Hypoplasia	< 1 in 1,000,000	< 1 in 1,000,000
X-linked Juvenile Retinoschisis	< 1 in 1,000,000	1 in 50,000
X-linked Myotubular Myopathy	Not calculated	Not calculated
X-linked Severe Combined Immunodeficiency	< 1 in 1,000,000	1 in 200,000
Xeroderma Pigmentosum Group A	1 in 10,000	< 1 in 1,000,000
Xeroderma Pigmentosum Group C	1 in 7,300	< 1 in 1,000,000

Copyright 2021 Myriad Women's Health, Inc. All rights reserved.

15 Crawford St., STE 100
Needham, MA 02494
(p) 626-350-0537
(f) 626-454-1667
Lab Director: Arash Radfar M.D.
CLIA: 22D0957540

fulgent | INFORM DIAGNOSTICS™ A FULGENT GENETICS COMPANY
Metroplex Pathology Associates
Lakewood Pathology Associates
Cohen Dermatopathology
Therapath Partners

Patient Information:
14184, Donor
DOB: [redacted]
Sex: M
MR#:
Patient#: FT-PT9004637

Partner Information:
Not Tested

Physician:
Kuan, James
ATTN: Martinez,Lupe
San Diego Sperm Bank
4915 25th Avenue NE, Ste 204W
Seattle, WA 98105
Phone: (206) 588-1484

Laboratory:
Fulgent Therapeutics LLC
CAP#: 8042697
CLIA#: 05D2043189
Laboratory Director:
Dr. Amar Jariwala
Report Date: **Mar 20,2025**

Accession:
FT-7380489
Test#: FT-TS15112425
Specimen Type: Saliva
Collected: Feb 22,2025

Accession:
N/A

FINAL RESULTS

No carrier mutations identified

TEST PERFORMED

Single Gene Carrier Screening: ATP8B1
(1 Gene Panel: *ATP8B1*; gene sequencing with deletion and duplication analysis)

INTERPRETATION:

Notes and Recommendations:

- No carrier mutations were identified in the submitted specimen. A negative result does not rule out the possibility of a genetic predisposition nor does it rule out any pathogenic mutations in areas not assessed by this test or in regions that were covered at a level too low to reliably assess. Also, it does not rule out mutations that are of the sort not queried by this test; see Methods and Limitations for more information. A negative result reduces, but does not eliminate, the chance to be a carrier for any condition included in this screen. Please see the supplemental table for details.
- This carrier screening test does not screen for all possible genetic conditions, nor for all possible mutations in every gene tested. This report does not include variants of uncertain significance; only variants classified as pathogenic or likely pathogenic at the time of testing, and considered relevant for reproductive carrier screening, are reported. Please see the gene specific notes for details. Please note that the classification of variants can change over time.
- Patients may wish to discuss any carrier results with blood relatives, as there is an increased chance that they are also carriers. These results should be interpreted in the context of this individual's clinical findings, biochemical profile, and family history.
- Gene specific notes and limitations may be present. See below.
- Genetic counseling is recommended. Available genetic counselors and additional resources can be found at the National Society of Genetic Counselors (NSGC; https://www.nsgc.org)

15 Crawford St., STE 100
Needham, MA 02494
(p) 626-350-0537
(f) 626-454-1667
Lab Director: Arash Radfar M.D.
CLIA: 22D0957540

GENES TESTED:

Custom Beacon Preconception Carrier Screening Panel - Gene

This analysis was run using the Custom Beacon Preconception Carrier Screening Panel gene list. 1 genes were tested with 100.0% of targets sequenced at >20x coverage. For more gene-specific information and assistance with residual risk calculation, see the SUPPLEMENTAL TABLE.

ATP8B1

METHODS:

Genomic DNA was isolated from the submitted specimen indicated above (if cellular material was submitted). DNA was barcoded, and enriched for the coding exons of targeted genes using hybrid capture technology. Prepared DNA libraries were then sequenced using a Next Generation Sequencing technology. Following alignment to the human genome reference sequence (assembly GRCh37), variants were detected in regions of at least 10x coverage. For this specimen, 100.00% and 100.00% of coding regions and splicing junctions of genes listed had been sequenced with coverage of at least 10x and 20x, respectively, by NGS or by Sanger sequencing. The remaining regions did not have 10x coverage, and were not evaluated. Variants were interpreted manually using locus specific databases, literature searches, and other molecular biological principles. To minimize false positive results, any variants that do not meet internal quality standards are confirmed by Sanger sequencing. Variants classified as pathogenic, likely pathogenic, or risk allele which are located in the coding regions and nearby intronic regions (+/- 20bp) of the genes listed above are reported. Variants outside these intervals may be reported but are typically not guaranteed. When a single pathogenic or likely pathogenic variant is identified in a clinically relevant gene with autosomal recessive inheritance, the laboratory will attempt to ensure 100% coverage of coding sequences either through NGS or Sanger sequencing technologies ("fill-in"). All genes listed were evaluated for large deletions and/or duplications. However, single exon deletions or duplications will not be detected in this assay, nor will copy number alterations in regions of genes with significant pseudogenes. Putative deletions or duplications are analyzed using Fulgent Germline proprietary pipeline for this specimen. Bioinformatics: The FPLMv2.0 pipeline was used to analyze this specimen.

LIMITATIONS:

General Limitations

These test results and variant interpretation are based on the proper identification of the submitted specimen, accuracy of any stated familial relationships, and use of the correct human reference sequences at the queried loci. In very rare instances, errors may result due to mix-up or co-mingling of specimens. Positive results do not imply that there are no other contributors, genetic or otherwise, to future pregnancies, and negative results do not rule out the genetic risk to a pregnancy. Official gene names change over time. Fulgent uses the most up to date gene names based on HUGO Gene Nomenclature Committee (https://www.genenames.org) recommendations. If the gene name on report does not match that of ordered gene, please contact the laboratory and details can be provided. Result interpretation is based on the available clinical and family history information for this individual, collected published information, and Alamut annotation available at the time of reporting. This assay is not designed or validated for the detection of low-level mosaicism or somatic mutations. This assay will not detect certain types of genomic aberrations such as translocations, inversions, or repeat expansions other than specified genes. DNA alterations in regulatory regions or deep intronic regions (greater than 20bp from an exon) may not be detected by this test. Unless otherwise indicated, no additional assays have been performed to evaluate genetic changes in this specimen. There are technical limitations on the ability of DNA sequencing to detect small insertions and deletions. Our laboratory uses a sensitive detection algorithm, however these types of alterations are not detected as reliably as single nucleotide variants. Rarely, due to systematic chemical, computational, or human error, DNA variants may be missed. Although next generation sequencing technologies and our bioinformatics analysis significantly reduce the confounding contribution of pseudogene sequences or other highly-homologous sequences, sometimes these may still interfere with the technical ability of the assay to identify pathogenic alterations in both sequencing and deletion/duplication analyses. Deletion/duplication analysis can identify alterations of genomic regions which include one whole gene (buccal swab specimens and whole blood specimens) and are two or more contiguous exons in size (whole blood specimens only); single exon deletions or duplications may occasionally be identified, but are not routinely detected by this test. When novel DNA duplications are identified, it is not possible to discern the genomic location or orientation of the duplicated segment, hence the effect of the duplication cannot be predicted. Where deletions are detected, it is not always possible to determine whether the predicted product will remain in-frame or not. Unless otherwise indicated, deletion/duplication analysis has not been performed in regions that have been sequenced by Sanger.

15 Crawford St., STE 100
Needham, MA 02494
(p) 626-350-0537
(f) 626-454-1667
Lab Director: Arash Radfar M.D.
CLIA: 22D0957540

fulgent | INFORM DIAGNOSTICS™ A FULGENT GENETICS COMPANY
Metroplex Pathology Associates
Lakewood Pathology Associates
Cohen Dermatopathology
Therapath Partners

Gene Specific Notes and Limitations

No gene specific limitations apply to the genes on the tested panel.

SIGNATURE:

Geetu Mendiratta-Vij, PhD, FACMG, CGMBS on **3/20/2025**
Laboratory Director, Fulgent

DISCLAIMER:

This test was developed and its performance characteristics determined by Fulgent Therapeutics LLC CAP #8042697 CLIA #05D2043189; 4399 Santa Anita Ave., El Monte, CA, 91731. It has not been cleared or approved by the FDA. The laboratory is regulated under CLIA as qualified to perform high-complexity testing. This test is used for clinical purposes. It should not be regarded as investigational or for research. Since genetic variation, as well as systematic and technical factors, can affect the accuracy of testing, the results of testing should always be interpreted in the context of clinical and familial data. For assistance with interpretation of these results, healthcare professionals may contact us directly at **626-350-0537** or by email at **info@fulgentgenetics.com**. It is recommended that patients receive appropriate genetic counseling to explain the implications of the test result, including its residual risks, uncertainties and reproductive or medical options.

To view the supplemental table describing the carrier frequencies, detection rates, and residual risks associated with the genes tested on any Beacon panel, please visit the following link:
Beacon Expanded Carrier Screening Supplemental Table

Integrated GENETICS
LabCorp Specialty Testing Group

Client/Sending Facility:
San Diego Sperm Bank

8950 Villa La Jolla Dr Ste B214
La Jolla, CA 92037
Ph: (858)732-8500
CAB-50

LCLS Specimen Number:	**346-253-0329-0**	Account Number:	04087470
Patient Name:	**14184, DONOR**	Ordering Physician:	**J OLLIFFE**
Date of Birth:	[redacted]	Specimen Type:	**BLOOD**
Gender:	M	Client Reference:	
Patient ID:		Date Collected:	12/11/2020
Lab Number:	YU20-93104 L	Date Received:	12/13/2020
Indications:	NOT GIVEN	Date Reported:	**12/24/2020**

Test: **Chromosome, Blood, Routine**

Cells Counted:	20	Cells Karyotyped:	2
Cells Analyzed:	20	Band Resolution:	500

CYTOGENETIC RESULT: 46,XY

INTERPRETATION: NORMAL MALE KARYOTYPE

Cytogenetic analysis of PHA stimulated cultures has revealed a MALE karyotype with an apparently normal GTG banding pattern in all cells observed.

This result does not exclude the possibility of subtle rearrangements below the resolution of cytogenetics or congenital anomalies due to other etiologies.

Chromosome analysis performed at Virtual Scientific, Inc-NE, CLIA# 26D2050153. 2835 S 27th Street, Lincoln, NE 68502. Laboratory Director Christine F Stephenson, M.D.

Integrated GENETICS
LabCorp Specialty Testing Group

Client/Sending Facility:
San Diego Sperm Bank

8950 Villa La Jolla Dr Ste B214
La Jolla, CA 92037
Ph: (858)732-8500
CAB-50

LCLS Specimen Number:	**346-253-0329-0**	Account Number:	04087470
Patient Name:	**14184, DONOR**	Ordering Physician:	**J OLLIFFE**
Date of Birth:	[redacted]	Specimen Type:	**BLOOD**
Gender:	M	Client Reference:	
Patient ID:		Date Collected:	12/11/2020
Lab Number:	YU20-93104 L	Date Received:	12/13/2020



Integrated GENETICS
LabCorp Specialty Testing Group

Client/Sending Facility:
San Diego Sperm Bank

8950 Villa La Jolla Dr Ste B214
La Jolla, CA 92037
Ph: (858)732-8500
CAB-50

LCLS Specimen Number:	**346-253-0329-0**	Account Number:	04087470
Patient Name:	**14184, DONOR**	Ordering Physician:	**J OLLIFFE**
Date of Birth:	[redacted]	Specimen Type:	**BLOOD**
Gender:	M	Client Reference:	
Patient ID:		Date Collected:	12/11/2020
Lab Number:	YU20-93104 L	Date Received:	12/13/2020

Venkateswara R Potluri, PhD

Anjen Chenn, M.D., Ph.D.
Medical Director
Stuart Schwartz, PhD
National Director of Cytogenetics

Technical component performed by Laboratory Corporation of America Holdings,
1904 TW Alexander Drive , RTP , NC , 27709-0153 (800) 345-4363
Professional component performed by LabCorp, CLIA 45D0674994. 7207 North Gessner Rd., Houston, TX 77040. Laboratory Director, Venkateswara R Potluri PhD.
Integrated Genetics is a brand used by Esoterix Genetic Laboratories, LLC, a wholly-owned subsidiary of Laboratory Corporation of America Holdings.

This document contains private and confidential health information **protected by state and federal law.**

LabCorp

Patient Report

Specimen ID: 346-253-0334-0
Control ID: B0109869531

Acct #: 04087470 **Phone:** (858) 732-8500 **Rte:** 00

14184, DONOR

San Diego Sperm Bank
8950 Villa La Jolla Dr Ste B214
La Jolla CA 92037

Patient Details
DOB: [redacted]
Age(y/m/d): [redacted]
Gender: M
Patient ID:

Specimen Details
Date collected: 12/11/2020 1452 Local
Date received: 12/12/2020
Date entered: 12/11/2020
Date reported: 12/15/2020 1105 ET

Physician Details
Ordering: J OLLIFFE
Referring:
ID:
NPI: 1306838271

General Comments & Additional Information
Alternate Control Number: B0109869531
Total Volume: Not Provided

Alternate Patient ID: Not Provided
Fasting: No

Ordered Items
LP+12AC+CBC/D/Plt+UA+Rh+ABO...

TESTS	RESULT	FLAG	UNITS	REFERENCE INTERVAL	LAB
LP+12AC+CBC/D/Plt+UA+Rh+ABO...					
Glucose	94		mg/dL	65-99	01
Uric Acid	6.3		mg/dL	3.8-8.4	01
	Therapeutic target for gout patients: <6.0				
	Please note reference interval change				
BUN	15		mg/dL	6-20	01
Creatinine	0.99		mg/dL	0.76-1.27	01
eGFR If NonAfricn Am	106		mL/min/1.73	>59	
eGFR If Africn Am	123		mL/min/1.73	>59	
Calcium	9.8		mg/dL	8.7-10.2	01
Protein, Total	7.9		g/dL	6.0-8.5	01
Albumin	5.1		g/dL	4.1-5.2	01
Bilirubin, Total	0.6		mg/dL	0.0-1.2	01
Alkaline Phosphatase	80		IU/L	39-117	01
LDH	145		IU/L	121-224	01
AST (SGOT)	20		IU/L	0-40	01
ALT (SGPT)	14		IU/L	0-44	01
Cholesterol, Total	140		mg/dL	100-199	01
Triglycerides	47		mg/dL	0-149	01
HDL Cholesterol	52		mg/dL	>39	01
LDL Chol Calc (NIH)	77		mg/dL	0-99	
LDL/HDL Ratio	1.5		ratio	0.0-3.6	
Please Note:					01

LDL/HDL Ratio

	Men	Women
1/2 Avg.Risk	1.0	1.5
Avg.Risk	3.6	3.2
2X Avg.Risk	6.2	5.0
3X Avg.Risk	8.0	6.1

Hemoglobin Fractionation 15 Feb 21 02
02

Date Issued: 12/17/20 1016 ET

FINAL REPORT

This document contains private and confidential health information protected by state and federal law.
If you have received this document in error, please call 858-668-3700

© 1995-2020 Laboratory Corporation of America® Holdings
All Rights Reserved - Enterprise Report Version: 1.00

Patient: 14184, DONOR
DOB: [redacted] Patient ID: Control ID: B0109869531
Specimen ID: 346-253-0334-0
Date collected: 12/11/2020 1452 Local

TESTS	RESULT	FLAG	UNITS	REFERENCE INTERVAL	LAB
Hgb F [A]	0.0		%	0.0-2.0	02
Hgb A [A]	97.9		%	96.4-98.8	02
Hgb S [A]	0.0		%	0.0	02
Hgb C [A]	0.0		%	0.0	02
Hgb A2 [A]	2.1		%	1.8-3.2	02
Hgb Variant [A]	0.0		%	0.0	02
Interpretation					
Normal adult hemoglobin present.					02
ABO Grouping	B				01
Rh Factor	Positive				01
Please note: Prior records for this patient's ABO / Rh type are not available for additional verification.					
					01
CBC, Platelet Ct, and Diff					01
WBC	7.2		x10E3/uL	3.4-10.8	01
RBC	5.58		x10E6/uL	4.14-5.80	01
Hemoglobin	16.3		g/dL	13.0-17.7	01
Hematocrit	48.3		%	37.5-51.0	01
MCV	87		fL	79-97	01
MCH	29.2		pg	26.6-33.0	01
MCHC	33.7		g/dL	31.5-35.7	01
RDW	11.8		%	11.6-15.4	01
Platelets	371		x10E3/uL	150-450	01
Neutrophils	57		%	Not Estab.	01
Lymphs	32		%	Not Estab.	01
Monocytes	8		%	Not Estab.	01
Eos	2		%	Not Estab.	01
Basos	1		%	Not Estab.	01
Neutrophils (Absolute)	4.2		x10E3/uL	1.4-7.0	01
Lymphs (Absolute)	2.3		x10E3/uL	0.7-3.1	01
Monocytes(Absolute)	0.6		x10E3/uL	0.1-0.9	01
Eos (Absolute)	0.1		x10E3/uL	0.0-0.4	01
Baso (Absolute)	0.1		x10E3/uL	0.0-0.2	01
Immature Granulocytes	0		%	Not Estab.	01
Immature Grans (Abs)	0.0		x10E3/uL	0.0-0.1	01
					01
Urinalysis Gross Exam					01
Specific Gravity	1.006			1.005-1.030	01
pH	7.5			5.0-7.5	01
Urine-Color	Yellow			Yellow	01
Appearance	Clear			Clear	01
WBC Esterase	Negative			Negative	01
Protein	Negative			Negative/Trace	01

This document contains private and confidential health information protected by state and federal law.
If you have received this document in error, please call 858-668-3700

© 1995-2020 Laboratory Corporation of America® Holdings
All Rights Reserved - Enterprise Report Version: 1.00

LabCorp

Patient Report

Patient: 14184 DONOR
DOB: [redacted]
Patient ID:
Control ID: B0109869531
Specimen ID: 346-253-0334-
Date collected: 12/11/2020 1452 Loca

TESTS	RESULT	FLAG	UNITS	REFERENCE INTERVAL	LAB
Glucose	Negative			Negative	01
Ketones	Negative			Negative	01
Occult Blood	Negative			Negative	01
Bilirubin	Negative			Negative	01
Urobilinogen, Semi-Qn	0.2		mg/dL	0.2-1.0	01
Nitrite, Urine	Negative			Negative	01
Microscopic Examination					
Microscopic follows if indicated.					

Comments: 01

A Results of this test are labeled for research purposes only by the assay's manufacturer. The performance characteristics of this assay have not been established by the manufacturer. The result should not be used for treatment or for diagnostic purposes without confirmation of the diagnosis by another medically established diagnostic product or procedure. The performance characteristics were determined by LabCorp.

01	SO	LabCorp San Diego 13112 Evening Creek Dr So Ste 200, San Diego, CA 92128-4108	Dir: Jenny Galloway, MD
02	BN	LabCorp Burlington 1447 York Court, Burlington, NC 27215-3361	Dir: Sanjai Nagendra, MD

For inquiries, the physician may contact **Branch: 800-859-6046 Lab: 858-668-3700**

15 Feb 21

Date Issued: 12/17/20 1016 ET

FINAL REPORT

This document contains private and confidential health information protected by state and federal law.
If you have received this document in error, please call 858-668-3700

© 1995-2020 Laboratory Corporation of America® Holdings
All Rights Reserved - Enterprise Report Version: 1.00