RESULTS RECIPIENT
SEATTLE SPERM BANK
Attn: Jeffrey Olliffe
4915 25th Ave NE Ste 204W
Seattle, WA 98105
Phone: (206) 588-1484
Fax: (206) 466-4696
NPI: 1306838271
Report Date: 11/06/2021

MALE
DONOR 14289
DOB:
Ethnicity: East Asian
Sample Type: EDTA Blood
Date of Collection: 10/27/2021
Date Received: 10/29/2021
Date Tested: 11/06/2021
Barcode: 11004513001910
Accession ID: CSLY6RGEUCLHQRY
Indication: Egg or sperm donor

FEMALE
N/A

Foresight® Carrier Screen

NEGATIVE

ABOUT THIS TEST

The **Myriad Foresight Carrier Screen** utilizes sequencing, maximizing coverage across all DNA regions tested, to help you learn about your chance to have a child with a genetic disease.

RESULTS SUMMARY

Risk Details	DONOR 14289	Partner
Panel Information	Foresight Carrier Screen Universal Panel Fundamental Plus Panel Fundamental Panel **(175 conditions tested)**	N/A
All conditions tested A complete list of all conditions tested can be found on page 5.	⊖ **NEGATIVE** No disease-causing mutations were detected.	N/A

CLINICAL NOTES

- None

NEXT STEPS

- If necessary, patients can discuss residual risks with their physician or a genetic counselor.

Copyright 2021 Myriad Women's Health, Inc. All rights reserved.

Methods and Limitations

DONOR 14289 [Foresight Carrier Screen]: Sequencing with copy number analysis, spinal muscular atrophy, analysis of homologous regions, and alpha thalassemia (HBA1/HBA2) sequencing with targeted copy number analysis (Assay(s): DTS v3.2).

Sequencing with copy number analysis

High-throughput sequencing and read-depth-based copy number analysis are used to analyze the genes listed in the Conditions Tested section of the report. Except where otherwise noted, the region of interest (ROI) comprises the indicated coding regions and 20 non-coding bases flanking each region. In a minority of cases where genomic features (e.g., long homopolymers) compromise calling fidelity, the affected non-coding bases are excluded from the ROI. The ROI is sequenced to a minimum acceptable read depth, and the sequences are compared to a reference genomic sequence (Genome Reference Consortium Human Build 37 [GRCh37]/hg19). On average, 99% of all bases in the ROI are sequenced at a read depth that is greater than the minimum read depth. Sequence variants may not be detected in areas of lower sequence coverage. Insertions and deletions may not be detected as accurately as single-nucleotide variants. Select genes or regions for which pseudogenes or other regions of homology impede reliable variant detection may be assayed using alternate technology, or they may be excluded from the ROI. *CFTR* and *DMD* testing includes analysis for exon-level deletions and duplications with an average sensitivity of ~99%. Only exon-level deletions are assayed for other genes on the panel and such deletions are detected with a sensitivity of ≥75%. Selected founder deletions may be detected at slightly higher sensitivity. Affected exons and/or breakpoints of copy number variants are estimated from junction reads, where available, or using the positions of affected probes. Only exons known to be included in the region affected by a copy number variant are provided in the variant nomenclature. In some cases, the copy number variant may be larger or smaller than indicated. If *GJB2* is tested, large upstream deletions involving the *GJB6* and/or *CRYL1* genes that may affect the expression of *GJB2* are also analyzed.

Spinal muscular atrophy

Targeted copy number analysis via high-throughput sequencing is used to determine the copy number of exon 7 of the *SMN1* gene. Other genetic variants may interfere with this analysis. Some individuals with two copies of *SMN1* are "silent" carriers with both *SMN1* genes on one chromosome and no copies of the gene on the other chromosome. This is more likely in individuals who have two copies of the *SMN1* gene and are positive for the g.27134T>G single-nucleotide polymorphism (SNP) (PMID: 9199562, 23788250, and 28676062), which affects the reported residual risk; Ashkenazi Jewish or Asian patients with this genotype have a high post-test likelihood of being carriers for SMA and are reported as carriers. The g.27134T>G SNP is only reported in individuals who have two copies of *SMN1*.

Analysis of homologous regions

A combination of high-throughput sequencing, read-depth-based copy number analysis, and targeted genotyping is used to determine the number of functional gene copies and/or the presence of selected loss-of-function variants in certain genes that have homology to other genomic regions. The precise breakpoints of large deletions in these genes cannot be determined but are instead estimated from copy number analysis. Pseudogenes may interfere with this analysis, especially when many pseudogene copies are present.

If *CYP21A2* is tested, patients who have one or more additional copies of the *CYP21A2* gene and a pathogenic variant may or may not be a carrier of 21-hydroxylase deficient CAH, depending on the chromosomal location of the variants (phase). Benign *CYP21A2* gene duplications and/or triplications will only be reported in this context. Some individuals with two functional *CYP21A2* gene copies may be "silent" carriers, with two gene copies resulting from a duplication on one chromosome and a gene deletion on the other chromosome. This and other similar rare carrier states, where complementary changes exist between the chromosomes, may not be detected by the assay. Given that the true incidence of non-classic CAH is unknown, the residual carrier and reproductive risk numbers on the report are based only on the published incidence for classic CAH. However, the published prevalence of non-classic CAH is highest in individuals of Ashkenazi Jewish, Hispanic, Italian, and Yugoslav descent. Therefore, the residual and reproductive risks are likely an underestimate for CAH, especially in the aforementioned populations, as they do not account for non-classic CAH.

Copyright 2021 Myriad Women's Health, Inc. All rights reserved.

Alpha thalassemia (HBA1/HBA2) sequencing with targeted copy number analysis

High-throughput sequencing and read-depth-based copy number analysis are used to identify sequence variation and functional gene copies within the region of interest (ROI) of *HBA1* and *HBA2*, which includes the listed exons plus 20 intronic flanking bases. In a minority of cases where genomic features (e.g , long homopolymers) compromise calling fidelity, the affected intronic bases are not included in the ROI. The ROI is sequenced to a minimum acceptable read depth, and the sequences are compared to a reference genomic sequence (Genome Reference Consortium Human Build 37 [GRCh37]/hg19). On average, 99% of all bases in the ROI are sequenced at a read depth that is greater than the minimum read depth. Sequence variants may not be detected in areas of lower sequence coverage. Insertions and deletions may not be detected as accurately as single-nucleotide variants. For large deletions or duplications in these genes, the precise breakpoints cannot be determined but are instead estimated from copy number analysis. This assay has been validated to detect up to two additional copies of each alpha globin gene. In rare instances where assay results suggest greater than two additional copies are present, this will be noted but the specific number of gene copies observed will not be provided.

Extensive sequence homology exists between *HBA1* and *HBA2*. This sequence homology can prevent certain variants from being localized to one gene over the other. In these instances, variant nomenclature will be provided for both genes. If follow-up testing is indicated for patients with the nomenclature provided for both genes, both *HBA1* and *HBA2* should be tested. Some individuals with four functional alpha globin gene copies may be "silent" carriers, with three gene copies resulting from triplication on one chromosome and a single gene deletion on the other chromosome. This and other similar rare carrier states, where complementary changes exist between the chromosomes, may not be detected by the assay.

Interpretation of reported variants

The classification and interpretation of all variants identified in this assay reflects the current state of Myriad's scientific understanding at the time this report was issued. Variants are classified according to internally defined criteria, which are compatible with the ACMG Standards and Guidelines for the Interpretation of Sequence Variants (PMID: 25741868). Variants that have been determined by Myriad to be disease-causing or likely disease-causing (i.e. pathogenic or likely pathogenic) are reported. Benign variants, variants of uncertain clinical significance (VUS), and variants not directly associated with the specified disease phenotype(s) are not reported. Variant classification and interpretation may change for a variety of reasons, including but not limited to, improvements to classification techniques, availability of additional scientific information, and observation of a variant in more patients. f the classification of one or more variants identified in this patient changes, an updated report reflecting the new classification generally will not be issued. If an updated report is issued, the variants reported may change based on their current classification. This can include changes to the variants displayed in gene specific 'variants tested' sections. Healthcare providers may contact Myriad directly to request updated variant classification information specific to this test result.

Limitations

The MWH Foresight Carrier Screen is designed to detect and report germline (constitutional) alterations. Mosaic (somatic) variation may not be detected, and if it is detected, it may not be reported. If more than one variant is detected in a gene, additional studies may be necessary to determine if those variants lie on the same chromosome or different chromosomes (phase). This test is not designed to detect sex-chromosome copy number variations. If present, sex-chromosome abnormalities may significantly reduce test sensitivity for X-linked conditions. Variant interpretation and residual and reproductive risk estimations assume a normal karyotype and may be different for individuals with abnormal karyotypes. The test does not fully address all inherited forms of intellectual disability, birth defects, or heritable diseases. Furthermore, not all forms of genetic variation are detected by this assay (i.e., duplications [except in specified genes], chromosomal rearrangements, structural abnormalities, etc.). Additional testing may be appropriate for some individuals. Pseudogenes and other regions of homology may interfere with this analysis. In an unknown number of cases, other genetic variation may interfere with variant detection. Rare carrier states where complementary changes exist between the chromosomes may not be detected by the assay. Other possible sources of diagnostic error include sample mix-up, trace contamination, bone marrow transplantation, blood transfusions, and technical or analytical errors.

Detection rates are determined using published scientific literature and/or reputable databases, when available, to estimate the fraction of disease alleles, weighted by frequency, that the methodology is predicted to be able or unable to detect. Detection rates are approximate and only account for analytical sensitivity. Certain variants that have been previously described in the literature may not be reported, if there is insufficient evidence for pathogenicity. Detection rates do not account for the disease specific rates of *de novo* variation.

This test was developed, and its performance characteristics determined by, Myriad Women's Health, Inc. It has not been cleared or approved by the US Food and Drug Administration (FDA). The FDA does not require this test to go through premarket review. This test is used for clinical purposes. It should not be regarded as investigational or for research. This laboratory is certified under the Clinical Laboratory Improvement Amendments of 1988 (CLIA) as qualified to perform high-complexity clinical testing. These results are adjunctive to the ordering physician's evaluation. CLIA Number: #05D1102604.

Copyright 2021 Myriad Women's Health, Inc. All rights reserved.

Incidental Findings

Unless otherwise indicated, these results and interpretations are limited to the specific disease panel(s) requested by the ordering healthcare provider. In some cases, standard data analyses may identify genetic findings beyond the region(s) of interest specified by the test, and such findings may not be reported. These findings may include genomic abnormalities with major, minor, or no, clinical significance.


If you have questions or would like more information about any of the test methods or limitations, please contact (888) 268-6795.

Resources

GENOME CONNECT | http://www.genomeconnect.org

Patients can share their reports using research registries such as Genome Connect, an online research registry building a genetics and health knowledge base. Genome Connect provides patients, physicians, and researchers an opportunity to share genetic information to support the study of the impact of genetic variation on health conditions.

SENIOR LABORATORY DIRECTOR



Karla R. Bowles, PhD, FACMG, CGMB

Report content approved by Karla Bowles, PhD, FACMG, CGMB on Nov 6, 2021

Copyright 2021 Myriad Women's Health, Inc. All rights reserved.

Conditions Tested

6-pyruvoyl-tetrahydropterin Synthase Deficiency - **Gene:** PTS. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000317:1-6. **Detection Rate:** East Asian >99%.

Adenosine Deaminase Deficiency - **Gene:** ADA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000022:1-12. **Detection Rate:** East Asian 98%.

Alpha Thalassemia, HBA1/HBA2-related - **Genes:** HBA1, HBA2. Autosomal Recessive. Alpha thalassemia (HBA1/HBA2) sequencing with targeted copy number analysis. **Exons:** NM_000517:1-3; NM_000558:1-3. **Variants (16):** -(alpha)20.5, --BRIT, --MEDI, --MEDII, --SEA, --THAI or --FIL, -alpha3.7, -alpha4.2, HBA1+HBA2 deletion, Hb Constant Spring, Poly(A) AATAAA>AATA--, Poly(A) AATAAA>AATAAG, Poly(A) AATAAA>AATGAA, anti3.7, anti4.2, del HS-40. **Detection Rate:** East Asian >99%.

Alpha-mannosidosis - **Gene:** MAN2B1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000528:1-23. **Detection Rate:** East Asian >99%.

Alpha-sarcoglycanopathy - **Gene:** SGCA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000023:1-9. **Detection Rate:** East Asian >99%.

Alstrom Syndrome - **Gene:** ALMS1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_015120:1-23. **Detection Rate:** East Asian >99%.

Andermann Syndrome - **Gene:** SLC12A6. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_133647:1-25. **Detection Rate:** East Asian >99%.

Argininemia - **Gene:** ARG1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000045:1-8. **Detection Rate:** East Asian 97%.

Argininosuccinic Aciduria - **Gene:** ASL. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001024943:1-16. **Detection Rate:** East Asian >99%.

Aspartylglucosaminuria - **Gene:** AGA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000027:1-9. **Detection Rate:** East Asian >99%.

Ataxia with Vitamin E Deficiency - **Gene:** TTPA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000370:1-5. **Detection Rate:** East Asian >99%.

Ataxia-telangiectasia - **Gene:** ATM. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000051:2-63. **Detection Rate:** East Asian 96%.

ATP7A-related Disorders - **Gene:** ATP7A. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000052:2-23. **Detection Rate:** East Asian 90%.

Autoimmune Polyglandular Syndrome Type 1 - **Gene:** AIRE. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000383:1-14. **Detection Rate:** East Asian >99%.

Autosomal Recessive Osteopetrosis Type 1 - **Gene:** TCIRG1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_006019:2-20. **Detection Rate:** East Asian 96%.

Autosomal Recessive Polycystic Kidney Disease, PKHD1-related - **Gene:** PKHD1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_138694 2-67. **Detection Rate:** East Asian >99%.

Autosomal Recessive Spastic Ataxia of Charlevoix-Saguenay - **Gene:** SACS. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_014363 2-10. **Detection Rate:** East Asian 99%.

Bardet-Biedl Syndrome, BBS1-related - **Gene:** BBS1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_024649:1-17. **Detection Rate:** East Asian >99%.

Bardet-Biedl Syndrome, BBS10-related - **Gene:** BBS10. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_024685:1-2. **Detection Rate:** East Asian >99%.

Bardet-Biedl Syndrome, BBS12-related - **Gene:** BBS12. Autosomal Recessive. Sequencing with copy number analysis. **Exon:** NM_152618:2. **Detection Rate:** East Asian >99%.

Bardet-Biedl Syndrome, BBS2-related - **Gene:** BBS2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_031885:1-17. **Detection Rate:** East Asian >99%.

BCS1L-related Disorders - **Gene:** BCS1L. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_004328:3-9. **Detection Rate:** East Asian >99%.

Beta-sarcoglycanopathy - **Gene:** SGCB. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000232:1-6. **Detection Rate:** East Asian >99%.

Biotinidase Deficiency - **Gene:** BTD. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000060:1-4. **Detection Rate:** East Asian >99%.

Bloom Syndrome - **Gene:** BLM. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000057:2-22. **Detection Rate:** East Asian >99%.

Calpainopathy - **Gene:** CAPN3. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000070:1-24. **Detection Rate:** East Asian 99%.

Canavan Disease - **Gene:** ASPA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000049:1-6. **Detection Rate:** East Asian 98%.

Carbamoylphosphate Synthetase I Deficiency - **Gene:** CPS1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001875:1-38. **Detection Rate:** East Asian >99%.

Carnitine Palmitoyltransferase IA Deficiency - **Gene:** CPT1A. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001876:2-19. **Detection Rate:** East Asian >99%.

Carnitine Palmitoyltransferase II Deficiency - **Gene:** CPT2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000098:1-5. **Detection Rate:** East Asian >99%.

Cartilage-hair Hypoplasia - **Gene:** RMRP. Autosomal Recessive. Sequencing with copy number analysis. **Exon:** NR_003051:1. **Detection Rate:** East Asian >99%.

Cerebrotendinous Xanthomatosis - **Gene:** CYP27A1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000784:1-9. **Detection Rate:** East Asian >99%.

Citrullinemia Type 1 - **Gene:** ASS1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000050:3-16. **Detection Rate:** East Asian 86%.

CLN3-related Neuronal Ceroid Lipofuscinosis - **Gene:** CLN3. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001042432 2-16. **Detection Rate:** East Asian >99%.

CLN5-related Neuronal Ceroid Lipofuscinosis - **Gene:** CLN5. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_006493:1-4. **Detection Rate:** East Asian >99%.

CLN8-related Neuronal Ceroid Lipofuscinosis - **Gene:** CLN8. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_018941:2-3. **Detection Rate:** East Asian >99%.

Cohen Syndrome - **Gene:** VPS13B. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_017890:2-62. **Detection Rate:** East Asian 97%.

COL4A3-related Alport Syndrome - **Gene:** COL4A3. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000091:1-52. **Detection Rate:** East Asian 94%.

COL4A4-related Alport Syndrome - **Gene:** COL4A4. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000092:2-48. **Detection Rate:** East Asian >99%.

Combined Pituitary Hormone Deficiency, PROP1-related - **Gene:** PROP1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_006261:1-3. **Detection Rate:** East Asian >99%.

Congenital Adrenal Hyperplasia, CYP11B1-related - **Gene:** CYP11B1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000497:1-9. **Detection Rate:** East Asian 97%.

Congenital Adrenal Hyperplasia, CYP21A2-related - **Gene:** CYP21A2. Autosomal Recessive. Analysis of homologous regions. **Variants (13):** CYP21A2 deletion, CYP21A2 duplication, CYP21A2 triplication, G111Vfs*21, I173N, L308Ffs*6, P31L, Q319*, Q319*+CYP21A2dup, R357W, V282L, [I237N;V238E;M240K], c.293-13C>G. **Detection Rate:** East Asian 88%.

Congenital Disorder of Glycosylation Type Ia - **Gene:** PMM2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000303:1-8. **Detection Rate:** East Asian >99%.

Copyright 2021 Myriad Women's Health, Inc. All rights reserved.

Congenital Disorder of Glycosylation Type Ic - **Gene:** ALG6. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_013339:2-15. **Detection Rate:** East Asian >99%.
Congenital Disorder of Glycosylation, MPI-related - **Gene:** MPI. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_002435:1-8. **Detection Rate:** East Asian >99%.
Costeff Optic Atrophy Syndrome - **Gene:** OPA3. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_025136:1-2. **Detection Rate:** East Asian >99%.
Cystic Fibrosis - **Gene:** CFTR. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000492:1-27. IVS8-5T allele analysis is only reported in the presence of the R117H mutation. **Detection Rate:** East Asian >99%.
Cystinosis - **Gene:** CTNS. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_004937:3-12. **Detection Rate:** East Asian >99%.
D-bifunctional Protein Deficiency - **Gene:** HSD17B4. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000414:1-24. **Detection Rate:** East Asian 98%.
Delta-sarcoglycanopathy - **Gene:** SGCD. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000337:2-9. **Detection Rate:** East Asian 96%.
Dihydrolipoamide Dehydrogenase Deficiency - **Gene:** DLD. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000108:1-14. **Detection Rate:** East Asian >99%.
Dysferlinopathy - **Gene:** DYSF. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_003494:1-55. **Detection Rate:** East Asian 98%.
Dystrophinopathy (Including Duchenne/Becker Muscular Dystrophy) - **Gene:** DMD. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_004006:1-79. **Detection Rate:** East Asian 99%.
ERCC6-related Disorders - **Gene:** ERCC6. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000124:2-21. **Detection Rate:** East Asian 96%.
ERCC8-related Disorders - **Gene:** ERCC8. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000082:1-12. **Detection Rate:** East Asian 78%.
EVC-related Ellis-van Creveld Syndrome - **Gene:** EVC. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_153717:1-21. **Detection Rate:** East Asian 96%.
EVC2-related Ellis-van Creveld Syndrome - **Gene:** EVC2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_147127:1-22. **Detection Rate:** East Asian 98%.
Fabry Disease - **Gene:** GLA. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000169:1-7. **Detection Rate:** East Asian 98%.
Familial Dysautonomia - **Gene:** ELP1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_003640:2-37. **Detection Rate:** East Asian >99%.
Familial Hyperinsulinism, ABCC8-related - **Gene:** ABCC8. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000352:1-39. **Detection Rate:** East Asian >99%.
Familial Hyperinsulinism, KCNJ11-related - **Gene:** KCNJ11. Autosomal Recessive. Sequencing with copy number analysis. **Exon:** NM_000525:1. **Detection Rate:** East Asian >99%.
Familial Mediterranean Fever - **Gene:** MEFV. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000243:1-10. **Detection Rate:** East Asian >99%.
Fanconi Anemia Complementation Group A - **Gene:** FANCA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000135:1-43. **Detection Rate:** East Asian 92%.
Fanconi Anemia, FANCC-related - **Gene:** FANCC. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000136:2-15. **Detection Rate:** East Asian >99%.
FKRP-related Disorders - **Gene:** FKRP. Autosomal Recessive. Sequencing with copy number analysis. **Exon:** NM_024301:4. **Detection Rate:** East Asian >99%.
FKTN-related Disorders - **Gene:** FKTN. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001079802:3-11. **Detection Rate:** East Asian 10%.
Free Sialic Acid Storage Disorders - **Gene:** SLC17A5. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_012434:1-11. **Detection Rate:** East Asian 98%.
Galactokinase Deficiency - **Gene:** GALK1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000154:1-8. **Detection Rate:** East Asian >99%.
Galactosemia - **Gene:** GALT. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000155:1-11. **Detection Rate:** East Asian >99%.
Gamma-sarcoglycanopathy - **Gene:** SGCG. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000231:2-8. **Detection Rate:** East Asian 87%.
Gaucher Disease - **Gene:** GBA. Autosomal Recessive. Analysis of homologous regions. **Variants (10):** D409V, D448H, IVS2+1G>A, L444P, N370S, R463C, R463H, R496H, V394L, p.L29Afs*18. **Detection Rate:** East Asian 60%.
GJB2-related DFNB1 Nonsyndromic Hearing Loss and Deafness - **Gene:** GJB2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_004004:1-2. **Detection Rate:** East Asian >99%.
GLB1-related Disorders - **Gene:** GLB1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000404:1-16. **Detection Rate:** East Asian >99%.
Glutaric Acidemia, GCDH-related - **Gene:** GCDH. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000159:2-12. **Detection Rate:** East Asian >99%.
Glycine Encephalopathy, AMT-related - **Gene:** AMT. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000481:1-9. **Detection Rate:** East Asian >99%.
Glycine Encephalopathy, GLDC-related - **Gene:** GLDC. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000170:1-25. **Detection Rate:** East Asian 94%.
Glycogen Storage Disease Type Ia - **Gene:** G6PC1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000151:1-5. **Detection Rate:** East Asian 98%.
Glycogen Storage Disease Type Ib - **Gene:** SLC37A4. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001164277 3-11. **Detection Rate:** East Asian >99%.
Glycogen Storage Disease Type III - **Gene:** AGL. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000642:2-34. **Detection Rate:** East Asian >99%.
GNE Myopathy - **Gene:** GNE. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001128227:1-12. **Detection Rate:** East Asian >99%.
GNPTAB-related Disorders - **Gene:** GNPTAB. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_024312:1-21. **Detection Rate:** East Asian 92%.
HADHA-related Disorders - **Gene:** HADHA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000182:1-20. **Detection Rate:** East Asian >99%.
Hb Beta Chain-related Hemoglobinopathy (Including Beta Thalassemia and Sickle Cell Disease) - **Gene:** HBB. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000518:1-3. **Detection Rate:** East Asian >99%.
Hereditary Fructose Intolerance - **Gene:** ALDOB. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000035:2-9. **Detection Rate:** East Asian >99%.
Hexosaminidase A Deficiency (Including Tay-Sachs Disease) - **Gene:** HEXA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000520:1-14. **Detection Rate:** East Asian >99%.
HMG-CoA Lyase Deficiency - **Gene:** HMGCL. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000191:1-9. **Detection Rate:** East Asian >99%.
Holocarboxylase Synthetase Deficiency - **Gene:** HLCS. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000411:4-12. **Detection Rate:** East Asian >99%.
Homocystinuria, CBS-related - **Gene:** CBS. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000071:3-17. **Detection Rate:** East Asian >99%.
Hydrolethalus Syndrome - **Gene:** HYLS1. Autosomal Recessive. Sequencing with copy number analysis. **Exon:** NM_145014:4. **Detection Rate:** East Asian >99%.
Hypophosphatasia - **Gene:** ALPL. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000478:2-12. **Detection Rate:** East Asian >99%.
Isovaleric Acidemia - **Gene:** IVD. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_002225:1-12. **Detection Rate:** East Asian >99%.

Copyright 2021 Myriad Women's Health, Inc. All rights reserved.

Joubert Syndrome 2 - **Gene:** TMEM216. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001173990:1-5. **Detection Rate:** East Asian >99%.
Junctional Epidermolysis Bullosa, LAMA3-related - **Gene:** LAMA3. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000227:1-38. **Detection Rate:** East Asian >99%.
Junctional Epidermolysis Bullosa, LAMB3-related - **Gene:** LAMB3. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000228:2-23. **Detection Rate:** East Asian >99%.
Junctional Epidermolysis Bullosa, LAMC2-related - **Gene:** LAMC2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_005562:1-23. **Detection Rate:** East Asian >99%.
Krabbe Disease - **Gene:** GALC. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000153:1-17. **Detection Rate:** East Asian >99%.
Leigh Syndrome, French-Canadian Type - **Gene:** LRPPRC. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_133259:1-38. **Detection Rate:** East Asian >99%.
Lipoid Congenital Adrenal Hyperplasia - **Gene:** STAR. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000349:1-7. **Detection Rate:** East Asian >99%.
Lysosomal Acid Lipase Deficiency - **Gene:** LIPA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000235:2-10. **Detection Rate:** East Asian 98%.
Maple Syrup Urine Disease Type Ia - **Gene:** BCKDHA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000709:1-9. **Detection Rate:** East Asian >99%.
Maple Syrup Urine Disease Type Ib - **Gene:** BCKDHB. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_183050:1-10. **Detection Rate:** East Asian >99%.
Maple Syrup Urine Disease Type II - **Gene:** DBT. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001918:1-11. **Detection Rate:** East Asian 97%.
Medium Chain Acyl-CoA Dehydrogenase Deficiency - **Gene:** ACADM. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000016:1-12. **Detection Rate:** East Asian >99%.
Megalencephalic Leukoencephalopathy with Subcortical Cysts - **Gene:** MLC1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_015166 2-12. **Detection Rate:** East Asian >99%.
Metachromatic Leukodystrophy - **Gene:** ARSA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000487:1-8. **Detection Rate:** East Asian >99%.
Methylmalonic Acidemia, cblA Type - **Gene:** MMAA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_172250:2-7. **Detection Rate:** East Asian >99%.
Methylmalonic Acidemia, cblB Type - **Gene:** MMAB. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_052845:1-9. **Detection Rate:** East Asian >99%.
Methylmalonic Acidemia, MMUT-related - **Gene:** MMUT. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000255:2-13. **Detection Rate:** East Asian >99%.
Methylmalonic Aciduria and Homocystinuria, cblC Type - **Gene:** MMACHC. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_015506:1-4. **Detection Rate:** East Asian >99%.
MKS1-related Disorders - **Gene:** MKS1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_017777:1-18. **Detection Rate:** East Asian >99%.
Mucolipidosis III Gamma - **Gene:** GNPTG. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_032520:1-11. **Detection Rate:** East Asian 98%.
Mucolipidosis IV - **Gene:** MCOLN1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_020533:1-14. **Detection Rate:** East Asian >99%.
Mucopolysaccharidosis Type I - **Gene:** IDUA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000203:1-14. **Detection Rate:** East Asian >99%.
Mucopolysaccharidosis Type II - **Gene:** IDS. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000202:1-9. **Detection Rate:** East Asian 89%.
Mucopolysaccharidosis Type IIIA - **Gene:** SGSH. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000199:1-8. **Detection Rate:** East Asian >99%.
Mucopolysaccharidosis Type IIIB - **Gene:** NAGLU. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000263:1-6. **Detection Rate:** East Asian >99%.
Mucopolysaccharidosis Type IIIC - **Gene:** HGSNAT. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_152419:1-18. **Detection Rate:** East Asian >99%.
Muscular Dystrophy, LAMA2-related - **Gene:** LAMA2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000426:1-43,45-65. **Detection Rate:** East Asian 98%.
MYO7A-related Disorders - **Gene:** MYO7A. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000260:2-49. **Detection Rate:** East Asian >99%.
NEB-related Nemaline Myopathy - **Gene:** NEB. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001271208:3-80,117-183. **Detection Rate:** East Asian 92%.
Nephrotic Syndrome, NPHS1-related - **Gene:** NPHS1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_004646:1-29. **Detection Rate:** East Asian >99%.
Nephrotic Syndrome, NPHS2-related - **Gene:** NPHS2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_014625:1-8. **Detection Rate:** East Asian >99%.
Neuronal Ceroid Lipofuscinosis, CLN6-related - **Gene:** CLN6. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_017882:1-7. **Detection Rate:** East Asian >99%.
Niemann-Pick Disease Type C1 - **Gene:** NPC1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000271:1-25. **Detection Rate:** East Asian >99%.
Niemann-Pick Disease Type C2 - **Gene:** NPC2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_006432:1-5. **Detection Rate:** East Asian >99%.
Niemann-Pick Disease, SMPD1-related - **Gene:** SMPD1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000543:1-6. **Detection Rate:** East Asian >99%.
Nijmegen Breakage Syndrome - **Gene:** NBN. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_002485:1-16. **Detection Rate:** East Asian >99%.
Ornithine Transcarbamylase Deficiency - **Gene:** OTC. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000531:1-10. **Detection Rate:** East Asian 97%.
PCCA-related Propionic Acidemia - **Gene:** PCCA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000282:1-24. **Detection Rate:** East Asian 95%.
PCCB-related Propionic Acidemia - **Gene:** PCCB. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000532:1-15. **Detection Rate:** East Asian >99%.
PCDH15-related Disorders - **Gene:** PCDH15. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_033056:2-33. **Detection Rate:** East Asian 93%.
Pendred Syndrome - **Gene:** SLC26A4. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000441:2-21. **Detection Rate:** East Asian >99%.
Peroxisome Biogenesis Disorder Type 1 - **Gene:** PEX1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000466:1-24. **Detection Rate:** East Asian >99%.
Peroxisome Biogenesis Disorder Type 3 - **Gene:** PEX12. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000286:1-3. **Detection Rate:** East Asian >99%.
Peroxisome Biogenesis Disorder Type 4 - **Gene:** PEX6. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000287:1-17. **Detection Rate:** East Asian 97%.

Copyright 2021 Myriad Women's Health, Inc. All rights reserved.

Peroxisome Biogenesis Disorder Type 5 - **Gene:** PEX2. Autosomal Recessive. Sequencing with copy number analysis. **Exon:** NM_000318:4. **Detection Rate:** East Asian >99%.
Peroxisome Biogenesis Disorder Type 6 - **Gene:** PEX10. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_153818:1-6. **Detection Rate:** East Asian >99%.
Phenylalanine Hydroxylase Deficiency - **Gene:** PAH. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000277:1-13. **Detection Rate:** East Asian >99%.
POMGNT-related Disorders - **Gene:** POMGNT1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_017739:2-22. **Detection Rate:** East Asian 96%.
Pompe Disease - **Gene:** GAA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000152:2-20. **Detection Rate:** East Asian >99%.
PPT1-related Neuronal Ceroid Lipofuscinosis - **Gene:** PPT1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000310:1-9. **Detection Rate:** East Asian >99%.
Primary Carnitine Deficiency - **Gene:** SLC22A5. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_003060:1-10. **Detection Rate:** East Asian >99%.
Primary Hyperoxaluria Type 1 - **Gene:** AGXT. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000030:1-11. **Detection Rate:** East Asian >99%.
Primary Hyperoxaluria Type 2 - **Gene:** GRHPR. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_012203:1-9. **Detection Rate:** East Asian >99%.
Primary Hyperoxaluria Type 3 - **Gene:** HOGA1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_138413:1-7. **Detection Rate:** East Asian >99%.
Pycnodysostosis - **Gene:** CTSK. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000396:2-8. **Detection Rate:** East Asian >99%.
Pyruvate Carboxylase Deficiency - **Gene:** PC. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000920:3-22. **Detection Rate:** East Asian >99%.
Rhizomelic Chondrodysplasia Punctata Type 1 - **Gene:** PEX7. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000288:1-10. **Detection Rate:** East Asian >99%.
RTEL1-related Disorders - **Gene:** RTEL1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_032957:2-35. **Detection Rate:** East Asian >99%.
Sandhoff Disease - **Gene:** HEXB. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000521:1-14. **Detection Rate:** East Asian 98%.
Short-chain Acyl-CoA Dehydrogenase Deficiency - **Gene:** ACADS. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000017:1-10. **Detection Rate:** East Asian >99%.
Sjogren-Larsson Syndrome - **Gene:** ALDH3A2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000382:1-10. **Detection Rate:** East Asian 96%.
SLC26A2-related Disorders - **Gene:** SLC26A2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000112:2-3. **Detection Rate:** East Asian >99%.
Smith-Lemli-Opitz Syndrome - **Gene:** DHCR7. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001360:3-9. **Detection Rate:** East Asian >99%.
Spastic Paraplegia Type 15 - **Gene:** ZFYVE26. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_015346:2-42. **Detection Rate:** East Asian >99%.
Spinal Muscular Atrophy - **Gene:** SMN1. Autosomal Recessive. Spinal muscular atrophy. **Variant (1):** SMN1 copy number. **Detection Rate:** East Asian 93%.
Spondylothoracic Dysostosis - **Gene:** MESP2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001039958:1-2. **Detection Rate:** East Asian >99%.
TGM1-related Autosomal Recessive Congenital Ichthyosis - **Gene:** TGM1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000359 2-15. **Detection Rate:** East Asian >99%.
TPP1-related Neuronal Ceroid Lipofuscinosis - **Gene:** TPP1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000391:1-13. **Detection Rate:** East Asian >99%.
Tyrosine Hydroxylase Deficiency - **Gene:** TH. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_199292:1-14. **Detection Rate:** East Asian >99%.
Tyrosinemia Type I - **Gene:** FAH. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000137:1-14. **Detection Rate:** East Asian >99%.
Tyrosinemia Type II - **Gene:** TAT. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000353:2-12. **Detection Rate:** East Asian >99%.
USH1C-related Disorders - **Gene:** USH1C. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_005709:1-21. **Detection Rate:** East Asian >99%.
USH2A-related Disorders - **Gene:** USH2A. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_206933:2-72. **Detection Rate:** East Asian 98%.
Usher Syndrome Type 3 - **Gene:** CLRN1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_174878:1-3. **Detection Rate:** East Asian >99%.
Very-long-chain Acyl-CoA Dehydrogenase Deficiency - **Gene:** ACADVL. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000018:1-20. **Detection Rate:** East Asian >99%.
Wilson Disease - **Gene:** ATP7B. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000053:1-21. **Detection Rate:** East Asian >99%.
X-linked Adrenal Hypoplasia Congenita - **Gene:** NR0B1. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000475:1-2. **Detection Rate:** East Asian 97%.
X-linked Adrenoleukodystrophy - **Gene:** ABCD1. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000033:1-6. **Detection Rate:** East Asian 77%.
X-linked Alport Syndrome - **Gene:** COL4A5. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000495:1-51. **Detection Rate:** East Asian 96%.
X-linked Juvenile Retinoschisis - **Gene:** RS1. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000330:1-6. **Detection Rate:** East Asian 98%.
X-linked Myotubular Myopathy - **Gene:** MTM1. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000252:2-15. **Detection Rate:** East Asian 96%.
X-linked Severe Combined Immunodeficiency - **Gene:** IL2RG. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000206:1-8. **Detection Rate:** East Asian >99%.
Xeroderma Pigmentosum Group A - **Gene:** XPA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000380:1-6. **Detection Rate:** East Asian >99%.
Xeroderma Pigmentosum Group C - **Gene:** XPC. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_004628:1-16. **Detection Rate:** East Asian 97%.

Copyright 2021 Myriad Women's Health, Inc. All rights reserved.

Risk Calculations

Below are the risk calculations for all conditions tested. Negative results do not rule out the possibility of being a carrier. Residual risk is an estimate of each patient's post-test likelihood of being a carrier, while the reproductive risk represents an estimated likelihood that the patients' future children could inherit each disease. These risks are inherent to all carrier-screening tests, may vary by ethnicity, are predicated on a negative family history, and are present even given a negative test result. Inaccurate reporting of ethnicity may cause errors in risk calculation. In addition, average carrier rates are estimated using incidence or prevalence data from published scientific literature and/or reputable databases, where available, and are incorporated into residual risk calculations for each population/ethnicity. When population-specific data is not available for a condition, average worldwide incidence or prevalence is used. Further, incidence and prevalence data are only collected for the specified phenotypes (which include primarily the classic or severe forms of disease) and may not include alternate or milder disease manifestations associated with the gene. Actual incidence rates, prevalence rates, and carrier rates, and therefore actual residual risks, may be higher or lower than the estimates provided. Carrier rates, incidence/prevalence, and/or residual risks are not provided for some genes with biological or heritable properties that would make these estimates inaccurate. A '†' symbol indicates a positive result. See the full clinical report for interpretation and details. The reproductive risk presented is based on a hypothetical pairing with a partner of the same ethnic group.

Disease	DONOR 14289 Residual Risk	Reproductive Risk
6-pyruvoyl-tetrahydropterin Synthase Deficiency	1 in 24,000	< 1 in 1,000,000
Adenosine Deaminase Deficiency	1 in 22,000	< 1 in 1,000,000
Alpha Thalassemia, HBA1/HBA2-related	Alpha globin status: aa/aa.	Not calculated
Alpha-mannosidosis	1 in 35,000	< 1 in 1,000,000
Alpha-sarcoglycanopathy	1 in 34,000	< 1 in 1,000,000
Alstrom Syndrome	< 1 in 50,000	< 1 in 1,000,000
Andermann Syndrome	< 1 in 50,000	< 1 in 1,000,000
Argininemia	1 in 6,300	< 1 in 1,000,000
Argininosuccinic Aciduria	1 in 45,000	< 1 in 1,000,000
Aspartylglucosaminuria	< 1 in 50,000	< 1 in 1,000,000
Ataxia with Vitamin E Deficiency	< 1 in 50,000	< 1 in 1,000,000
Ataxia-telangiectasia	1 in 4,200	< 1 in 1,000,000
ATP7A-related Disorders	< 1 in 1,000,000	1 in 410,000
Autoimmune Polyglandular Syndrome Type 1	1 in 18,000	< 1 in 1,000,000
Autosomal Recessive Osteopetrosis Type 1	1 in 8,900	< 1 in 1,000,000
Autosomal Recessive Polycystic Kidney Disease, PKHD1-related	1 in 8,100	< 1 in 1,000,000
Autosomal Recessive Spastic Ataxia of Charlevoix-Saguenay	< 1 in 44,000	< 1 in 1,000,000
Bardet-Biedl Syndrome, BBS1-related	< 1 in 50,000	< 1 in 1,000,000
Bardet-Biedl Syndrome, BBS10-related	< 1 in 50,000	< 1 in 1,000,000
Bardet-Biedl Syndrome, BBS12-related	< 1 in 50,000	< 1 in 1,000,000
Bardet-Biedl Syndrome, BBS2-related	< 1 in 50,000	< 1 in 1,000,000
BCS1L-related Disorders	< 1 in 50,000	< 1 in 1,000,000
Beta-sarcoglycanopathy	1 in 39,000	< 1 in 1,000,000
Biotinidase Deficiency	1 in 67,000	< 1 in 1,000,000
Bloom Syndrome	< 1 in 50,000	< 1 in 1,000,000
Calpainopathy	1 in 11,000	< 1 in 1,000,000
Canavan Disease	1 in 9,700	< 1 in 1,000,000
Carbamoylphosphate Synthetase I Deficiency	1 in 45,000	< 1 in 1,000,000
Carnitine Palmitoyltransferase IA Deficiency	< 1 in 50,000	< 1 in 1,000,000
Carnitine Palmitoyltransferase II Deficiency	1 in 31,000	< 1 in 1,000,000
Cartilage-hair Hypoplasia	< 1 in 50,000	< 1 in 1,000,000
Cerebrotendinous Xanthomatosis	1 in 13,000	< 1 in 1,000,000
Citrullinemia Type 1	1 in 700	1 in 270,000
CLN3-related Neuronal Ceroid Lipofuscinosis	1 in 13,000	< 1 in 1,000,000
CLN5-related Neuronal Ceroid Lipofuscinosis	< 1 in 50,000	< 1 in 1,000,000
CLN8-related Neuronal Ceroid Lipofuscinosis	< 1 in 50,000	< 1 in 1,000,000
Cohen Syndrome	< 1 in 15,000	< 1 in 1,000,000
COL4A3-related Alport Syndrome	1 in 5,800	< 1 in 1,000,000
COL4A4-related Alport Syndrome	1 in 35,000	< 1 in 1,000,000
Combined Pituitary Hormone Deficiency, PROP1-related	1 in 6,100	< 1 in 1,000,000
Congenital Adrenal Hyperplasia, CYP11B1-related	1 in 8,400	< 1 in 1,000,000
Congenital Adrenal Hyperplasia, CYP21A2-related	1 in 550	1 in 140,000
Congenital Disorder of Glycosylation Type Ia	1 in 16,000	< 1 in 1,000,000
Congenital Disorder of Glycosylation Type Ic	< 1 in 50,000	< 1 in 1,000,000
Congenital Disorder of Glycosylation, MPI-related	< 1 in 50,000	< 1 in 1,000,000

Copyright 2021 Myriad Women's Health, Inc. All rights reserved.

Disease	DONOR 14289 Residual Risk	Reproductive Risk
Costeff Optic Atrophy Syndrome	< 1 in 50,000	< 1 in 1,000,000
Cystic Fibrosis	1 in 9,000	< 1 in 1,000,000
Cystinosis	1 in 22,000	< 1 in 1,000,000
D-bifunctional Protein Deficiency	1 in 9,000	< 1 in 1,000,000
Delta-sarcoglycanopathy	< 1 in 13,000	< 1 in 1,000,000
Dihydrolipoamide Dehydrogenase Deficiency	< 1 in 50,000	< 1 in 1,000,000
Dysferlinopathy	1 in 11,000	< 1 in 1,000,000
Dystrophinopathy (Including Duchenne/Becker Muscular Dystrophy)	Not calculated	Not calculated
ERCC6-related Disorders	1 in 8,400	< 1 in 1,000,000
ERCC8-related Disorders	< 1 in 2,300	< 1 in 1,000,000
EVC-related Ellis-van Creveld Syndrome	1 in 7,800	< 1 in 1,000,000
EVC2-related Ellis-van Creveld Syndrome	1 in 9,800	< 1 in 1,000,000
Fabry Disease	< 1 in 1,000,000	1 in 80,000
Familial Dysautonomia	< 1 in 50,000	< 1 in 1,000,000
Familial Hyperinsulinism, ABCC8-related	1 in 14,000	< 1 in 1,000,000
Familial Hyperinsulinism, KCNJ11-related	1 in 42,000	< 1 in 1,000,000
Familial Mediterranean Fever	< 1 in 50,000	< 1 in 1,000,000
Fanconi Anemia Complementation Group A	1 in 3,100	< 1 in 1,000,000
Fanconi Anemia, FANCC-related	< 1 in 50,000	< 1 in 1,000,000
FKRP-related Disorders	1 in 32,000	< 1 in 1,000,000
FKTN-related Disorders	1 in 110	1 in 40,000
Free Sialic Acid Storage Disorders	< 1 in 30,000	< 1 in 1,000,000
Galactokinase Deficiency	< 1 in 50,000	< 1 in 1,000,000
Galactosemia	1 in 32,000	< 1 in 1,000,000
Gamma-sarcoglycanopathy	< 1 in 3,800	< 1 in 1,000,000
Gaucher Disease	1 in 450	1 in 320,000
GJB2-related DFNB1 Nonsyndromic Hearing Loss and Deafness	1 in 3,300	1 in 450,000
GLB1-related Disorders	1 in 17,000	< 1 in 1,000,000
Glutaric Acidemia, GCDH-related	1 in 13,000	< 1 in 1,000,000
Glycine Encephalopathy, AMT-related	1 in 26,000	< 1 in 1,000,000
Glycine Encephalopathy, GLDC-related	1 in 4,200	< 1 in 1,000,000
Glycogen Storage Disease Type Ia	1 in 8,700	< 1 in 1,000,000
Glycogen Storage Disease Type Ib	1 in 35,000	< 1 in 1,000,000
Glycogen Storage Disease Type III	1 in 16,000	< 1 in 1,000,000
GNE Myopathy	< 1 in 50,000	< 1 in 1,000,000
GNPTAB-related Disorders	1 in 2,600	< 1 in 1,000,000
HADHA-related Disorders	1 in 25,000	< 1 in 1,000,000
Hb Beta Chain-related Hemoglobinopathy (Including Beta Thalassemia and Sickle Cell Disease)	1 in 900	1 in 34,000
Hereditary Fructose Intolerance	1 in 7,900	< 1 in 1,000,000
Hexosaminidase A Deficiency (Including Tay-Sachs Disease)	1 in 30,000	< 1 in 1,000,000
HMG-CoA Lyase Deficiency	< 1 in 50,000	< 1 in 1,000,000
Holocarboxylase Synthetase Deficiency	1 in 16,000	< 1 in 1,000,000
Homocystinuria, CBS-related	< 1 in 50,000	< 1 in 1,000,000
Hydrolethalus Syndrome	< 1 in 50,000	< 1 in 1,000,000
Hypophosphatasia	1 in 24,000	< 1 in 1,000,000
Isovaleric Acidemia	1 in 39,000	< 1 in 1,000,000
Joubert Syndrome 2	< 1 in 50,000	< 1 in 1,000,000
Junctional Epidermolysis Bullosa, LAMA3-related	< 1 in 50,000	< 1 in 1,000,000
Junctional Epidermolysis Bullosa, LAMB3-related	1 in 31,000	< 1 in 1,000,000
Junctional Epidermolysis Bullosa, LAMC2-related	< 1 in 50,000	< 1 in 1,000,000
Krabbe Disease	1 in 17,000	< 1 in 1,000,000
Leigh Syndrome, French-Canadian Type	< 1 in 50,000	< 1 in 1,000,000
Lipoid Congenital Adrenal Hyperplasia	1 in 40,000	< 1 in 1,000,000
Lysosomal Acid Lipase Deficiency	< 1 in 34,000	< 1 in 1,000,000
Maple Syrup Urine Disease Type Ia	1 in 43,000	< 1 in 1,000,000
Maple Syrup Urine Disease Type Ib	1 in 23,000	< 1 in 1,000,000
Maple Syrup Urine Disease Type II	1 in 15,000	< 1 in 1,000,000
Medium Chain Acyl-CoA Dehydrogenase Deficiency	1 in 11,000	< 1 in 1,000,000
Megalencephalic Leukoencephalopathy with Subcortical Cysts	< 1 in 50,000	< 1 in 1,000,000
Metachromatic Leukodystrophy	1 in 16,000	< 1 in 1,000,000
Methylmalonic Acidemia, cblA Type	< 1 in 50,000	< 1 in 1,000,000

Copyright 2021 Myriad Women's Health, Inc. All rights reserved.

Disease	DONOR 14289 Residual Risk	Reproductive Risk
Methylmalonic Acidemia, cblB Type	< 1 in 50,000	< 1 in 1,000,000
Methylmalonic Acidemia, MMUT-related	1 in 12,000	< 1 in 1,000,000
Methylmalonic Aciduria and Homocystinuria, cblC Type	1 in 33,000	< 1 in 1,000,000
MKS1-related Disorders	< 1 in 50,000	< 1 in 1,000,000
Mucolipidosis III Gamma	< 1 in 20,000	< 1 in 1,000,000
Mucolipidosis IV	< 1 in 50,000	< 1 in 1,000,000
Mucopolysaccharidosis Type I	1 in 16,000	< 1 in 1,000,000
Mucopolysaccharidosis Type II	1 in 520,000	1 in 120,000
Mucopolysaccharidosis Type IIIA	1 in 16,000	< 1 in 1,000,000
Mucopolysaccharidosis Type IIIB	1 in 42,000	< 1 in 1,000,000
Mucopolysaccharidosis Type IIIC	< 1 in 50,000	< 1 in 1,000,000
Muscular Dystrophy, LAMA2-related	1 in 5,700	< 1 in 1,000,000
MYO7A-related Disorders	1 in 15,000	< 1 in 1,000,000
NEB-related Nemaline Myopathy	1 in 1,200	1 in 400,000
Nephrotic Syndrome, NPHS1-related	< 1 in 50,000	< 1 in 1,000,000
Nephrotic Syndrome, NPHS2-related	1 in 35,000	< 1 in 1,000,000
Neuronal Ceroid Lipofuscinosis, CLN6-related	< 1 in 50,000	< 1 in 1,000,000
Niemann-Pick Disease Type C1	1 in 17,000	< 1 in 1,000,000
Niemann-Pick Disease Type C2	< 1 in 50,000	< 1 in 1,000,000
Niemann-Pick Disease, SMPD1-related	1 in 25,000	< 1 in 1,000,000
Nijmegen Breakage Syndrome	< 1 in 50,000	< 1 in 1,000,000
Ornithine Transcarbamylase Deficiency	< 1 in 1,000,000	1 in 140,000
PCCA-related Propionic Acidemia	1 in 4,200	< 1 in 1,000,000
PCCB-related Propionic Acidemia	1 in 6,500	< 1 in 1,000,000
PCDH15-related Disorders	1 in 3,300	< 1 in 1,000,000
Pendred Syndrome	1 in 6,400	< 1 in 1,000,000
Peroxisome Biogenesis Disorder Type 1	1 in 16,000	< 1 in 1,000,000
Peroxisome Biogenesis Disorder Type 3	< 1 in 50,000	< 1 in 1,000,000
Peroxisome Biogenesis Disorder Type 4	1 in 9,300	< 1 in 1,000,000
Peroxisome Biogenesis Disorder Type 5	< 1 in 71,000	< 1 in 1,000,000
Peroxisome Biogenesis Disorder Type 6	< 1 in 50,000	< 1 in 1,000,000
Phenylalanine Hydroxylase Deficiency	1 in 7,700	< 1 in 1,000,000
POMGNT-related Disorders	< 1 in 12,000	< 1 in 1,000,000
Pompe Disease	1 in 10,000	< 1 in 1,000,000
PPT1-related Neuronal Ceroid Lipofuscinosis	1 in 7,700	< 1 in 1,000,000
Primary Carnitine Deficiency	1 in 10,000	< 1 in 1,000,000
Primary Hyperoxaluria Type 1	1 in 13,000	< 1 in 1,000,000
Primary Hyperoxaluria Type 2	< 1 in 50,000	< 1 in 1,000,000
Primary Hyperoxaluria Type 3	1 in 20,000	< 1 in 1,000,000
Pycnodysostosis	1 in 43,000	< 1 in 1,000,000
Pyruvate Carboxylase Deficiency	1 in 25,000	< 1 in 1,000,000
Rhizomelic Chondrodysplasia Punctata Type 1	1 in 16,000	< 1 in 1,000,000
RTEL1-related Disorders	< 1 in 50,000	< 1 in 1,000,000
Sandhoff Disease	1 in 18,000	< 1 in 1,000,000
Short-chain Acyl-CoA Dehydrogenase Deficiency	1 in 9,700	< 1 in 1,000,000
Sjogren-Larsson Syndrome	< 1 in 12,000	< 1 in 1,000,000
SLC26A2-related Disorders	1 in 16,000	< 1 in 1,000,000
Smith-Lemli-Opitz Syndrome	< 1 in 50,000	< 1 in 1,000,000
Spastic Paraplegia Type 15	< 1 in 50,000	< 1 in 1,000,000
Spinal Muscular Atrophy	Negative for g.27134T>G SNP SMN1: 2 copies 1 in 700	1 in 150,000
Spondylothoracic Dysostosis	< 1 in 50,000	< 1 in 1,000,000
TGM1-related Autosomal Recessive Congenital Ichthyosis	1 in 22,000	< 1 in 1,000,000
TPP1-related Neuronal Ceroid Lipofuscinosis	1 in 30,000	< 1 in 1,000,000
Tyrosine Hydroxylase Deficiency	< 1 in 50,000	< 1 in 1,000,000
Tyrosinemia Type I	1 in 16,000	< 1 in 1,000,000
Tyrosinemia Type II	1 in 25,000	< 1 in 1,000,000
USH1C-related Disorders	1 in 44,000	< 1 in 1,000,000
USH2A-related Disorders	1 in 5,900	< 1 in 1,000,000
Usher Syndrome Type 3	1 in 41,000	< 1 in 1,000,000
Very-long-chain Acyl-CoA Dehydrogenase Deficiency	1 in 12,000	< 1 in 1,000,000

Copyright 2021 Myriad Women's Health, Inc. All rights reserved.

Disease	DONOR 14289 Residual Risk	Reproductive Risk
Wilson Disease	1 in 6,500	< 1 in 1,000,000
X-linked Adrenal Hypoplasia Congenita	< 1 in 1,000,000	< 1 in 1,000,000
X-linked Adrenoleukodystrophy	1 in 170,000	1 in 80,000
X-linked Alport Syndrome	Not calculated	Not calculated
X-linked Juvenile Retinoschisis	< 1 in 1,000,000	1 in 50,000
X-linked Myotubular Myopathy	Not calculated	Not calculated
X-linked Severe Combined Immunodeficiency	< 1 in 1,000,000	1 in 200,000
Xeroderma Pigmentosum Group A	1 in 10,000	< 1 in 1,000,000
Xeroderma Pigmentosum Group C	1 in 7,300	< 1 in 1,000,000

Copyright 2021 Myriad Women's Health, Inc. All rights reserved.

4399 Santa Anita Ave.
El Monte, CA, 91731
(p) 626-350-0537 (f) 626-454-1667
info@fulgentgenetics.com
www.fulgentgenetics.com

Patient Information:
14289, Donor
DOB: [redacted]
Sex: M
MR#:
Patient#: FT-PT8948392

Partner Information:
Not Tested

Physician:
Kuan, James
ATTN: Martinez, Lupe
San Diego Sperm Bank
4915 25th Avenue NE, Ste 204W
Seattle, WA 98105
Phone: (206) 588-1484

Laboratory:
Fulgent Therapeutics LLC
CAP#: 8042697
CLIA#: 05D2043189
Laboratory Director:
Dr. Amar Jariwala
Report Date: **Jun 18,2026**

Accession:
FT-7290010
Test#: FT-TS15617432
Specimen Type: Blood (EDTA)
Collected: Not Provided

Accession:
N/A

FINAL RESULTS

No carrier mutations identified

TEST PERFORMED

Single Gene Carrier Screening: TMEM38B
(1 Gene Panel: *TMEM38B*; gene sequencing with deletion and duplication analysis)

INTERPRETATION:

Notes and Recommendations:

- No carrier mutations were identified in the submitted specimen. A negative result does not rule out the possibility of a genetic predisposition nor does it rule out any pathogenic mutations in areas not assessed by this test or in regions that were covered at a level too low to reliably assess. Also, it does not rule out mutations that are of the sort not queried by this test; see Methods and Limitations for more information. A negative result reduces, but does not eliminate, the chance to be a carrier for any condition included in this screen. Please see the supplemental table for details.
- This carrier screening test does not screen for all possible genetic conditions, nor for all possible mutations in every gene tested. This report does not include variants of uncertain significance; only variants classified as pathogenic or likely pathogenic at the time of testing, and considered relevant for reproductive carrier screening, are reported. Please see the gene specific notes for details. Please note that the classification of variants can change over time.
- Patients may wish to discuss any carrier results with blood relatives, as there is an increased chance that they are also carriers. These results should be interpreted in the context of this individual's clinical findings, biochemical profile, and family history.
- Gene specific notes and limitations may be present. See below.
- Genetic counseling is recommended. Available genetic counselors and additional resources can be found at the National Society of Genetic Counselors (NSGC; https://www.nsgc.org).

4399 Santa Anita Ave.
El Monte, CA, 91731
(p) 626-350-0537 (f) 626-454-1667
info@fulgentgenetics.com
www.fulgentgenetics.com

GENES TESTED:

Custom Beacon Preconception Carrier Screening Panel - 1 Gene

This analysis was run using the Custom Beacon Preconception Carrier Screening Panel gene list. 1 genes were tested with 100.0% of targets sequenced at >20x coverage. For more gene-specific information and assistance with residual risk calculation, see the SUPPLEMENTAL TABLE.

TMEM38B

METHODS:

Genomic DNA was isolated from the submitted specimen indicated above (if cellular material was submitted). DNA was barcoded, and enriched for the coding exons of targeted genes using hybrid capture technology. Prepared DNA libraries were then sequenced using a Next Generation Sequencing technology. Following alignment to the human genome reference sequence (assembly GRCh37), variants were detected in regions of at least 10x coverage. For this specimen, 100.00% and 100.00% of coding regions and splicing junctions of genes listed had been sequenced with coverage of at least 10x and 20x, respectively, by NGS or by Sanger sequencing. The remaining regions did not have 10x coverage, and were not evaluated. Variants were interpreted manually using locus specific databases, literature searches, and other molecular biological principles. To minimize false positive results, any variants that do not meet internal quality standards are confirmed by Sanger sequencing. Variants classified as pathogenic, likely pathogenic, or risk allele which are located in the coding regions and nearby intronic regions (+/- 20bp) of the genes listed above are reported. Variants outside these intervals may be reported but are typically not guaranteed. When a single pathogenic or likely pathogenic variant is identified in a clinically relevant gene with autosomal recessive inheritance, the laboratory will attempt to ensure 100% coverage of coding sequences either through NGS or Sanger sequencing technologies ("fill-in"). All genes listed were evaluated for large deletions and/or duplications. However, single exon deletions or duplications will not be detected in this assay, nor will copy number alterations in regions of genes with significant pseudogenes. Putative deletions or duplications are analyzed using Fulgent Germline proprietary pipeline for this specimen. Bioinformatics: The FPLMv2.0 pipeline was used to analyze this specimen.

LIMITATIONS:

General Limitations

These test results and variant interpretation are based on the proper identification of the submitted specimen, accuracy of any stated familial relationships, and use of the correct human reference sequences at the queried loci. In very rare instances, errors may result due to mix-up or co-mingling of specimens. Positive results do not imply that there are no other contributors, genetic or otherwise, to future pregnancies, and negative results do not rule out the genetic risk to a pregnancy. Official gene names change over time. Fulgent uses the most up to date gene names based on HUGO Gene Nomenclature Committee (https://www.genenames.org) recommendations. If the gene name on report does not match that of ordered gene, please contact the laboratory and details can be provided. Result interpretation is based on the available clinical and family history information for this individual, collected published information, and Alamut annotation available at the time of reporting. This assay is not designed or validated for the detection of low-level mosaicism or somatic mutations. This assay will not detect certain types of genomic aberrations such as translocations, inversions, or repeat expansions other than specified genes. DNA alterations in regulatory regions or deep intronic regions (greater than 20bp from an exon) may not be detected by this test. Unless otherwise indicated, no additional assays have been performed to evaluate genetic changes in this specimen. There are technical limitations on the ability of DNA sequencing to detect small insertions and deletions. Our laboratory uses a sensitive detection algorithm, however these types of alterations are not detected as reliably as single nucleotide variants. Rarely, due to systematic chemical, computational, or human error, DNA variants may be missed. Although next generation sequencing technologies and our bioinformatics analysis significantly reduce the confounding contribution of pseudogene sequences or other highly-homologous sequences, sometimes these may still interfere with the technical ability of the assay to identify pathogenic alterations in both sequencing and deletion/duplication analyses. Deletion/duplication analysis can identify alterations of genomic regions which include two or more contiguous exons in size (whole blood specimens only); single exon deletions or duplications may occasionally be identified, but are not routinely detected by this test. Saliva swab samples may have reduced sensitivity for CNVs and a higher CNV failure rate. When novel DNA duplications are identified, it is not possible to discern the genomic location or orientation of the duplicated segment, hence the effect of the duplication cannot be predicted. Where deletions are detected, it is not always possible to determine whether the predicted product will remain in-frame or not. Unless otherwise indicated, deletion/duplication analysis has not been performed in regions that have been sequenced by Sanger.

4399 Santa Anita Ave.
El Monte, CA, 91731
(p) 626-350-0537 (f) 626-454-1667
info@fulgentgenetics.com
www.fulgentgenetics.com

Gene Specific Notes and Limitations

No gene specific limitations apply to the genes on the tested panel.

SIGNATURE:

Geetu Mendiratta-Vij, PhD, FACMG, CGMBS on **06/18/2026**
Laboratory Director, Fulgent

DISCLAIMER:

This test was developed, performed, and its performance characteristics determined by**Fulgent Therapeutics LLC** (CAP# 8042697, CLIA# 05D2043189), 4399 Santa Anita Ave., El Monte, CA 91731. It has not been cleared or approved by the FDA. The laboratory is regulated under CLIA as qualified to perform high-complexity testing. This test is used for clinical purposes. It should not be regarded as investigational or for research. Since genetic variation, as well as systematic and technical factors, can affect the accuracy of testing, the results of testing should always be interpreted in the context of clinical and familial data. For assistance with interpretation of these results, healthcare professionals may contact us directly at **(626) 350-0537** or **info@fulgentgenetics.com**. It is recommended that patients receive appropriate genetic counseling to explain the implications of the test result, including its residual risks, uncertainties and reproductive or medical options.

4399 Santa Anita Ave.
El Monte, CA, 91731
(p) 626-350-0537 (f) 626-454-1667
info@fulgentgenetics.com
www.fulgentgenetics.com

To view the supplemental table describing the carrier frequencies, detection rates, and residual risks associated with the genes tested on any Beacon panel, please visit the following link:
Beacon Expanded Carrier Screening Supplemental Table

4399 Santa Anita Ave.
El Monte, CA, 91731
(p) 626-350-0537 (f) 626-454-1667
info@fulgentgenetics.com
www.fulgentgenetics.com

Patient Information:
14289, Donor
DOB: [redacted]
Sex: M
MR#:
Patient#: FT-PT8948392

Partner Information:
Not Tested

Physician:
Kuan, James
ATTN: Martinez, Lupe
San Diego Sperm Bank
4915 25th Avenue NE, Ste 204W
Seattle, WA 98105
Phone: (206) 588-1484

Laboratory:
Fulgent Therapeutics LLC
CAP#: 8042697
CLIA#: 05D2043189
Laboratory Director:
Dr. Amar Jariwala
Report Date: **Sep 11,2025**

Accession:
FT-7290010
Test#: FT-TS15328675
Specimen Type: Blood (EDTA)
Collected: Not Provided

Accession:
N/A

FINAL RESULTS

No carrier mutations identified

TEST PERFORMED

Custom Beacon Preconception Carrier Screening Panel
(3 Gene Panel: *ACSF3, OCA2, and SLC12A3*; gene sequencing with deletion and duplication analysis)

INTERPRETATION:

Notes and Recommendations:

- No carrier mutations were identified in the submitted specimen. A negative result does not rule out the possibility of a genetic predisposition nor does it rule out any pathogenic mutations in areas not assessed by this test or in regions that were covered at a level too low to reliably assess. Also, it does not rule out mutations that are of the sort not queried by this test; see Methods and Limitations for more information. A negative result reduces, but does not eliminate, the chance to be a carrier for any condition included in this screen. Please see the supplemental table for details.
- This carrier screening test does not screen for all possible genetic conditions, nor for all possible mutations in every gene tested. This report does not include variants of uncertain significance; only variants classified as pathogenic or likely pathogenic at the time of testing, and considered relevant for reproductive carrier screening, are reported. Please see the gene specific notes for details. Please note that the classification of variants can change over time.
- Patients may wish to discuss any carrier results with blood relatives, as there is an increased chance that they are also carriers. These results should be interpreted in the context of this individual's clinical findings, biochemical profile, and family history.
- Gene specific notes and limitations may be present. See below.
- Genetic counseling is recommended. Available genetic counselors and additional resources can be found at the National Society of Genetic Counselors (NSGC; https://www.nsgc.org)

4399 Santa Anita Ave.
El Monte, CA, 91731
(p) 626-350-0537 (f) 626-454-1667
info@fulgentgenetics.com
www.fulgentgenetics.com

GENES TESTED:

Custom Beacon Preconception Carrier Screening Panel - 3 Genes

This analysis was run using the Custom Beacon Preconception Carrier Screening Panel gene list. 3 genes were tested with 100.0% of targets sequenced at >20x coverage. For more gene-specific information and assistance with residual risk calculation, see the SUPPLEMENTAL TABLE.

ACSF3, OCA2, SLC12A3

METHODS:

Genomic DNA was isolated from the submitted specimen indicated above (if cellular material was submitted). DNA was barcoded, and enriched for the coding exons of targeted genes using hybrid capture technology. Prepared DNA libraries were then sequenced using a Next Generation Sequencing technology. Following alignment to the human genome reference sequence (assembly GRCh37), variants were detected in regions of at least 10x coverage. For this specimen, 100.00% and 100.00% of coding regions and splicing junctions of genes listed had been sequenced with coverage of at least 10x and 20x, respectively, by NGS or by Sanger sequencing. The remaining regions did not have 10x coverage, and were not evaluated. Variants were interpreted manually using locus specific databases, literature searches, and other molecular biological principles. To minimize false positive results, any variants that do not meet internal quality standards are confirmed by Sanger sequencing. Variants classified as pathogenic, likely pathogenic, or risk allele which are located in the coding regions and nearby intronic regions (+/- 20bp) of the genes listed above are reported. Variants outside these intervals may be reported but are typically not guaranteed. When a single pathogenic or likely pathogenic variant is identified in a clinically relevant gene with autosomal recessive inheritance, the laboratory will attempt to ensure 100% coverage of coding sequences either through NGS or Sanger sequencing technologies ("fill-in"). All genes listed were evaluated for large deletions and/or duplications. However, single exon deletions or duplications will not be detected in this assay, nor will copy number alterations in regions of genes with significant pseudogenes. Putative deletions or duplications are analyzed using Fulgent Germline proprietary pipeline for this specimen. Bioinformatics: The FPLMv2.0 pipeline was used to analyze this specimen.

LIMITATIONS:

General Limitations

These test results and variant interpretation are based on the proper identification of the submitted specimen, accuracy of any stated familial relationships, and use of the correct human reference sequences at the queried loci. In very rare instances, errors may result due to mix-up or co-mingling of specimens. Positive results do not imply that there are no other contributors, genetic or otherwise, to future pregnancies, and negative results do not rule out the genetic risk to a pregnancy. Official gene names change over time. Fulgent uses the most up to date gene names based on HUGO Gene Nomenclature Committee (https://www.genenames.org) recommendations. If the gene name on report does not match that of ordered gene, please contact the laboratory and details can be provided. Result interpretation is based on the available clinical and family history information for this individual, collected published information, and Alamut annotation available at the time of reporting. This assay is not designed or validated for the detection of low-level mosaicism or somatic mutations. This assay will not detect certain types of genomic aberrations such as translocations, inversions, or repeat expansions other than specified genes. DNA alterations in regulatory regions or deep intronic regions (greater than 20bp from an exon) may not be detected by this test. Unless otherwise indicated, no additional assays have been performed to evaluate genetic changes in this specimen. There are technical limitations on the ability of DNA sequencing to detect small insertions and deletions. Our laboratory uses a sensitive detection algorithm, however these types of alterations are not detected as reliably as single nucleotide variants. Rarely, due to systematic chemical, computational, or human error, DNA variants may be missed. Although next generation sequencing technologies and our bioinformatics analysis significantly reduce the confounding contribution of pseudogene sequences or other highly-homologous sequences, sometimes these may still interfere with the technical ability of the assay to identify pathogenic alterations in both sequencing and deletion/duplication analyses. Deletion/duplication analysis can identify alterations of genomic regions which include one whole gene (buccal swab specimens and whole blood specimens) and are two or more contiguous exons in size (whole blood specimens only); single exon deletions or duplications may occasionally be identified, but are not routinely detected by this test. When novel DNA duplications are identified, it is not possible to discern the genomic location or orientation of the duplicated segment, hence the effect of the duplication cannot be predicted. Where deletions are detected, it is not always possible to determine whether the predicted product will remain in-frame or not. Unless otherwise indicated, deletion/duplication analysis has not been performed in regions that have been sequenced by Sanger.

4399 Santa Anita Ave.
El Monte, CA, 91731
(p) 626-350-0537 (f) 626-454-1667
info@fulgentgenetics.com
www.fulgentgenetics.com

Gene Specific Notes and Limitations

No gene specific limitations apply to the genes on the tested panel.

SIGNATURE:

Wei Kelly M.D., Ph.D., FACMG, CGMBS on **09/11/2025**
Laboratory Director, Fulgent

DISCLAIMER:

This test was developed, performed, and its performance characteristics determined by**Fulgent Therapeutics LLC** (CAP# 8042697, CLIA# 05D2043189), 4399 Santa Anita Ave., El Monte, CA 91731. It has not been cleared or approved by the FDA. The laboratory is regulated under CLIA as qualified to perform high-complexity testing. This test is used for clinical purposes. It should not be regarded as investigational or for research. Since genetic variation, as well as systematic and technical factors, can affect the accuracy of testing, the results of testing should always be interpreted in the context of clinical and familial data. For assistance with interpretation of these results, healthcare professionals may contact us directly at **(626) 350-0537** or **info@fulgentgenetics.com**. It is recommended that patients receive appropriate genetic counseling to explain the implications of the test result, including its residual risks, uncertainties and reproductive or medical options.

4399 Santa Anita Ave.
El Monte, CA, 91731
(p) 626-350-0537 (f) 626-454-1667
info@fulgentgenetics.com
www.fulgentgenetics.com

To view the supplemental table describing the carrier frequencies, detection rates, and residual risks associated with the genes tested on any Beacon panel, please visit the following link:
Beacon Expanded Carrier Screening Supplemental Table

15 Crawford St., STE 100
Needham, MA 02494
(p) 626-350-0537
(f) 626-454-1667
Lab Director: Arash Radfar M.D.
CLIA: 22D0957540

fulgent | INFORM DIAGNOSTICS™ A FULGENT GENETICS COMPANY
Metroplex Pathology Associates
Lakewood Pathology Associates
Cohen Dermatopathology
Therapath Partners

Patient Information:
14289, Donor
DOB: [redacted]
Sex: M
MR#:
Patient#: FT-PT8948392

Partner Information:
Not Tested

Physician:
Kuan, James
ATTN: SSB Genetics, Dept
San Diego Sperm Bank
4915 25th Avenue NE, Ste 204W
Seattle, WA 98105
Phone: (206) 588-1484

Laboratory:
Fulgent Therapeutics LLC
CAP#: 8042697
CLIA#: 05D2043189
Laboratory Director:
Dr. Amar Jariwala
Report Date: **Jan 22,2025**

Accession:
FT-7290010
Test#: FT-TS15056689
Specimen Type: Blood (EDTA)
Collected: Not Provided

Accession:
N/A

FINAL RESULTS

No carrier mutations identified

TEST PERFORMED

Single Gene Carrier Screening: OTOF

(1 Gene Panel: *OTOF*; gene sequencing with deletion and duplication analysis)

INTERPRETATION:

Notes and Recommendations:

- No carrier mutations were identified in the submitted specimen. A negative result does not rule out the possibility of a genetic predisposition nor does it rule out any pathogenic mutations in areas not assessed by this test or in regions that were covered at a level too low to reliably assess. Also, it does not rule out mutations that are of the sort not queried by this test; see Methods and Limitations for more information. A negative result reduces, but does not eliminate, the chance to be a carrier for any condition included in this screen. Please see the supplemental table for details.
- This carrier screening test does not screen for all possible genetic conditions, nor for all possible mutations in every gene tested. This report does not include variants of uncertain significance; only variants classified as pathogenic or likely pathogenic at the time of testing, and considered relevant for reproductive carrier screening, are reported. Please see the gene specific notes for details. Please note that the classification of variants can change over time.
- Patients may wish to discuss any carrier results with blood relatives, as there is an increased chance that they are also carriers. These results should be interpreted in the context of this individual's clinical findings, biochemical profile, and family history.
- Gene specific notes and limitations may be present. See below.
- Genetic counseling is recommended. Available genetic counselors and additional resources can be found at the National Society of Genetic Counselors (NSGC; https://www.nsgc.org)

15 Crawford St., STE 100
Needham, MA 02494
(p) 626-350-0537
(f) 626-454-1667
Lab Director: Arash Radfar M.D.
CLIA: 22D0957540

GENES TESTED:

Custom Beacon Preconception Carrier Screening Panel - Gene

This analysis was run using the Custom Beacon Preconception Carrier Screening Panel gene list. 1 genes were tested with 100.0% of targets sequenced at >20x coverage. For more gene-specific information and assistance with residual risk calculation, see the SUPPLEMENTAL TABLE.

OTOF

METHODS:

Genomic DNA was isolated from the submitted specimen indicated above (if cellular material was submitted). DNA was barcoded, and enriched for the coding exons of targeted genes using hybrid capture technology. Prepared DNA libraries were then sequenced using a Next Generation Sequencing technology. Following alignment to the human genome reference sequence (assembly GRCh37), variants were detected in regions of at least 10x coverage. For this specimen, 100.00% and 100.00% of coding regions and splicing junctions of genes listed had been sequenced with coverage of at least 10x and 20x, respectively, by NGS or by Sanger sequencing. The remaining regions did not have 10x coverage, and were not evaluated. Variants were interpreted manually using locus specific databases, literature searches, and other molecular biological principles. To minimize false positive results, any variants that do not meet internal quality standards are confirmed by Sanger sequencing. Variants classified as pathogenic, likely pathogenic, or risk allele which are located in the coding regions and nearby intronic regions (+/- 20bp) of the genes listed above are reported. Variants outside these intervals may be reported but are typically not guaranteed. When a single pathogenic or likely pathogenic variant is identified in a clinically relevant gene with autosomal recessive inheritance, the laboratory will attempt to ensure 100% coverage of coding sequences either through NGS or Sanger sequencing technologies ("fill-in"). All genes listed were evaluated for large deletions and/or duplications. However, single exon deletions or duplications will not be detected in this assay, nor will copy number alterations in regions of genes with significant pseudogenes. Putative deletions or duplications are analyzed using Fulgent Germline proprietary pipeline for this specimen. Bioinformatics: The Fulgent Germline v2019.2 pipeline was used to analyze this specimen.

LIMITATIONS:

General Limitations

These test results and variant interpretation are based on the proper identification of the submitted specimen, accuracy of any stated familial relationships, and use of the correct human reference sequences at the queried loci. In very rare instances, errors may result due to mix-up or co-mingling of specimens. Positive results do not imply that there are no other contributors, genetic or otherwise, to future pregnancies, and negative results do not rule out the genetic risk to a pregnancy. Official gene names change over time. Fulgent uses the most up to date gene names based on HUGO Gene Nomenclature Committee (https://www.genenames.org) recommendations. If the gene name on report does not match that of ordered gene, please contact the laboratory and details can be provided. Result interpretation is based on the available clinical and family history information for this individual, collected published information, and Alamut annotation available at the time of reporting. This assay is not designed or validated for the detection of low-level mosaicism or somatic mutations. This assay will not detect certain types of genomic aberrations such as translocations, inversions, or repeat expansions other than specified genes. DNA alterations in regulatory regions or deep intronic regions (greater than 20bp from an exon) may not be detected by this test. Unless otherwise indicated, no additional assays have been performed to evaluate genetic changes in this specimen. There are technical limitations on the ability of DNA sequencing to detect small insertions and deletions. Our laboratory uses a sensitive detection algorithm, however these types of alterations are not detected as reliably as single nucleotide variants. Rarely, due to systematic chemical, computational, or human error, DNA variants may be missed. Although next generation sequencing technologies and our bioinformatics analysis significantly reduce the confounding contribution of pseudogene sequences or other highly-homologous sequences, sometimes these may still interfere with the technical ability of the assay to identify pathogenic alterations in both sequencing and deletion/duplication analyses. Deletion/duplication analysis can identify alterations of genomic regions which include one whole gene (buccal swab specimens and whole blood specimens) and are two or more contiguous exons in size (whole blood specimens only); single exon deletions or duplications may occasionally be identified, but are not routinely detected by this test. When novel DNA duplications are identified, it is not possible to discern the genomic location or orientation of the duplicated segment, hence the effect of the duplication cannot be predicted. Where deletions are detected, it is not always possible to determine whether the predicted product will remain in-frame or not. Unless otherwise indicated, deletion/duplication analysis has not been performed in regions that have been sequenced by Sanger.

15 Crawford St., STE 100
Needham, MA 02494
(p) 626-350-0537
(f) 626-454-1667
Lab Director: Arash Radfar M.D.
CLIA: 22D0957540

fulgent | INFORM DIAGNOSTICS™ A FULGENT GENETICS COMPANY
Metroplex Pathology Associates
Lakewood Pathology Associates
Cohen Dermatopathology
Therapath Partners

Gene Specific Notes and Limitations

No gene specific limitations apply to the genes on the tested panel.

SIGNATURE:

Dr. Harry Gao, DABMG, FACMG on **1/22/2025**
Laboratory Director, Fulgent

DISCLAIMER:

This test was developed and its performance characteristics determined by Fulgent Therapeutics LLC CAP #8042697 CLIA #05D2043189; 4399 Santa Anita Ave., El Monte, CA, 91731. It has not been cleared or approved by the FDA. The laboratory is regulated under CLIA as qualified to perform high-complexity testing. This test is used for clinical purposes. It should not be regarded as investigational or for research. Since genetic variation, as well as systematic and technical factors, can affect the accuracy of testing, the results of testing should always be interpreted in the context of clinical and familial data. For assistance with interpretation of these results, healthcare professionals may contact us directly at **626-350-0537** or by email at **info@fulgentgenetics.com**. It is recommended that patients receive appropriate genetic counseling to explain the implications of the test result, including its residual risks, uncertainties and reproductive or medical options.

To view the supplemental table describing the carrier frequencies, detection rates, and residual risks associated with the genes tested on any Beacon panel, please visit the following link:
Beacon Expanded Carrier Screening Supplemental Table

INVITAE

INVITAE CARRIER SCREEN RESULTS

Patient name:	DONOR 14289	**Sample type:**	Blood	**Report date:**	16-JAN-2024
DOB:	[redacted]	**Sample collection date:**	08-JAN-2024	**Invitae #:**	RQ5975707
Sex assigned at birth:	Male	**Sample accession date:**	09-JAN-2024	**Clinical team:**	Guadalupe Martinez Dr. James Kuan
Gender:					
Patient ID (MRN):					

Reason for testing

Gamete donor

Test performed

Invitae Carrier Screen

RESULT: NEGATIVE

This carrier test evaluated 6 gene(s) for genetic changes (variants) that are associated with an increased risk of having a child with a genetic condition. Knowledge of carrier status for one of these conditions may provide information that can be used to assist with family planning and/or preparation. Carrier screening is not intended for diagnostic purposes. To identify a potential genetic basis for a condition in the individual being tested, diagnostic testing for the gene(s) of interest is recommended.

This test did not identify any genetic changes in the gene(s) analyzed that are currently recognized as clinically significant. This negative result reduces, but does not eliminate, the chance that this individual is a carrier for conditions caused by any of the genes tested. This individual may still be a carrier for a genetic condition that is not evaluated by this test.

Next steps

- Even for genes that have a negative test result, there is always a small risk that an individual could still be a carrier. This is called “residual risk.” See the Carrier detection rates and residual risks document.
- Discussion with a physician and/or genetic counselor is recommended to further review the implications of this test result and to understand these results in the context of any family history of a genetic condition.
- All patients, regardless of result, may wish to consider additional screening for hemoglobinopathies by complete blood count (CBC) and hemoglobin electrophoresis, if this has not already been completed.
- Individuals can register their tests at **https://www.invitae.com/patients/** to access online results, educational resources, and next steps.

Laboratory Director: Jeana DaRe, PhD, FACMG
Invitae 1400 16th Street, San Francisco, CA 94103 · clientservices@invitae.com · 800.436.3037
© 2024 Invitae Corporation

Residual risk

No carrier test can detect 100% of carriers. There still remains a small risk of being a carrier after a negative test (residual risk). Residual risk values assume a negative family history and are inferred from published carrier frequencies and estimated detection rates based on testing technologies used at Invitae. You can view Invitae's complete Carrier detection rates and residual risks document (containing all carrier genes) online at https://www.invitae.com/carrier-residual-risks/. Additionally, the order-specific information for this report is available to download in the portal (under this order's documents) or can be requested by contacting Invitae Client Services. The complete Carrier detection rates and residual risks document will not be applicable for any genes with specimen-specific limitations in sequencing and/or deletion/duplication coverage. Please see the final bullet point in the Limitations section of this report to view if this specimen had any gene-specific coverage gaps.

© 2024 Invitae Corporation

Genes analyzed

This table represents a complete list of genes analyzed for this individual, including the relevant gene transcript(s). If more than one transcript is listed for a single gene, variants were reported using the first transcript listed unless otherwise indicated in the report. An asterisk (*) indicates that this gene has a limitation. Please see the Limitations section for details. Results are negative, unless otherwise indicated in the report.

GENE	TRANSCRIPT
ABCA4	NM_000350.2
CDH23	NM_022124.5

GENE	TRANSCRIPT
DNAH11	NM_001277115.1
GBE1	NM_000158.3

GENE	TRANSCRIPT
NDUFV1	NM_007103.3
TMEM67	NM_153704.5

© 2024 Invitae Corporation

Methods

- Genomic DNA obtained from the submitted sample is enriched for targeted regions using a hybridization-based protocol, and sequenced using Illumina technology. Unless otherwise indicated, all targeted regions are sequenced with ≥50x depth or are supplemented with additional analysis. Reads are aligned to a reference sequence (GRCh37), and sequence changes are identified and interpreted in the context of a single clinically relevant transcript, indicated in the Genes Analyzed table. Enrichment and analysis focus on the coding sequence of the indicated transcripts, 20bp of flanking intronic sequence, and other specific genomic regions demonstrated to be causative of disease at the time of assay design. Promoters, untranslated regions, and other non-coding regions are not otherwise interrogated. Exonic deletions and duplications are called using an in-house algorithm that determines copy number at each target by comparing the read depth for each target in the proband sequence with both mean read-depth and read-depth distribution, obtained from a set of clinical samples. Markers across the X and Y chromosomes are analyzed for quality control purposes and may detect deviations from the expected sex chromosome complement. Such deviations may be included in the report in accordance with internal guidelines. Variants are reported according to the Human Genome Variation Society (HGVS) guidelines. Confirmation of the presence and location of reportable variants is performed as needed based on stringent criteria, using one of several validated orthogonal approaches (PubMed ID 30610921). Sequencing is performed by Invitae Corporation (1400 16th Street, San Francisco, CA 94103, #05D2040778). Confirmatory sequencing is performed by Invitae Corporation (1400 16th Street, San Francisco, CA 94103, #05D2040778).

  The following additional analyses are performed if relevant to the requisition. For GBA the reference genome has been modified to mask the sites of polymorphic paralog sequence variants (PSVs) in both the gene and pseudogene. For CYP21A2 and GBA, if one or more reportable variants, gene conversion, or fusion event is identified via our NGS pipeline (see Limitations), these variants are confirmed by PacBio sequencing of an amplicon generated by long-range PCR and subsequent short-range PCR. In some cases, it may not be possible to disambiguate between the gene and pseudogene. For GJB2, the reportable range includes large upstream deletions overlapping GJB6. For HBA1/2, the reference genome has been modified to force some sequencing reads derived from HBA1 to align to HBA2, and variant calling algorithms are modified to support an expectation of 4 alleles in these regions. HBA1/2 copy number calling is performed by a custom hypothesis testing algorithm which generates diplotype calls. If sequence data for a sample does not support a unique high confidence match from among hypotheses tested, that sample is flagged for manual review. Copy number variation is only reported for coding sequence of HBA1 and HBA2 and the HS-40 region. This assay does not distinguish among the -α3.7 subtypes, and all -α3.7 variants are called as HBA1 deletions. This assay may not detect overlapping copy gain and copy loss events when the breakpoints of those events are similar. For FMR1, cytosine-guanine-guanine (CGG) triplet repeats in the 5' untranslated region (5' UTR) of the FMR1 gene are detected by triplet repeat-primed PCR (RP-PCR) with fluorescently labeled primers followed by capillary electrophoresis. Reference ranges: Normal: <45 CGG repeats, intermediate: 45-54 CGG repeats, premutation: 55-200 CGG repeats, full mutation: >200 CGG repeats. For alleles with 55-90 triplet repeats, the region surrounding the FMR1 repeat is amplified by PCR. The PCR amplicons are then processed through PacBio SMRTBell library prep and sequenced using PacBio long read technology. The number of AGG interruptions within the 55-90 triplet repeat is read directly from the resulting DNA sequences.

- This report only includes variants that have a clinically significant association with the conditions tested as of the report date. Variants of uncertain significance, benign variants, and likely benign variants are not included in this report. However, if additional evidence becomes available to indicate that the clinical significance of a variant has changed, Invitae may update this report and provide notification.
- A PMID is a unique identifier referring to a published, scientific paper. Search by PMID at **http://www.ncbi.nlm.nih.gov/pubmed**.
- An rsID is a unique identifier referring to a single genomic position, and is used to associate population frequency information with sequence changes at that position. Reported population frequencies are derived from a number of public sites that aggregate data from large-scale population sequencing projects, including ExAC (**http://exac.broadinstitute.org**), gnomAD (**http://gnomad.broadinstitute.org**), and dbSNP (**http://ncbi.nlm.nih.gov/SNP**).

Disclaimer

DNA studies do not constitute a definitive test for the selected condition(s) in all individuals. It should be realized that there are possible sources of error. Errors can result from trace contamination, rare technical errors, rare genetic variants that interfere with analysis, recent scientific developments, and alternative classification systems. This test should be one of many aspects used by the healthcare provider to help with a diagnosis and treatment plan, but it is not a diagnosis itself. This test was developed and its performance characteristics determined by Invitae. It has not been cleared or approved by

© 2024 Invitae Corporation

the FDA. The laboratory is regulated under the Clinical Laboratory Improvement Act (CLIA) as qualified to perform high-complexity clinical tests (CLIA ID: 05D2040778). This test is used for clinical purposes. It should not be regarded as investigational or for research.

Limitations

- Based on validation study results, this assay achieves >99% analytical sensitivity and specificity for single nucleotide variants, insertions and deletions <15bp in length, and exon-level deletions and duplications. Invitae's methods also detect insertions and deletions larger than 15bp but smaller than a full exon but sensitivity for these may be marginally reduced. Invitae's deletion/duplication analysis determines copy number at a single exon resolution at virtually all targeted exons. However, in rare situations, single-exon copy number events may not be analyzed due to inherent sequence properties or isolated reduction in data quality. Certain types of variants, such as structural rearrangements (e.g. inversions, gene conversion events, translocations, etc.) or variants embedded in sequence with complex architecture (e.g. short tandem repeats or segmental duplications), may not be detected. Additionally, it may not be possible to fully resolve certain details about variants, such as mosaicism, phasing, or mapping ambiguity. Unless explicitly guaranteed, sequence changes in the promoter, non-coding exons, and other non-coding regions are not covered by this assay. Please consult the test definition on our website for details regarding regions or types of variants that are covered or excluded for this test. This report reflects the analysis of an extracted genomic DNA sample. While this test is intended to reflect the analysis of extracted genomic DNA from a referred patient, in very rare cases the analyzed DNA may not represent that individual's constitutional genome, such as in the case of a circulating hematolymphoid neoplasm, bone marrow transplant, blood transfusion, chimerism, culture artifact or maternal cell contamination. Interpretations are made on the assumption that any clinical information provided, including specimen identity, is accurate.

This report has been released utilizing a validated procedure approved by:

Jeana DaRe

Jeana DaRe, Ph.D., FACMG
Laboratory Director

jd_0835_pr

© 2024 Invitae Corporation

INVITAE

CARRIER DETECTION RATES AND RESIDUAL RISKS

This table is relevant to patient report RQ5975707
Issue date: 01/16/2024

This table displays residual risks after a negative result for each of the genes and corresponding disorders. The values provided assume a negative family history and the absence of symptoms for each disorder. For genes associated with both dominant and recessive inheritance, the numbers in this table apply to the recessive condition(s) associated with the gene, unless otherwise noted. Residual risk values are provided for disorders when carrier frequency is greater than 1 in 500. For disorders with carrier frequency equal to, or less than, 1 in 500, residual risk is considered to be reduced substantially. When provided, residual risk values are inferred from published carrier frequencies, and estimated detection rates are based on testing technologies used at Invitae. Residual risks are provided only as a guide for assessing approximate risk given a negative result; values may vary based on the ethnic background(s) of an individual. For any genes marked with an asterisk*, refer to the Limitations section of the patient report for detailed coverage information. In the case of a sample-specific limitation, "N/A" indicates that a residual risk value could not be calculated. AR = autosomal recessive, XL = X-linked, AD = autosomal dominant.

DISORDER (INHERITANCE)	GENE	ETHNICITY	CARRIER FREQUENCY	DETECTION RATE	RISK TO BE A CARRIER AFTER NEGATIVE RESULT
ABCA4-related conditions (AR) NM_000350.2	ABCA4	Pan-ethnic	1 in 45	90%	1 in 441
CDH23-related conditions (AR) NM_022124.5	CDH23	Pan-ethnic	1 in 202	95%	1 in 4020
GBE1-related conditions (AR) NM_000158.3	GBE1	Pan-ethnic	1 in 387	99%	1 in 38600
Mitochondrial complex I deficiency 4 (AR) NM_007103.3	NDUFV1	Pan-ethnic	1 in 387	99%	1 in 38600
Primary ciliary dyskinesia (DNAH11-related) (AR) NM_001277115.1	DNAH11	Pan-ethnic	1 in 211	99%	1 in 21000
TMEM67-related conditions (AR) NM_153704.5	TMEM67	Pan-ethnic	1 in 316	99%	1 in 31500

WITHIN THE US | p: 800-436-3037 | f: 415-276-4164 | e: clientservices@invitae.com | OUTSIDE THE US | p: +1 415-930-4018 | f: +1 510 417-3922 | e: globalsupport@invitae.com
INVITAE | 1400 16th Street, San Francisco, CA 94103, USA | www.invitae.com/contact | © 2024 Invitae Corporation. All Rights Reserved.

Integrated GENETICS
LabCorp Specialty Testing Group

Client/Sending Facility:
San Diego Sperm Bank

8950 Villa La Jolla Dr Ste B214
La Jolla, CA 92037
Ph: (858)732-8500
CAB-50

LCLS Specimen Number:	**300-253-2173-0**	Account Number:	04087470
Patient Name:	**14289, DONOR**	Ordering Physician:	**J OLLIFFE**
Date of Birth:	[redacted]	Specimen Type:	**BLOOD**
Gender:	M	Client Reference:	
Patient ID:		Date Collected:	10/27/2021
Lab Number:	YU21-94923 L	Date Received:	10/29/2021
Indications:	NOT GIVEN	Date Reported:	**11/17/2021**

Test: **Chromosome, Blood, Routine**

Cells Counted:	20	Cells Karyotyped:	2
Cells Analyzed:	20	Band Resolution:	500

CYTOGENETIC RESULT: 46,XY

INTERPRETATION: NORMAL MALE KARYOTYPE

Cytogenetic analysis of PHA stimulated cultures has revealed a MALE karyotype with an apparently normal GTG banding pattern in all cells observed.

This result does not exclude the possibility of subtle rearrangements below the resolution of cytogenetics or congenital anomalies due to other etiologies.

Technical Component-Processing performed by LabCorp CLIA 34D1008914, 1904 TW Alexander Dr, Research Triangle Park, NC 27709. Medical Director, Anjen Chenn, M.D., Ph.D.


Technical Component-Chromosome analysis performed by Laboratory Corporation of America Inc., CLIA 29D0539775. 3600 Mill Street, Suite 100, Reno, NV 89502. Laboratory Director, Frank Ryan, Ph.D., (ABCC).

Integrated GENETICS
LabCorp Specialty Testing Group

Client/Sending Facility:
San Diego Sperm Bank

8950 Villa La Jolla Dr Ste B214
La Jolla, CA 92037
Ph: (858)732-8500
CAB-50

LCLS Specimen Number:	**300-253-2173-0**	Account Number:	04087470
Patient Name:	**14289, DONOR**	Ordering Physician:	**J OLLIFFE**
Date of Birth:	[redacted]	Specimen Type:	**BLOOD**
Gender:	M	Client Reference:	
Patient ID:		Date Collected:	10/27/2021
Lab Number:	YU21-94923 L	Date Received:	10/29/2021



Integrated GENETICS
LabCorp Specialty Testing Group

Client/Sending Facility:
San Diego Sperm Bank

8950 Villa La Jolla Dr Ste B214
La Jolla, CA 92037
Ph: (858)732-8500
CAB-50

LCLS Specimen Number:	**300-253-2173-0**	Account Number:	04087470
Patient Name:	**14289, DONOR**	Ordering Physician:	**J OLLIFFE**
Date of Birth:	[redacted]	Specimen Type:	**BLOOD**
Gender:	M	Client Reference:	
Patient ID:		Date Collected:	10/27/2021
Lab Number:	YU21-94923 L	Date Received:	10/29/2021



PATRICK A. LENNON, PHD, FACMG

Anjen Chenn, M.D., Ph.D.
Medical Director
Stuart Schwartz, PhD
National Director of Cytogenetics

Technical component performed by Laboratory Corporation of America Holdings,
1904 TW Alexander Drive , RTP , NC , 27709-0153 (800) 345-4363
Professional Component performed by LabCorp CLIA 34D1008914, 1904 TW Alexander Dr, Research Triangle Park, NC 27709. Medical Director, Anjen Chenn, M.D., Ph.D.
Integrated Genetics is a brand used by Esoterix Genetic Laboratories, LLC, a wholly-owned subsidiary of Laboratory Corporation of America Holdings.
This document contains private and confidential health information **protected by state and federal law.**

14289, Donor
Patient ID:
Specimen ID: **300-253-2172-0**

DOB: [redacted]
Age: [redacted]
Sex: **Male**

Patient Report
Account Number: **04087470**
Ordering Physician: **J OLLIFFE**

labcorp

Ordered Items: **LP+12AC+CBC/D/Plt+UA+Rh+ABO...**

Date Collected: **10/27/2021** Date Received: **10/28/2021** Date Reported: **11/01/2021** Fasting: **No**

LP+12AC+CBC/D/Plt+UA+Rh+ABO...

Test	Current Result and Flag	Previous Result and Date	Units	Reference Interval
Glucose [01]	79		mg/dL	65-99
Uric Acid [01]	6.0		mg/dL	3.8-8.4
		Therapeutic target for gout patients: <6.0 **Please note reference interval change**		
BUN [01]	13		mg/dL	6-20
Creatinine [01]	0.80		mg/dL	0.76-1.27
eGFR If NonAfricn Am	130		mL/min/1.73	>59
eGFR If Africn Am	151		mL/min/1.73	>59
	In accordance with recommendations from the NKF-ASN Task force, Labcorp is in the process of updating its eGFR calculation to the 2021 CKD-EPI creatinine equation that estimates kidney function without a race variable.			
Calcium [01]	9.4		mg/dL	8.7-10.2
Protein, Total [01]	6.8		g/dL	6.0-8.5
Albumin [01]	4.6		g/dL	4.1-5.2
Bilirubin, Total [01]	0.4		mg/dL	0.0-1.2
Alkaline Phosphatase [01]	85		IU/L	51-125
		Please note reference interval change		
LDH [01]	172		IU/L	121-224
AST (SGOT) [01]	23		IU/L	0-40
ALT (SGPT) [01]	31		IU/L	0-44
Cholesterol, Total [01]	143		mg/dL	100-169
Triglycerides [01]	57		mg/dL	0-89
HDL Cholesterol [01]	40		mg/dL	>39
LDL Chol Calc (NIH)	91		mg/dL	0-109
LDL/HDL Ratio	2.3		ratio	0.0-3.6
Please Note: [01]				
		LDL/HDL Ratio Men Women 1/2 Avg.Risk 1.0 1.5 Avg.Risk 3.6 3.2 2X Avg.Risk 6.2 5.0 3X Avg.Risk 8.0 6.1		
Hgb Fractionation by CE: [02]				
Hgb F [02]	0.0		%	0.0-2.0
Hgb A [02]	97.1		%	96.4-98.8
Hgb A2 [02]	2.9		%	1.8-3.2
Hgb S [02]	0.0		%	0.0
Interpretation: [02]				
	Normal hemoglobin present; no hemoglobin variant or thalassemia observed.			

©2021 Laboratory Corporation of America® Holdings
All Rights Reserved - Enterprise Report Version 2.00

This document contains private and confidential health information protected by state and federal law.
If you have received this document in error please call 858-668-3700

LP+12AC+CBC/D/Plt+UA+Rh+ABO... (Cont.)

ABO Grouping [01]	O		
Rh Factor [01]	Positive		
Please note: Prior records for this patient's ABO / Rh type are not available for additional verification.			
. [01]			
CBC, Platelet Ct, and Diff [01]			
WBC [01]	6.5	x10E3/uL	3.4-10.8
RBC [01]	4.66	x10E6/uL	4.14-5.80
Hemoglobin [01]	13.3	g/dL	13.0-17.7
Hematocrit [01]	41.5	%	37.5-51.0
MCV [01]	89	fL	79-97
MCH [01]	28.5	pg	26.6-33.0
MCHC [01]	32.0	g/dL	31.5-35.7
RDW [01]	12.3	%	11.6-15.4
Platelets [01]	217	x10E3/uL	150-450
Neutrophils [01]	79	%	Not Estab.
Lymphs [01]	14	%	Not Estab.
Monocytes [01]	5	%	Not Estab.
Eos [01]	1	%	Not Estab.
Basos [01]	1	%	Not Estab.
Neutrophils (Absolute) [01]	5.2	x10E3/uL	1.4-7.0
Lymphs (Absolute) [01]	0.9	x10E3/uL	0.7-3.1
Monocytes(Absolute) [01]	0.3	x10E3/uL	0.1-0.9
Eos (Absolute) [01]	0.0	x10E3/uL	0.0-0.4
Baso (Absolute) [01]	0.0	x10E3/uL	0.0-0.2
Immature Granulocytes [01]	0	%	Not Estab.
Immature Grans (Abs) [01]	0.0	x10E3/uL	0.0-0.1
. [01]			
Urinalysis Gross Exam [01]			
Specific Gravity [01]	1.013		1.005-1.030
pH [01]	6.0		5.0-7.5
Urine-Color [01]	Yellow		Yellow
Appearance [01]	Clear		Clear
WBC Esterase [01]	Negative		Negative
Protein [01]	Negative		Negative/Trace
Glucose [01]	Negative		Negative
Ketones [01]	Negative		Negative
Occult Blood [01]	Negative		Negative
Bilirubin [01]	Negative		Negative
Urobilinogen,Semi-Qn [01]	0.2	mg/dL	0.2-1.0
Nitrite, Urine [01]	Negative		Negative
Microscopic Examination [01]			
Microscopic not indicated and not performed.			

©2021 Laboratory Corporation of America® Holdings
All Rights Reserved - Enterprise Report Version 2.00
This document contains private and confidential health information protected by state and federal law.
If you have received this document in error please call 858-668-3700