myriad WOMEN'S HEALTH

RESULTS RECIPIENT
SEATTLE SPERM BANK
Attn: Jeffrey Olliffe
4915 25th Ave NE Ste 204W
Seattle, WA 98105
Phone: (206) 588-1484
Fax: (206) 466-4696
NPI: 1306838271
Report Date: 11/18/2021

MALE
DONOR 16010
DOB:
Ethnicity: East Asian
Sample Type: EDTA Blood
Date of Collection: 11/10/2021
Date Received: 11/11/2021
Date Tested: 11/17/2021
Barcode: 11004513015652
Accession ID:
CSLKWYNUGWXLEFV
Indication: Egg or sperm donor

FEMALE
N/A

Foresight® Carrier Screen

POSITIVE: CARRIER

ABOUT THIS TEST

The **Myriad Foresight Carrier Screen** utilizes sequencing, maximizing coverage across all DNA regions tested, to help you learn about your chance to have a child with a genetic disease.

RESULTS SUMMARY

Risk Details	DONOR 16010	Partner
Panel Information	Foresight Carrier Screen Universal Panel Fundamental Plus Panel Fundamental Panel **(175 conditions tested)**	N/A
POSITIVE: CARRIER Wilson Disease **Reproductive Risk: 1 in 260** **Inheritance: Autosomal Recessive**	**CARRIER*** NM_000053.3(ATP7B):c.2621C>T(A874V) heterozygote	The reproductive risk presented is based on a hypothetical pairing with a partner of the same ethnic group. Carrier testing should be considered. See "Next Steps".

*Carriers generally do not experience symptoms.

No disease-causing mutations were detected in any other gene tested. A complete list of all conditions tested can be found on page 7.

CLINICAL NOTES

- None

NEXT STEPS

- Carrier testing should be considered for the diseases specified above for the patient's partner.
- Patients are recommended to discuss reproductive risks with their health care provider or a genetic counselor. Patients may also wish to discuss any positive results with blood relatives, as there is an increased chance that they are also carriers.

Copyright 2021 Myriad Women's Health, Inc. All rights reserved.

POSITIVE: CARRIER

Wilson Disease

Reproductive risk: 1 in 260
Risk before testing: 1 in 17,000

Gene: ATP7B | **Inheritance Pattern:** Autosomal Recessive

	Patient	**DONOR 16010**	**No partner tested**
Result		✚ Carrier	N/A
Variant(s)		NM_000053.3(ATP7B):c.2621C>T(A874V) heterozygote	N/A
Methodology		Sequencing with copy number analysis (v3.1)	N/A
Interpretation		This individual is a carrier of Wilson disease. Carriers generally do not experience symptoms.	N/A
Detection rate		>99%	N/A
Exons tested		NM_000053:1-21.	N/A

What Is Wilson Disease?

Wilson disease, caused by mutations in the *ATP7B* gene, is an inherited condition that causes the body to retain too much copper. The extra copper deposits in the liver, brain, kidneys, and eyes leading to damage and scarring in the tissues and causing the affected organs to stop working properly.

Symptoms typically first appear in childhood or early adolescence, but they can appear as early as age 3 or as late as age 70. The most common symptoms are liver disease and neurological impairment. Liver disease can first appear as fatigue, abdominal pain, or a yellowing of the skin and the whites of the eye (jaundice). Sometimes the result is liver failure, which requires a liver transplant. Neurological impairment can include tremors, clumsiness, problems walking, trouble swallowing, and impaired thinking.

Some individuals with Wilson disease also develop psychiatric problems including depression, anxiety, behavioral problems, mood changes, and difficulty with attention. Extra copper in the kidneys may also cause problems that sometimes lead to kidney failure. Individuals with Wilson disease may also have arthritis, weaker bones, heart problems, pancreatitis, and endocrine disorders. Extra copper in the eyes can cause brown circles, referred to as Kayser-Fleischer rings, around the colored part of the eyes, but this does not affect vision.

How Common Is Wilson Disease?

The prevalence of Wilson disease is approximately 1 in 30,000 individuals worldwide. In China, Japan, and Sardinia, Wilson disease is more common and may affect as many as 1 in 10,000 individuals.

How Is Wilson Disease Treated?

Wilson disease should be treated as soon as possible. Most individuals with the condition take D-penicillamine or trientine by mouth several times a day. This medicine traps (chelates) the excessive copper and helps remove it from the body through the urine. This can help prevent or reduce some of the liver, neurological, and psychiatric symptoms. People on this medication often also need to take

Copyright 2021 Myriad Women's Health, Inc. All rights reserved.

vitamin B6 (pyridoxine) as a supplement. Treatment should continue for the whole life of the patient. Sometimes a liver transplant will still be needed. People with Wilson disease should also avoid eating food that contains a lot of copper, such as organs, chocolate, mushrooms, shellfish, and nuts.

What Is the Prognosis for an Individual with Wilson Disease?

Frequent monitoring of the blood and urine and lifelong treatment are important. Without proper treatment, an individual with Wilson disease usually suffers progressively worse liver, neurological, and psychiatric symptoms until they die from liver or neurological disease. With proper treatment, individuals with Wilson disease can often have normal lifespans.

Copyright 2021 Myriad Women's Health, Inc. All rights reserved.

Methods and Limitations

DONOR 16010 [Foresight Carrier Screen]: Sequencing with copy number analysis, spinal muscular atrophy, analysis of homologous regions, and alpha thalassemia (HBA1/HBA2) sequencing with targeted copy number analysis (Assay(s): DTS v3.2).

Sequencing with copy number analysis

High-throughput sequencing and read-depth-based copy number analysis are used to analyze the genes listed in the Conditions Tested section of the report. Except where otherwise noted, the region of interest (ROI) comprises the indicated coding regions and 20 non-coding bases flanking each region. In a minority of cases where genomic features (e.g., long homopolymers) compromise calling fidelity, the affected non-coding bases are excluded from the ROI. The ROI is sequenced to a minimum acceptable read depth, and the sequences are compared to a reference genomic sequence (Genome Reference Consortium Human Build 37 [GRCh37]/hg19). On average, 99% of all bases in the ROI are sequenced at a read depth that is greater than the minimum read depth. Sequence variants may not be detected in areas of lower sequence coverage. Insertions and deletions may not be detected as accurately as single-nucleotide variants. Select genes or regions for which pseudogenes or other regions of homology impede reliable variant detection may be assayed using alternate technology, or they may be excluded from the ROI. *CFTR* and *DMD* testing includes analysis for exon-level deletions and duplications with an average sensitivity of ~99%. Only exon-level deletions are assayed for other genes on the panel and such deletions are detected with a sensitivity of ≥75%. Selected founder deletions may be detected at slightly higher sensitivity. Affected exons and/or breakpoints of copy number variants are estimated from junction reads, where available, or using the positions of affected probes. Only exons known to be included in the region affected by a copy number variant are provided in the variant nomenclature. In some cases, the copy number variant may be larger or smaller than indicated. If *GJB2* is tested, large upstream deletions involving the *GJB6* and/or *CRYL1* genes that may affect the expression of *GJB2* are also analyzed.

Spinal muscular atrophy

Targeted copy number analysis via high-throughput sequencing is used to determine the copy number of exon 7 of the *SMN1* gene. Other genetic variants may interfere with this analysis. Some individuals with two copies of *SMN1* are "silent" carriers with both *SMN1* genes on one chromosome and no copies of the gene on the other chromosome. This is more likely in individuals who have two copies of the *SMN1* gene and are positive for the g.27134T>G single-nucleotide polymorphism (SNP) (PMID: 9199562, 23788250, and 28676062), which affects the reported residual risk; Ashkenazi Jewish or Asian patients with this genotype have a high post-test likelihood of being carriers for SMA and are reported as carriers. The g.27134T>G SNP is only reported in individuals who have two copies of *SMN1*.

Analysis of homologous regions

A combination of high-throughput sequencing, read-depth-based copy number analysis, and targeted genotyping is used to determine the number of functional gene copies and/or the presence of selected loss-of-function variants in certain genes that have homology to other genomic regions. The precise breakpoints of large deletions in these genes cannot be determined but are instead estimated from copy number analysis. Pseudogenes may interfere with this analysis, especially when many pseudogene copies are present.

If *CYP21A2* is tested, patients who have one or more additional copies of the *CYP21A2* gene and a pathogenic variant may or may not be a carrier of 21-hydroxylase deficient CAH, depending on the chromosomal location of the variants (phase). Benign *CYP21A2* gene duplications and/or triplications will only be reported in this context. Some individuals with two functional *CYP21A2* gene copies may be "silent" carriers, with two gene copies resulting from a duplication on one chromosome and a gene deletion on the other chromosome. This and other similar rare carrier states, where complementary changes exist between the chromosomes, may not be detected by the assay. Given that the true incidence of non-classic CAH is unknown, the residual carrier and reproductive risk numbers on the report are based only on the published incidence for classic CAH. However, the published prevalence of non-classic CAH is highest in individuals of Ashkenazi Jewish, Hispanic, Italian, and Yugoslav descent. Therefore, the residual and reproductive risks are likely an underestimate for CAH, especially in the aforementioned populations, as they do not account for non-classic CAH.

Copyright 2021 Myriad Women's Health, Inc. All rights reserved.

Alpha thalassemia (HBA1/HBA2) sequencing with targeted copy number analysis

High-throughput sequencing and read-depth-based copy number analysis are used to identify sequence variation and functional gene copies within the region of interest (ROI) of *HBA1* and *HBA2*, which includes the listed exons plus 20 intronic flanking bases. In a minority of cases where genomic features (e.g , long homopolymers) compromise calling fidelity, the affected intronic bases are not included in the ROI. The ROI is sequenced to a minimum acceptable read depth, and the sequences are compared to a reference genomic sequence (Genome Reference Consortium Human Build 37 [GRCh37]/hg19). On average, 99% of all bases in the ROI are sequenced at a read depth that is greater than the minimum read depth. Sequence variants may not be detected in areas of lower sequence coverage. Insertions and deletions may not be detected as accurately as single-nucleotide variants. For large deletions or duplications in these genes, the precise breakpoints cannot be determined but are instead estimated from copy number analysis. This assay has been validated to detect up to two additional copies of each alpha globin gene. In rare instances where assay results suggest greater than two additional copies are present, this will be noted but the specific number of gene copies observed will not be provided.

Extensive sequence homology exists between *HBA1* and *HBA2*. This sequence homology can prevent certain variants from being localized to one gene over the other. In these instances, variant nomenclature will be provided for both genes. If follow-up testing is indicated for patients with the nomenclature provided for both genes, both *HBA1* and *HBA2* should be tested. Some individuals with four functional alpha globin gene copies may be "silent" carriers, with three gene copies resulting from triplication on one chromosome and a single gene deletion on the other chromosome. This and other similar rare carrier states, where complementary changes exist between the chromosomes, may not be detected by the assay.

Interpretation of reported variants

The classification and interpretation of all variants identified in this assay reflects the current state of Myriad's scientific understanding at the time this report was issued. Variants are classified according to internally defined criteria, which are compatible with the ACMG Standards and Guidelines for the Interpretation of Sequence Variants (PMID: 25741868). Variants that have been determined by Myriad to be disease-causing or likely disease-causing (i.e. pathogenic or likely pathogenic) are reported. Benign variants, variants of uncertain clinical significance (VUS), and variants not directly associated with the specified disease phenotype(s) are not reported. Variant classification and interpretation may change for a variety of reasons, including but not limited to, improvements to classification techniques, availability of additional scientific information, and observation of a variant in more patients. f the classification of one or more variants identified in this patient changes, an updated report reflecting the new classification generally will not be issued. If an updated report is issued, the variants reported may change based on their current classification. This can include changes to the variants displayed in gene specific 'variants tested' sections. Healthcare providers may contact Myriad directly to request updated variant classification information specific to this test result.

Limitations

The MWH Foresight Carrier Screen is designed to detect and report germline (constitutional) alterations. Mosaic (somatic) variation may not be detected, and if it is detected, it may not be reported. If more than one variant is detected in a gene, additional studies may be necessary to determine if those variants lie on the same chromosome or different chromosomes (phase). This test is not designed to detect sex-chromosome copy number variations. If present, sex-chromosome abnormalities may significantly reduce test sensitivity for X-linked conditions. Variant interpretation and residual and reproductive risk estimations assume a normal karyotype and may be different for individuals with abnormal karyotypes. The test does not fully address all inherited forms of intellectual disability, birth defects, or heritable diseases. Furthermore, not all forms of genetic variation are detected by this assay (i.e., duplications [except in specified genes], chromosomal rearrangements, structural abnormalities, etc.). Additional testing may be appropriate for some individuals. Pseudogenes and other regions of homology may interfere with this analysis. In an unknown number of cases, other genetic variation may interfere with variant detection. Rare carrier states where complementary changes exist between the chromosomes may not be detected by the assay. Other possible sources of diagnostic error include sample mix-up, trace contamination, bone marrow transplantation, blood transfusions, and technical or analytical errors.

Detection rates are determined using published scientific literature and/or reputable databases, when available, to estimate the fraction of disease alleles, weighted by frequency, that the methodology is predicted to be able or unable to detect. Detection rates are approximate and only account for analytical sensitivity. Certain variants that have been previously described in the literature may not be reported, if there is insufficient evidence for pathogenicity. Detection rates do not account for the disease specific rates of *de novo* variation.

This test was developed, and its performance characteristics determined by, Myriad Women's Health, Inc. It has not been cleared or approved by the US Food and Drug Administration (FDA). The FDA does not require this test to go through premarket review. This test is used for clinical purposes. It should not be regarded as investigational or for research. This laboratory is certified under the Clinical Laboratory Improvement Amendments of 1988 (CLIA) as qualified to perform high-complexity clinical testing. These results are adjunctive to the ordering physician's evaluation. CLIA Number: #05D1102604.

Copyright 2021 Myriad Women's Health, Inc. All rights reserved.

Incidental Findings

Unless otherwise indicated, these results and interpretations are limited to the specific disease panel(s) requested by the ordering healthcare provider. In some cases, standard data analyses may identify genetic findings beyond the region(s) of interest specified by the test, and such findings may not be reported. These findings may include genomic abnormalities with major, minor, or no, clinical significance.


If you have questions or would like more information about any of the test methods or limitations, please contact (888) 268-6795.

Resources

GENOME CONNECT | http://www.genomeconnect.org

Patients can share their reports using research registries such as Genome Connect, an online research registry building a genetics and health knowledge base. Genome Connect provides patients, physicians, and researchers an opportunity to share genetic information to support the study of the impact of genetic variation on health conditions.

SENIOR LABORATORY DIRECTOR



Karla R. Bowles, PhD, FACMG, CGMB

Report content approved by Maria Alfaro, PhD, FACMG, CGMB on Nov 18, 2021

Copyright 2021 Myriad Women's Health, Inc. All rights reserved.

Conditions Tested

6-pyruvoyl-tetrahydropterin Synthase Deficiency - **Gene:** PTS. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000317:1-6. **Detection Rate:** East Asian >99%.

Adenosine Deaminase Deficiency - **Gene:** ADA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000022:1-12. **Detection Rate:** East Asian 98%.

Alpha Thalassemia, HBA1/HBA2-related - **Genes:** HBA1, HBA2. Autosomal Recessive. Alpha thalassemia (HBA1/HBA2) sequencing with targeted copy number analysis. **Exons:** NM_000517:1-3; NM_000558:1-3. **Variants (16):** -(alpha)20.5, --BRIT, --MEDI, --MEDII, --SEA, --THAI or --FIL, -alpha3.7, -alpha4.2, HBA1+HBA2 deletion, Hb Constant Spring, Poly(A) AATAAA>AATA--, Poly(A) AATAAA>AATAAG, Poly(A) AATAAA>AATGAA, anti3.7, anti4.2, del HS-40. **Detection Rate:** East Asian >99%.

Alpha-mannosidosis - **Gene:** MAN2B1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000528:1-23. **Detection Rate:** East Asian >99%.

Alpha-sarcoglycanopathy - **Gene:** SGCA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000023:1-9. **Detection Rate:** East Asian >99%.

Alstrom Syndrome - **Gene:** ALMS1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_015120:1-23. **Detection Rate:** East Asian >99%.

Andermann Syndrome - **Gene:** SLC12A6. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_133647:1-25. **Detection Rate:** East Asian >99%.

Argininemia - **Gene:** ARG1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000045:1-8. **Detection Rate:** East Asian 97%.

Argininosuccinic Aciduria - **Gene:** ASL. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001024943:1-16. **Detection Rate:** East Asian >99%.

Aspartylglucosaminuria - **Gene:** AGA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000027:1-9. **Detection Rate:** East Asian >99%.

Ataxia with Vitamin E Deficiency - **Gene:** TTPA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000370:1-5. **Detection Rate:** East Asian >99%.

Ataxia-telangiectasia - **Gene:** ATM. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000051:2-63. **Detection Rate:** East Asian 96%.

ATP7A-related Disorders - **Gene:** ATP7A. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000052:2-23. **Detection Rate:** East Asian 90%.

Autoimmune Polyglandular Syndrome Type 1 - **Gene:** AIRE. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000383:1-14. **Detection Rate:** East Asian >99%.

Autosomal Recessive Osteopetrosis Type 1 - **Gene:** TCIRG1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_006019:2-20. **Detection Rate:** East Asian 96%.

Autosomal Recessive Polycystic Kidney Disease, PKHD1-related - **Gene:** PKHD1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_138694 2-67. **Detection Rate:** East Asian >99%.

Autosomal Recessive Spastic Ataxia of Charlevoix-Saguenay - **Gene:** SACS. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_014363 2-10. **Detection Rate:** East Asian 99%.

Bardet-Biedl Syndrome, BBS1-related - **Gene:** BBS1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_024649:1-17. **Detection Rate:** East Asian >99%.

Bardet-Biedl Syndrome, BBS10-related - **Gene:** BBS10. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_024685:1-2. **Detection Rate:** East Asian >99%.

Bardet-Biedl Syndrome, BBS12-related - **Gene:** BBS12. Autosomal Recessive. Sequencing with copy number analysis. **Exon:** NM_152618:2. **Detection Rate:** East Asian >99%.

Bardet-Biedl Syndrome, BBS2-related - **Gene:** BBS2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_031885:1-17. **Detection Rate:** East Asian >99%.

BCS1L-related Disorders - **Gene:** BCS1L. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_004328:3-9. **Detection Rate:** East Asian >99%.

Beta-sarcoglycanopathy - **Gene:** SGCB. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000232:1-6. **Detection Rate:** East Asian >99%.

Biotinidase Deficiency - **Gene:** BTD. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000060:1-4. **Detection Rate:** East Asian >99%.

Bloom Syndrome - **Gene:** BLM. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000057:2-22. **Detection Rate:** East Asian >99%.

Calpainopathy - **Gene:** CAPN3. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000070:1-24. **Detection Rate:** East Asian 99%.

Canavan Disease - **Gene:** ASPA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000049:1-6. **Detection Rate:** East Asian 98%.

Carbamoylphosphate Synthetase I Deficiency - **Gene:** CPS1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001875:1-38. **Detection Rate:** East Asian >99%.

Carnitine Palmitoyltransferase IA Deficiency - **Gene:** CPT1A. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001876:2-19. **Detection Rate:** East Asian >99%.

Carnitine Palmitoyltransferase II Deficiency - **Gene:** CPT2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000098:1-5. **Detection Rate:** East Asian >99%.

Cartilage-hair Hypoplasia - **Gene:** RMRP. Autosomal Recessive. Sequencing with copy number analysis. **Exon:** NR_003051:1. **Detection Rate:** East Asian >99%.

Cerebrotendinous Xanthomatosis - **Gene:** CYP27A1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000784:1-9. **Detection Rate:** East Asian >99%.

Citrullinemia Type 1 - **Gene:** ASS1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000050:3-16. **Detection Rate:** East Asian 86%.

CLN3-related Neuronal Ceroid Lipofuscinosis - **Gene:** CLN3. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001042432 2-16. **Detection Rate:** East Asian >99%.

CLN5-related Neuronal Ceroid Lipofuscinosis - **Gene:** CLN5. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_006493:1-4. **Detection Rate:** East Asian >99%.

CLN8-related Neuronal Ceroid Lipofuscinosis - **Gene:** CLN8. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_018941:2-3. **Detection Rate:** East Asian >99%.

Cohen Syndrome - **Gene:** VPS13B. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_017890:2-62. **Detection Rate:** East Asian 97%.

COL4A3-related Alport Syndrome - **Gene:** COL4A3. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000091:1-52. **Detection Rate:** East Asian 94%.

COL4A4-related Alport Syndrome - **Gene:** COL4A4. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000092:2-48. **Detection Rate:** East Asian >99%.

Combined Pituitary Hormone Deficiency, PROP1-related - **Gene:** PROP1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_006261:1-3. **Detection Rate:** East Asian >99%.

Congenital Adrenal Hyperplasia, CYP11B1-related - **Gene:** CYP11B1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000497:1-9. **Detection Rate:** East Asian 97%.

Congenital Adrenal Hyperplasia, CYP21A2-related - **Gene:** CYP21A2. Autosomal Recessive. Analysis of homologous regions. **Variants (13):** CYP21A2 deletion, CYP21A2 duplication, CYP21A2 triplication, G111Vfs*21, I173N, L308Ffs*6, P31L, Q319*, Q319*+CYP21A2dup, R357W, V282L, [I237N;V238E;M240K], c.293-13C>G. **Detection Rate:** East Asian 88%.

Congenital Disorder of Glycosylation Type Ia - **Gene:** PMM2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000303:1-8. **Detection Rate:** East Asian >99%.

Copyright 2021 Myriad Women's Health, Inc. All rights reserved.

Congenital Disorder of Glycosylation Type Ic - **Gene:** ALG6. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_013339:2-15. **Detection Rate:** East Asian >99%.
Congenital Disorder of Glycosylation, MPI-related - **Gene:** MPI. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_002435:1-8. **Detection Rate:** East Asian >99%.
Costeff Optic Atrophy Syndrome - **Gene:** OPA3. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_025136:1-2. **Detection Rate:** East Asian >99%.
Cystic Fibrosis - **Gene:** CFTR. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000492:1-27. IVS8-5T allele analysis is only reported in the presence of the R117H mutation. **Detection Rate:** East Asian >99%.
Cystinosis - **Gene:** CTNS. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_004937:3-12. **Detection Rate:** East Asian >99%.
D-bifunctional Protein Deficiency - **Gene:** HSD17B4. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000414:1-24. **Detection Rate:** East Asian 98%.
Delta-sarcoglycanopathy - **Gene:** SGCD. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000337:2-9. **Detection Rate:** East Asian 96%.
Dihydrolipoamide Dehydrogenase Deficiency - **Gene:** DLD. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000108:1-14. **Detection Rate:** East Asian >99%.
Dysferlinopathy - **Gene:** DYSF. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_003494:1-55. **Detection Rate:** East Asian 98%.
Dystrophinopathy (Including Duchenne/Becker Muscular Dystrophy) - **Gene:** DMD. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_004006:1-79. **Detection Rate:** East Asian 99%.
ERCC6-related Disorders - **Gene:** ERCC6. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000124:2-21. **Detection Rate:** East Asian 96%.
ERCC8-related Disorders - **Gene:** ERCC8. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000082:1-12. **Detection Rate:** East Asian 78%.
EVC-related Ellis-van Creveld Syndrome - **Gene:** EVC. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_153717:1-21. **Detection Rate:** East Asian 96%.
EVC2-related Ellis-van Creveld Syndrome - **Gene:** EVC2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_147127:1-22. **Detection Rate:** East Asian 98%.
Fabry Disease - **Gene:** GLA. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000169:1-7. **Detection Rate:** East Asian 98%.
Familial Dysautonomia - **Gene:** ELP1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_003640:2-37. **Detection Rate:** East Asian >99%.
Familial Hyperinsulinism, ABCC8-related - **Gene:** ABCC8. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000352:1-39. **Detection Rate:** East Asian >99%.
Familial Hyperinsulinism, KCNJ11-related - **Gene:** KCNJ11. Autosomal Recessive. Sequencing with copy number analysis. **Exon:** NM_000525:1. **Detection Rate:** East Asian >99%.
Familial Mediterranean Fever - **Gene:** MEFV. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000243:1-10. **Detection Rate:** East Asian >99%.
Fanconi Anemia Complementation Group A - **Gene:** FANCA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000135:1-43. **Detection Rate:** East Asian 92%.
Fanconi Anemia, FANCC-related - **Gene:** FANCC. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000136:2-15. **Detection Rate:** East Asian >99%.
FKRP-related Disorders - **Gene:** FKRP. Autosomal Recessive. Sequencing with copy number analysis. **Exon:** NM_024301:4. **Detection Rate:** East Asian >99%.
FKTN-related Disorders - **Gene:** FKTN. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001079802:3-11. **Detection Rate:** East Asian 10%.
Free Sialic Acid Storage Disorders - **Gene:** SLC17A5. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_012434:1-11. **Detection Rate:** East Asian 98%.
Galactokinase Deficiency - **Gene:** GALK1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000154:1-8. **Detection Rate:** East Asian >99%.
Galactosemia - **Gene:** GALT. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000155:1-11. **Detection Rate:** East Asian >99%.
Gamma-sarcoglycanopathy - **Gene:** SGCG. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000231:2-8. **Detection Rate:** East Asian 87%.
Gaucher Disease - **Gene:** GBA. Autosomal Recessive. Analysis of homologous regions. **Variants (10):** D409V, D448H, IVS2+1G>A, L444P, N370S, R463C, R463H, R496H, V394L, p.L29Afs*18. **Detection Rate:** East Asian 60%.
GJB2-related DFNB1 Nonsyndromic Hearing Loss and Deafness - **Gene:** GJB2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_004004:1-2. **Detection Rate:** East Asian >99%.
GLB1-related Disorders - **Gene:** GLB1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000404:1-16. **Detection Rate:** East Asian >99%.
Glutaric Acidemia, GCDH-related - **Gene:** GCDH. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000159:2-12. **Detection Rate:** East Asian >99%.
Glycine Encephalopathy, AMT-related - **Gene:** AMT. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000481:1-9. **Detection Rate:** East Asian >99%.
Glycine Encephalopathy, GLDC-related - **Gene:** GLDC. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000170:1-25. **Detection Rate:** East Asian 94%.
Glycogen Storage Disease Type Ia - **Gene:** G6PC1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000151:1-5. **Detection Rate:** East Asian 98%.
Glycogen Storage Disease Type Ib - **Gene:** SLC37A4. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001164277 3-11. **Detection Rate:** East Asian >99%.
Glycogen Storage Disease Type III - **Gene:** AGL. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000642:2-34. **Detection Rate:** East Asian >99%.
GNE Myopathy - **Gene:** GNE. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001128227:1-12. **Detection Rate:** East Asian >99%.
GNPTAB-related Disorders - **Gene:** GNPTAB. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_024312:1-21. **Detection Rate:** East Asian 92%.
HADHA-related Disorders - **Gene:** HADHA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000182:1-20. **Detection Rate:** East Asian >99%.
Hb Beta Chain-related Hemoglobinopathy (Including Beta Thalassemia and Sickle Cell Disease) - **Gene:** HBB. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000518:1-3. **Detection Rate:** East Asian >99%.
Hereditary Fructose Intolerance - **Gene:** ALDOB. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000035:2-9. **Detection Rate:** East Asian >99%.
Hexosaminidase A Deficiency (Including Tay-Sachs Disease) - **Gene:** HEXA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000520:1-14. **Detection Rate:** East Asian >99%.
HMG-CoA Lyase Deficiency - **Gene:** HMGCL. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000191:1-9. **Detection Rate:** East Asian >99%.
Holocarboxylase Synthetase Deficiency - **Gene:** HLCS. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000411:4-12. **Detection Rate:** East Asian >99%.
Homocystinuria, CBS-related - **Gene:** CBS. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000071:3-17. **Detection Rate:** East Asian >99%.
Hydrolethalus Syndrome - **Gene:** HYLS1. Autosomal Recessive. Sequencing with copy number analysis. **Exon:** NM_145014:4. **Detection Rate:** East Asian >99%.
Hypophosphatasia - **Gene:** ALPL. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000478:2-12. **Detection Rate:** East Asian >99%.
Isovaleric Acidemia - **Gene:** IVD. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_002225:1-12. **Detection Rate:** East Asian >99%.

Copyright 2021 Myriad Women's Health, Inc. All rights reserved.

Joubert Syndrome 2 - **Gene:** TMEM216. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001173990:1-5. **Detection Rate:** East Asian >99%.
Junctional Epidermolysis Bullosa, LAMA3-related - **Gene:** LAMA3. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000227:1-38. **Detection Rate:** East Asian >99%.
Junctional Epidermolysis Bullosa, LAMB3-related - **Gene:** LAMB3. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000228:2-23. **Detection Rate:** East Asian >99%.
Junctional Epidermolysis Bullosa, LAMC2-related - **Gene:** LAMC2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_005562:1-23. **Detection Rate:** East Asian >99%.
Krabbe Disease - **Gene:** GALC. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000153:1-17. **Detection Rate:** East Asian >99%.
Leigh Syndrome, French-Canadian Type - **Gene:** LRPPRC. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_133259:1-38. **Detection Rate:** East Asian >99%.
Lipoid Congenital Adrenal Hyperplasia - **Gene:** STAR. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000349:1-7. **Detection Rate:** East Asian >99%.
Lysosomal Acid Lipase Deficiency - **Gene:** LIPA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000235:2-10. **Detection Rate:** East Asian 98%.
Maple Syrup Urine Disease Type Ia - **Gene:** BCKDHA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000709:1-9. **Detection Rate:** East Asian >99%.
Maple Syrup Urine Disease Type Ib - **Gene:** BCKDHB. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_183050:1-10. **Detection Rate:** East Asian >99%.
Maple Syrup Urine Disease Type II - **Gene:** DBT. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001918:1-11. **Detection Rate:** East Asian 97%.
Medium Chain Acyl-CoA Dehydrogenase Deficiency - **Gene:** ACADM. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000016:1-12. **Detection Rate:** East Asian >99%.
Megalencephalic Leukoencephalopathy with Subcortical Cysts - **Gene:** MLC1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_015166 2-12. **Detection Rate:** East Asian >99%.
Metachromatic Leukodystrophy - **Gene:** ARSA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000487:1-8. **Detection Rate:** East Asian >99%.
Methylmalonic Acidemia, cblA Type - **Gene:** MMAA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_172250:2-7. **Detection Rate:** East Asian >99%.
Methylmalonic Acidemia, cblB Type - **Gene:** MMAB. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_052845:1-9. **Detection Rate:** East Asian >99%.
Methylmalonic Acidemia, MMUT-related - **Gene:** MMUT. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000255:2-13. **Detection Rate:** East Asian >99%.
Methylmalonic Aciduria and Homocystinuria, cblC Type - **Gene:** MMACHC. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_015506:1-4. **Detection Rate:** East Asian >99%.
MKS1-related Disorders - **Gene:** MKS1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_017777:1-18. **Detection Rate:** East Asian >99%.
Mucolipidosis III Gamma - **Gene:** GNPTG. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_032520:1-11. **Detection Rate:** East Asian 98%.
Mucolipidosis IV - **Gene:** MCOLN1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_020533:1-14. **Detection Rate:** East Asian >99%.
Mucopolysaccharidosis Type I - **Gene:** IDUA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000203:1-14. **Detection Rate:** East Asian >99%.
Mucopolysaccharidosis Type II - **Gene:** IDS. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000202:1-9. **Detection Rate:** East Asian 89%.
Mucopolysaccharidosis Type IIIA - **Gene:** SGSH. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000199:1-8. **Detection Rate:** East Asian >99%.
Mucopolysaccharidosis Type IIIB - **Gene:** NAGLU. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000263:1-6. **Detection Rate:** East Asian >99%.
Mucopolysaccharidosis Type IIIC - **Gene:** HGSNAT. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_152419:1-18. **Detection Rate:** East Asian >99%.
Muscular Dystrophy, LAMA2-related - **Gene:** LAMA2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000426:1-43,45-65. **Detection Rate:** East Asian 98%.
MYO7A-related Disorders - **Gene:** MYO7A. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000260:2-49. **Detection Rate:** East Asian >99%.
NEB-related Nemaline Myopathy - **Gene:** NEB. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001271208:3-80,117-183. **Detection Rate:** East Asian 92%.
Nephrotic Syndrome, NPHS1-related - **Gene:** NPHS1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_004646:1-29. **Detection Rate:** East Asian >99%.
Nephrotic Syndrome, NPHS2-related - **Gene:** NPHS2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_014625:1-8. **Detection Rate:** East Asian >99%.
Neuronal Ceroid Lipofuscinosis, CLN6-related - **Gene:** CLN6. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_017882:1-7. **Detection Rate:** East Asian >99%.
Niemann-Pick Disease Type C1 - **Gene:** NPC1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000271:1-25. **Detection Rate:** East Asian >99%.
Niemann-Pick Disease Type C2 - **Gene:** NPC2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_006432:1-5. **Detection Rate:** East Asian >99%.
Niemann-Pick Disease, SMPD1-related - **Gene:** SMPD1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000543:1-6. **Detection Rate:** East Asian >99%.
Nijmegen Breakage Syndrome - **Gene:** NBN. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_002485:1-16. **Detection Rate:** East Asian >99%.
Ornithine Transcarbamylase Deficiency - **Gene:** OTC. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000531:1-10. **Detection Rate:** East Asian 97%.
PCCA-related Propionic Acidemia - **Gene:** PCCA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000282:1-24. **Detection Rate:** East Asian 95%.
PCCB-related Propionic Acidemia - **Gene:** PCCB. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000532:1-15. **Detection Rate:** East Asian >99%.
PCDH15-related Disorders - **Gene:** PCDH15. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_033056:2-33. **Detection Rate:** East Asian 93%.
Pendred Syndrome - **Gene:** SLC26A4. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000441:2-21. **Detection Rate:** East Asian >99%.
Peroxisome Biogenesis Disorder Type 1 - **Gene:** PEX1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000466:1-24. **Detection Rate:** East Asian >99%.
Peroxisome Biogenesis Disorder Type 3 - **Gene:** PEX12. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000286:1-3. **Detection Rate:** East Asian >99%.
Peroxisome Biogenesis Disorder Type 4 - **Gene:** PEX6. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000287:1-17. **Detection Rate:** East Asian 97%.

Copyright 2021 Myriad Women's Health, Inc. All rights reserved.

Peroxisome Biogenesis Disorder Type 5 - **Gene:** PEX2. Autosomal Recessive. Sequencing with copy number analysis. **Exon:** NM_000318:4. **Detection Rate:** East Asian >99%.
Peroxisome Biogenesis Disorder Type 6 - **Gene:** PEX10. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_153818:1-6. **Detection Rate:** East Asian >99%.
Phenylalanine Hydroxylase Deficiency - **Gene:** PAH. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000277:1-13. **Detection Rate:** East Asian >99%.
POMGNT-related Disorders - **Gene:** POMGNT1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_017739:2-22. **Detection Rate:** East Asian 96%.
Pompe Disease - **Gene:** GAA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000152:2-20. **Detection Rate:** East Asian >99%.
PPT1-related Neuronal Ceroid Lipofuscinosis - **Gene:** PPT1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000310:1-9. **Detection Rate:** East Asian >99%.
Primary Carnitine Deficiency - **Gene:** SLC22A5. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_003060:1-10. **Detection Rate:** East Asian >99%.
Primary Hyperoxaluria Type 1 - **Gene:** AGXT. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000030:1-11. **Detection Rate:** East Asian >99%.
Primary Hyperoxaluria Type 2 - **Gene:** GRHPR. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_012203:1-9. **Detection Rate:** East Asian >99%.
Primary Hyperoxaluria Type 3 - **Gene:** HOGA1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_138413:1-7. **Detection Rate:** East Asian >99%.
Pycnodysostosis - **Gene:** CTSK. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000396:2-8. **Detection Rate:** East Asian >99%.
Pyruvate Carboxylase Deficiency - **Gene:** PC. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000920:3-22. **Detection Rate:** East Asian >99%.
Rhizomelic Chondrodysplasia Punctata Type 1 - **Gene:** PEX7. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000288:1-10. **Detection Rate:** East Asian >99%.
RTEL1-related Disorders - **Gene:** RTEL1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_032957:2-35. **Detection Rate:** East Asian >99%.
Sandhoff Disease - **Gene:** HEXB. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000521:1-14. **Detection Rate:** East Asian 98%.
Short-chain Acyl-CoA Dehydrogenase Deficiency - **Gene:** ACADS. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000017:1-10. **Detection Rate:** East Asian >99%.
Sjogren-Larsson Syndrome - **Gene:** ALDH3A2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000382:1-10. **Detection Rate:** East Asian 96%.
SLC26A2-related Disorders - **Gene:** SLC26A2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000112:2-3. **Detection Rate:** East Asian >99%.
Smith-Lemli-Opitz Syndrome - **Gene:** DHCR7. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001360:3-9. **Detection Rate:** East Asian >99%.
Spastic Paraplegia Type 15 - **Gene:** ZFYVE26. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_015346:2-42. **Detection Rate:** East Asian >99%.
Spinal Muscular Atrophy - **Gene:** SMN1. Autosomal Recessive. Spinal muscular atrophy. **Variant (1):** SMN1 copy number. **Detection Rate:** East Asian 93%.
Spondylothoracic Dysostosis - **Gene:** MESP2. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_001039958:1-2. **Detection Rate:** East Asian >99%.
TGM1-related Autosomal Recessive Congenital Ichthyosis - **Gene:** TGM1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000359 2-15. **Detection Rate:** East Asian >99%.
TPP1-related Neuronal Ceroid Lipofuscinosis - **Gene:** TPP1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000391:1-13. **Detection Rate:** East Asian >99%.
Tyrosine Hydroxylase Deficiency - **Gene:** TH. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_199292:1-14. **Detection Rate:** East Asian >99%.
Tyrosinemia Type I - **Gene:** FAH. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000137:1-14. **Detection Rate:** East Asian >99%.
Tyrosinemia Type II - **Gene:** TAT. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000353:2-12. **Detection Rate:** East Asian >99%.
USH1C-related Disorders - **Gene:** USH1C. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_005709:1-21. **Detection Rate:** East Asian >99%.
USH2A-related Disorders - **Gene:** USH2A. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_206933:2-72. **Detection Rate:** East Asian 98%.
Usher Syndrome Type 3 - **Gene:** CLRN1. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_174878:1-3. **Detection Rate:** East Asian >99%.
Very-long-chain Acyl-CoA Dehydrogenase Deficiency - **Gene:** ACADVL. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000018:1-20. **Detection Rate:** East Asian >99%.
Wilson Disease - **Gene:** ATP7B. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000053:1-21. **Detection Rate:** East Asian >99%.
X-linked Adrenal Hypoplasia Congenita - **Gene:** NR0B1. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000475:1-2. **Detection Rate:** East Asian 97%.
X-linked Adrenoleukodystrophy - **Gene:** ABCD1. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000033:1-6. **Detection Rate:** East Asian 77%.
X-linked Alport Syndrome - **Gene:** COL4A5. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000495:1-51. **Detection Rate:** East Asian 96%.
X-linked Juvenile Retinoschisis - **Gene:** RS1. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000330:1-6. **Detection Rate:** East Asian 98%.
X-linked Myotubular Myopathy - **Gene:** MTM1. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000252:2-15. **Detection Rate:** East Asian 96%.
X-linked Severe Combined Immunodeficiency - **Gene:** IL2RG. X-linked Recessive. Sequencing with copy number analysis. **Exons:** NM_000206:1-8. **Detection Rate:** East Asian >99%.
Xeroderma Pigmentosum Group A - **Gene:** XPA. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_000380:1-6. **Detection Rate:** East Asian >99%.
Xeroderma Pigmentosum Group C - **Gene:** XPC. Autosomal Recessive. Sequencing with copy number analysis. **Exons:** NM_004628:1-16. **Detection Rate:** East Asian 97%.

Copyright 2021 Myriad Women's Health, Inc. All rights reserved.

Risk Calculations

Below are the risk calculations for all conditions tested. Negative results do not rule out the possibility of being a carrier. Residual risk is an estimate of each patient's post-test likelihood of being a carrier, while the reproductive risk represents an estimated likelihood that the patients' future children could inherit each disease. These risks are inherent to all carrier-screening tests, may vary by ethnicity, are predicated on a negative family history, and are present even given a negative test result. Inaccurate reporting of ethnicity may cause errors in risk calculation. In addition, average carrier rates are estimated using incidence or prevalence data from published scientific literature and/or reputable databases, where available, and are incorporated into residual risk calculations for each population/ethnicity. When population-specific data is not available for a condition, average worldwide incidence or prevalence is used. Further, incidence and prevalence data are only collected for the specified phenotypes (which include primarily the classic or severe forms of disease) and may not include alternate or milder disease manifestations associated with the gene. Actual incidence rates, prevalence rates, and carrier rates, and therefore actual residual risks, may be higher or lower than the estimates provided. Carrier rates, incidence/prevalence, and/or residual risks are not provided for some genes with biological or heritable properties that would make these estimates inaccurate. A '†' symbol indicates a positive result. See the full clinical report for interpretation and details. The reproductive risk presented is based on a hypothetical pairing with a partner of the same ethnic group.

Disease	DONOR 16010 Residual Risk	Reproductive Risk
6-pyruvoyl-tetrahydropterin Synthase Deficiency	1 in 24,000	< 1 in 1,000,000
Adenosine Deaminase Deficiency	1 in 22,000	< 1 in 1,000,000
Alpha Thalassemia, HBA1/HBA2-related	Alpha globin status: aa/aa.	Not calculated
Alpha-mannosidosis	1 in 35,000	< 1 in 1,000,000
Alpha-sarcoglycanopathy	1 in 34,000	< 1 in 1,000,000
Alstrom Syndrome	< 1 in 50,000	< 1 in 1,000,000
Andermann Syndrome	< 1 in 50,000	< 1 in 1,000,000
Argininemia	1 in 6,300	< 1 in 1,000,000
Argininosuccinic Aciduria	1 in 45,000	< 1 in 1,000,000
Aspartylglucosaminuria	< 1 in 50,000	< 1 in 1,000,000
Ataxia with Vitamin E Deficiency	< 1 in 50,000	< 1 in 1,000,000
Ataxia-telangiectasia	1 in 4,200	< 1 in 1,000,000
ATP7A-related Disorders	< 1 in 1,000,000	1 in 410,000
Autoimmune Polyglandular Syndrome Type 1	1 in 18,000	< 1 in 1,000,000
Autosomal Recessive Osteopetrosis Type 1	1 in 8,900	< 1 in 1,000,000
Autosomal Recessive Polycystic Kidney Disease, PKHD1-related	1 in 8,100	< 1 in 1,000,000
Autosomal Recessive Spastic Ataxia of Charlevoix-Saguenay	< 1 in 44,000	< 1 in 1,000,000
Bardet-Biedl Syndrome, BBS1-related	< 1 in 50,000	< 1 in 1,000,000
Bardet-Biedl Syndrome, BBS10-related	< 1 in 50,000	< 1 in 1,000,000
Bardet-Biedl Syndrome, BBS12-related	< 1 in 50,000	< 1 in 1,000,000
Bardet-Biedl Syndrome, BBS2-related	< 1 in 50,000	< 1 in 1,000,000
BCS1L-related Disorders	< 1 in 50,000	< 1 in 1,000,000
Beta-sarcoglycanopathy	1 in 39,000	< 1 in 1,000,000
Biotinidase Deficiency	1 in 67,000	< 1 in 1,000,000
Bloom Syndrome	< 1 in 50,000	< 1 in 1,000,000
Calpainopathy	1 in 11,000	< 1 in 1,000,000
Canavan Disease	1 in 9,700	< 1 in 1,000,000
Carbamoylphosphate Synthetase I Deficiency	1 in 45,000	< 1 in 1,000,000
Carnitine Palmitoyltransferase IA Deficiency	< 1 in 50,000	< 1 in 1,000,000
Carnitine Palmitoyltransferase II Deficiency	1 in 31,000	< 1 in 1,000,000
Cartilage-hair Hypoplasia	< 1 in 50,000	< 1 in 1,000,000
Cerebrotendinous Xanthomatosis	1 in 13,000	< 1 in 1,000,000
Citrullinemia Type 1	1 in 700	1 in 270,000
CLN3-related Neuronal Ceroid Lipofuscinosis	1 in 13,000	< 1 in 1,000,000
CLN5-related Neuronal Ceroid Lipofuscinosis	< 1 in 50,000	< 1 in 1,000,000
CLN8-related Neuronal Ceroid Lipofuscinosis	< 1 in 50,000	< 1 in 1,000,000
Cohen Syndrome	< 1 in 15,000	< 1 in 1,000,000
COL4A3-related Alport Syndrome	1 in 5,800	< 1 in 1,000,000
COL4A4-related Alport Syndrome	1 in 35,000	< 1 in 1,000,000
Combined Pituitary Hormone Deficiency, PROP1-related	1 in 6,100	< 1 in 1,000,000
Congenital Adrenal Hyperplasia, CYP11B1-related	1 in 8,400	< 1 in 1,000,000
Congenital Adrenal Hyperplasia, CYP21A2-related	1 in 550	1 in 140,000
Congenital Disorder of Glycosylation Type Ia	1 in 16,000	< 1 in 1,000,000
Congenital Disorder of Glycosylation Type Ic	< 1 in 50,000	< 1 in 1,000,000
Congenital Disorder of Glycosylation, MPI-related	< 1 in 50,000	< 1 in 1,000,000

Copyright 2021 Myriad Women's Health, Inc. All rights reserved.

Disease	DONOR 16010 Residual Risk	Reproductive Risk
Costeff Optic Atrophy Syndrome	< 1 in 50,000	< 1 in 1,000,000
Cystic Fibrosis	1 in 9,000	< 1 in 1,000,000
Cystinosis	1 in 22,000	< 1 in 1,000,000
D-bifunctional Protein Deficiency	1 in 9,000	< 1 in 1,000,000
Delta-sarcoglycanopathy	< 1 in 13,000	< 1 in 1,000,000
Dihydrolipoamide Dehydrogenase Deficiency	< 1 in 50,000	< 1 in 1,000,000
Dysferlinopathy	1 in 11,000	< 1 in 1,000,000
Dystrophinopathy (Including Duchenne/Becker Muscular Dystrophy)	Not calculated	Not calculated
ERCC6-related Disorders	1 in 8,400	< 1 in 1,000,000
ERCC8-related Disorders	< 1 in 2,300	< 1 in 1,000,000
EVC-related Ellis-van Creveld Syndrome	1 in 7,800	< 1 in 1,000,000
EVC2-related Ellis-van Creveld Syndrome	1 in 9,800	< 1 in 1,000,000
Fabry Disease	< 1 in 1,000,000	1 in 80,000
Familial Dysautonomia	< 1 in 50,000	< 1 in 1,000,000
Familial Hyperinsulinism, ABCC8-related	1 in 14,000	< 1 in 1,000,000
Familial Hyperinsulinism, KCNJ11-related	1 in 42,000	< 1 in 1,000,000
Familial Mediterranean Fever	< 1 in 50,000	< 1 in 1,000,000
Fanconi Anemia Complementation Group A	1 in 3,100	< 1 in 1,000,000
Fanconi Anemia, FANCC-related	< 1 in 50,000	< 1 in 1,000,000
FKRP-related Disorders	1 in 32,000	< 1 in 1,000,000
FKTN-related Disorders	1 in 110	1 in 40,000
Free Sialic Acid Storage Disorders	< 1 in 30,000	< 1 in 1,000,000
Galactokinase Deficiency	< 1 in 50,000	< 1 in 1,000,000
Galactosemia	1 in 32,000	< 1 in 1,000,000
Gamma-sarcoglycanopathy	< 1 in 3,800	< 1 in 1,000,000
Gaucher Disease	1 in 450	1 in 320,000
GJB2-related DFNB1 Nonsyndromic Hearing Loss and Deafness	1 in 3,300	1 in 450,000
GLB1-related Disorders	1 in 17,000	< 1 in 1,000,000
Glutaric Acidemia, GCDH-related	1 in 13,000	< 1 in 1,000,000
Glycine Encephalopathy, AMT-related	1 in 26,000	< 1 in 1,000,000
Glycine Encephalopathy, GLDC-related	1 in 4,200	< 1 in 1,000,000
Glycogen Storage Disease Type Ia	1 in 8,700	< 1 in 1,000,000
Glycogen Storage Disease Type Ib	1 in 35,000	< 1 in 1,000,000
Glycogen Storage Disease Type III	1 in 16,000	< 1 in 1,000,000
GNE Myopathy	< 1 in 50,000	< 1 in 1,000,000
GNPTAB-related Disorders	1 in 2,600	< 1 in 1,000,000
HADHA-related Disorders	1 in 25,000	< 1 in 1,000,000
Hb Beta Chain-related Hemoglobinopathy (Including Beta Thalassemia and Sickle Cell Disease)	1 in 900	1 in 34,000
Hereditary Fructose Intolerance	1 in 7,900	< 1 in 1,000,000
Hexosaminidase A Deficiency (Including Tay-Sachs Disease)	1 in 30,000	< 1 in 1,000,000
HMG-CoA Lyase Deficiency	< 1 in 50,000	< 1 in 1,000,000
Holocarboxylase Synthetase Deficiency	1 in 16,000	< 1 in 1,000,000
Homocystinuria, CBS-related	< 1 in 50,000	< 1 in 1,000,000
Hydrolethalus Syndrome	< 1 in 50,000	< 1 in 1,000,000
Hypophosphatasia	1 in 24,000	< 1 in 1,000,000
Isovaleric Acidemia	1 in 39,000	< 1 in 1,000,000
Joubert Syndrome 2	< 1 in 50,000	< 1 in 1,000,000
Junctional Epidermolysis Bullosa, LAMA3-related	< 1 in 50,000	< 1 in 1,000,000
Junctional Epidermolysis Bullosa, LAMB3-related	1 in 31,000	< 1 in 1,000,000
Junctional Epidermolysis Bullosa, LAMC2-related	< 1 in 50,000	< 1 in 1,000,000
Krabbe Disease	1 in 17,000	< 1 in 1,000,000
Leigh Syndrome, French-Canadian Type	< 1 in 50,000	< 1 in 1,000,000
Lipoid Congenital Adrenal Hyperplasia	1 in 40,000	< 1 in 1,000,000
Lysosomal Acid Lipase Deficiency	< 1 in 34,000	< 1 in 1,000,000
Maple Syrup Urine Disease Type Ia	1 in 43,000	< 1 in 1,000,000
Maple Syrup Urine Disease Type Ib	1 in 23,000	< 1 in 1,000,000
Maple Syrup Urine Disease Type II	1 in 15,000	< 1 in 1,000,000
Medium Chain Acyl-CoA Dehydrogenase Deficiency	1 in 11,000	< 1 in 1,000,000
Megalencephalic Leukoencephalopathy with Subcortical Cysts	< 1 in 50,000	< 1 in 1,000,000
Metachromatic Leukodystrophy	1 in 16,000	< 1 in 1,000,000
Methylmalonic Acidemia, cblA Type	< 1 in 50,000	< 1 in 1,000,000

Copyright 2021 Myriad Women's Health, Inc. All rights reserved.

Disease	DONOR 16010 Residual Risk	Reproductive Risk
Methylmalonic Acidemia, cblB Type	< 1 in 50,000	< 1 in 1,000,000
Methylmalonic Acidemia, MMUT-related	1 in 12,000	< 1 in 1,000,000
Methylmalonic Aciduria and Homocystinuria, cblC Type	1 in 33,000	< 1 in 1,000,000
MKS1-related Disorders	< 1 in 50,000	< 1 in 1,000,000
Mucolipidosis III Gamma	< 1 in 20,000	< 1 in 1,000,000
Mucolipidosis IV	< 1 in 50,000	< 1 in 1,000,000
Mucopolysaccharidosis Type I	1 in 16,000	< 1 in 1,000,000
Mucopolysaccharidosis Type II	1 in 520,000	1 in 120,000
Mucopolysaccharidosis Type IIIA	1 in 16,000	< 1 in 1,000,000
Mucopolysaccharidosis Type IIIB	1 in 42,000	< 1 in 1,000,000
Mucopolysaccharidosis Type IIIC	< 1 in 50,000	< 1 in 1,000,000
Muscular Dystrophy, LAMA2-related	1 in 5,700	< 1 in 1,000,000
MYO7A-related Disorders	1 in 15,000	< 1 in 1,000,000
NEB-related Nemaline Myopathy	1 in 1,200	1 in 400,000
Nephrotic Syndrome, NPHS1-related	< 1 in 50,000	< 1 in 1,000,000
Nephrotic Syndrome, NPHS2-related	1 in 35,000	< 1 in 1,000,000
Neuronal Ceroid Lipofuscinosis, CLN6-related	< 1 in 50,000	< 1 in 1,000,000
Niemann-Pick Disease Type C1	1 in 17,000	< 1 in 1,000,000
Niemann-Pick Disease Type C2	< 1 in 50,000	< 1 in 1,000,000
Niemann-Pick Disease, SMPD1-related	1 in 25,000	< 1 in 1,000,000
Nijmegen Breakage Syndrome	< 1 in 50,000	< 1 in 1,000,000
Ornithine Transcarbamylase Deficiency	< 1 in 1,000,000	1 in 140,000
PCCA-related Propionic Acidemia	1 in 4,200	< 1 in 1,000,000
PCCB-related Propionic Acidemia	1 in 6,500	< 1 in 1,000,000
PCDH15-related Disorders	1 in 3,300	< 1 in 1,000,000
Pendred Syndrome	1 in 6,400	< 1 in 1,000,000
Peroxisome Biogenesis Disorder Type 1	1 in 16,000	< 1 in 1,000,000
Peroxisome Biogenesis Disorder Type 3	< 1 in 50,000	< 1 in 1,000,000
Peroxisome Biogenesis Disorder Type 4	1 in 9,300	< 1 in 1,000,000
Peroxisome Biogenesis Disorder Type 5	< 1 in 71,000	< 1 in 1,000,000
Peroxisome Biogenesis Disorder Type 6	< 1 in 50,000	< 1 in 1,000,000
Phenylalanine Hydroxylase Deficiency	1 in 7,700	< 1 in 1,000,000
POMGNT-related Disorders	< 1 in 12,000	< 1 in 1,000,000
Pompe Disease	1 in 10,000	< 1 in 1,000,000
PPT1-related Neuronal Ceroid Lipofuscinosis	1 in 7,700	< 1 in 1,000,000
Primary Carnitine Deficiency	1 in 10,000	< 1 in 1,000,000
Primary Hyperoxaluria Type 1	1 in 13,000	< 1 in 1,000,000
Primary Hyperoxaluria Type 2	< 1 in 50,000	< 1 in 1,000,000
Primary Hyperoxaluria Type 3	1 in 20,000	< 1 in 1,000,000
Pycnodysostosis	1 in 43,000	< 1 in 1,000,000
Pyruvate Carboxylase Deficiency	1 in 25,000	< 1 in 1,000,000
Rhizomelic Chondrodysplasia Punctata Type 1	1 in 16,000	< 1 in 1,000,000
RTEL1-related Disorders	< 1 in 50,000	< 1 in 1,000,000
Sandhoff Disease	1 in 18,000	< 1 in 1,000,000
Short-chain Acyl-CoA Dehydrogenase Deficiency	1 in 9,700	< 1 in 1,000,000
Sjogren-Larsson Syndrome	< 1 in 12,000	< 1 in 1,000,000
SLC26A2-related Disorders	1 in 16,000	< 1 in 1,000,000
Smith-Lemli-Opitz Syndrome	< 1 in 50,000	< 1 in 1,000,000
Spastic Paraplegia Type 15	< 1 in 50,000	< 1 in 1,000,000
Spinal Muscular Atrophy	Negative for g.27134T>G SNP SMN1: 2 copies 1 in 700	1 in 150,000
Spondylothoracic Dysostosis	< 1 in 50,000	< 1 in 1,000,000
TGM1-related Autosomal Recessive Congenital Ichthyosis	1 in 22,000	< 1 in 1,000,000
TPP1-related Neuronal Ceroid Lipofuscinosis	1 in 30,000	< 1 in 1,000,000
Tyrosine Hydroxylase Deficiency	< 1 in 50,000	< 1 in 1,000,000
Tyrosinemia Type I	1 in 16,000	< 1 in 1,000,000
Tyrosinemia Type II	1 in 25,000	< 1 in 1,000,000
USH1C-related Disorders	1 in 44,000	< 1 in 1,000,000
USH2A-related Disorders	1 in 5,900	< 1 in 1,000,000
Usher Syndrome Type 3	1 in 41,000	< 1 in 1,000,000
Very-long-chain Acyl-CoA Dehydrogenase Deficiency	1 in 12,000	< 1 in 1,000,000

Copyright 2021 Myriad Women's Health, Inc. All rights reserved.

Disease	DONOR 16010 Residual Risk	Reproductive Risk
Wilson Disease	NM_000053.3(ATP7B):c.2621C>T(A874V) heterozygote †	1 in 260
X-linked Adrenal Hypoplasia Congenita	< 1 in 1,000,000	< 1 in 1,000,000
X-linked Adrenoleukodystrophy	1 in 170,000	1 in 80,000
X-linked Alport Syndrome	Not calculated	Not calculated
X-linked Juvenile Retinoschisis	< 1 in 1,000,000	1 in 50,000
X-linked Myotubular Myopathy	Not calculated	Not calculated
X-linked Severe Combined Immunodeficiency	< 1 in 1,000,000	1 in 200,000
Xeroderma Pigmentosum Group A	1 in 10,000	< 1 in 1,000,000
Xeroderma Pigmentosum Group C	1 in 7,300	< 1 in 1,000,000

Copyright 2021 Myriad Women's Health, Inc. All rights reserved.

Patient Information	
Patient Name:	Donor 16010
Date Of Birth:	[redacted]
Gender:	Male
Patient ID:	N/A
Medical Record #:	N/A
Collection Kit:	40845274-2-C
Accession ID:	N/A
Case File ID:	21180696
Ethnicity:	Other

Test Information	
Ordering Physician:	Dr. James Kuan
Clinic Information:	Seattle Sperm Bank
Phone:	(206) 588-1484
Report Date:	03/12/2026
Sample Collected:	02/26/2026
Sample Received:	03/03/2026
Sample Type:	Saliva

natera | Horizon™ Advanced carrier screening

CARRIER SCREENING REPORT

ABOUT THIS SCREEN: Horizon™ is a carrier screen for specific autosomal recessive and X-linked diseases. This information can help patients learn their risk of having a child with specific genetic conditions.

ORDER SELECTED: The Horizon Custom panel was ordered for this patient. Males are not screened for X-linked diseases

FINAL RESULTS SUMMARY:

Negative for 1 out of 1 diseases

No Pathogenic variants were detected in the genes that were screened. The patient's remaining carrier risk after the negative screening results is listed for each disease/gene on the Horizon website at https://www.natera.com/panel-option/h-all/. Please see the following pages of this report for a comprehensive list of all conditions included on this individual's screen.

Carrier screening is not diagnostic and may not detect all possible pathogenic variants in a given gene.

RECOMMENDATIONS

Individuals who would like to review their Horizon report with a Natera Laboratory Genetic Counselor may schedule a telephone genetic information session by calling 650-249-9090 or visiting naterasession.com. Clinicians with questions may contact Natera at 650-249-9090 or email support@natera.com.

Christine M. Eng, M.D.
Medical Director, Baylor Genetics

Linyan Meng, Ph.D.
Laboratory Director, Baylor Genetics

J. Dianne Keen-Kim, Ph.D., FACMGG
Senior Laboratory Director, Natera

Yang Wang, Ph.D., FACMGG
Laboratory Director, Natera

The pre-analytic and post-analytic phases of this test were performed by NSTX, Inc., 13011 McCallen Pass, Building A Suite 110, Austin, TX 78753 (CLIA ID 45D2093704). This test was performed by Baylor Miraca Genetics, DBA Baylor Genetics, 2450 Holcombe Blvd. Houston, TX 77021 (CLIA ID 45D0660090). The performance characteristics of this test were developed by Baylor Miraca Genetics, DBA Baylor Genetics (CLIA ID 45D0660090). This test has not been cleared or approved by the U.S. Food and Drug Administration (FDA). These laboratories are regulated under CLIA as qualified to perform high-complexity testing. © Natera, Inc. 2022. All Rights Reserved.

natera

Patient Information
Patient Name: Donor 16010

Date Of Birth: [redacted]
Case File ID: 21180696

Test Information
Ordering Physician: Dr. James Kuan

Clinic Information: Seattle Sperm Bank

Report Date: 03/12/2026

DISEASES SCREENED

Below is a list of all diseases screened and the result. Certain conditions have unique patient-specific numerical values, therefore, results for those conditions are formatted differently.

Autosomal Recessive

F
FRIEDREICH ATAXIA (*FXN*) **negative**

Testing Methodology, Limitations, and Comments:

Next-generation sequencing (NGS)

Sequencing library prepared from genomic DNA isolated from a patient sample is enriched for targets of interest using standard hybridization capture protocols and PCR amplification (for targets specified below). NGS is then performed to achieve the standards of quality control metrics, including a minimum coverage of 99% of targeted regions at 20X sequencing depth. Sequencing data is aligned to human reference sequence, followed by deduplication, metric collection and variant calling (coding region +/- 20bp). Variants are then classified according to ACMGG/AMP standards of interpretation using publicly available databases including but not limited to ENSEMBL, HGMD Pro, ClinGen, ClinVar, 1000G, ESP and gnomAD. Variants predicted to be pathogenic or likely pathogenic for the specified diseases are reported. It should be noted that the data interpretation is based on our current understanding of the genes and variants at the time of reporting. Putative positive sequencing variants that do not meet internal quality standards or are within highly homologous regions are confirmed by Sanger sequencing or gene-specific long-range PCR as needed prior to reporting.

Copy Number Variant (CNV) analysis is limited to deletions involving two or more exons for all genes on the panel, in addition to specific known recurrent single-exon deletions. CNVs of small size may have reduced detection rate. This method does not detect gene inversions, single-exonic and sub-exonic deletions (unless otherwise specified), and duplications of all sizes (unless otherwise specified). Additionally, this method does not define the exact breakpoints of detected CNV events. Confirmation testing for copy number variation is performed by specific PCR, Multiplex Ligation-dependent Probe Amplification (MLPA), next generation sequencing, or other methodology.

This test may not detect certain variants due to local sequence characteristics, high/low genomic complexity, homologous sequence, or allele dropout (PCR-based assays). Variants within noncoding regions (promoter, 5'UTR, 3'UTR, deep intronic regions, unless otherwise specified), small deletions or insertions larger than 25bp, low-level mosaic variants, structural variants such as inversions, and/or balanced translocations may not be detected with this technology.

Friedreich Ataxia (FXN)

The GAA repeat region of the FXN gene is assessed by trinucleotide PCR assay and capillary electrophoresis. Variances of +/-1 repeat for normal alleles and up to +/-3 repeats for premutation alleles may occur. For fully penetrant expanded alleles, the precise repeat size cannot be determined, therefore the approximate allele size is reported. Sequencing and copy number variants are analyzed by next-generation sequencing analysis.

Friedreich Ataxia Repeat Categories

Categories	GAA Repeat Sizes
Normal	<34
Premutation	34 - 65
Full	>65

Variant Classification

Only pathogenic or likely pathogenic variants are reported. Other variants including benign variants, likely benign variants, variants of uncertain significance, or inconclusive variants identified during this analysis may be reported in certain circumstances. Our laboratory's variant classification criteria are based on the ACMG and internal guidelines and our current understanding of the specific genes. This interpretation may change over time as more information about a gene and/or variant becomes available. Natera and its lab partner(s) may reclassify variants at certain intervals but may not release updated reports without a specific request made to Natera by the ordering provider. Natera may disclose incidental findings if deemed clinically pertinent to the test performed.

Negative Results

A negative carrier screening result reduces the risk for a patient to be a carrier of a specific disease but does not completely rule out carrier status. Please visit https://www.natera.com/panel-option/h-all/ for a table of carrier rates, detection rates, residual risks and promised variants/exons per gene. Carrier rates before and after testing vary by ethnicity and assume a negative family history for each disease screened and the absence of clinical symptoms in the patient. Any patient with a family history for a specific genetic disease will have a higher carrier risk prior to testing and, if the disease-causing mutation in their family is not included on the test, their carrier risk would remain unchanged. Genetic counseling is recommended for patients with a family history of genetic disease so that risk figures based on actual family history can be determined and discussed along with potential implications for reproduction. Horizon carrier screening has been developed to identify the reproductive risks for monogenic inherited conditions. Even when one or both members of a couple screen negative for pathogenic variants in a specific gene, the disease risk for their offspring is not zero. There is still a low risk for the condition in their offspring due to a number of different mechanisms that are not detected by Horizon including, but not limited to, pathogenic variant(s) in the tested gene or in a different gene not included on Horizon, pathogenic variant(s) in an upstream regulator, uniparental disomy, de novo mutation(s), or digenic or polygenic inheritance.

Patient Information
Patient Name: Donor 16010

Date Of Birth:
Case File ID: 21180696

Test Information
Ordering Physician: Dr. James Kuan

Clinic Information: Seattle Sperm Bank

Report Date: 03/12/2026

Additional Comments

These analyses generally provide highly accurate information regarding the patient's carrier status. Despite this high level of accuracy, it should be kept in mind that there are many potential sources of diagnostic error, including misidentification of samples, polymorphisms, or other rare genetic variants that interfere with analysis. Families should understand that rare diagnostic errors may occur for these reasons.

15 Crawford St., STE 100
Needham, MA 02494
(p) 626-350-0537
(f) 626-454-1667
Lab Director: Arash Radfar M.D.
CLIA: 22D0957540

fulgent | INFORM DIAGNOSTICS™ A FULGENT GENETICS COMPANY
Metroplex Pathology Associates
Lakewood Pathology Associates
Cohen Dermatopathology
Therapath Partners

Patient Information:
16010, Donor
DOB: [redacted]
Sex: M
MR#:
Patient#: FT-PT8772017

Partner Information:
Not Tested

Physician:
Kuan, James
ATTN: SSB Genetics, Dept
Bellevue Sperm Bank
4915 25th Avenue NE, Ste 204W
Seattle, WA 98105
Phone: (206) 588-1484

Laboratory:
Fulgent Therapeutics LLC
CAP#: 8042697
CLIA#: 05D2043189
Laboratory Director:
Lawrence M. Weiss, MD
Report Date: **Jul 10,2024**

Accession:
FT-7101563
Test#: FT-TS14878861
Specimen Type: Saliva
Collected: Jun 25,2024

Accession:
N/A

FINAL RESULTS

No carrier mutations identified

TEST PERFORMED

Custom Beacon Preconception Carrier Screening Panel
(2 Gene Panel: *CDH23 and SBDS*; gene sequencing with deletion and duplication analysis)

INTERPRETATION:

Notes and Recommendations:

- No carrier mutations were identified in the submitted specimen. A negative result does not rule out the possibility of a genetic predisposition nor does it rule out any pathogenic mutations in areas not assessed by this test or in regions that were covered at a level too low to reliably assess. Also, it does not rule out mutations that are of the sort not queried by this test; see Methods and Limitations for more information. A negative result reduces, but does not eliminate, the chance to be a carrier for any condition included in this screen. Please see the supplemental table for details.
- This carrier screening test does not screen for all possible genetic conditions, nor for all possible mutations in every gene tested. This report does not include variants of uncertain significance; only variants classified as pathogenic or likely pathogenic at the time of testing, and considered relevant for reproductive carrier screening, are reported. Please see the gene specific notes for details. Please note that the classification of variants can change over time.
- Patients may wish to discuss any carrier results with blood relatives, as there is an increased chance that they are also carriers. These results should be interpreted in the context of this individual's clinical findings, biochemical profile, and family history.
- Gene specific notes and limitations may be present. See below.
- Genetic counseling is recommended. Available genetic counselors and additional resources can be found at the National Society of Genetic Counselors (NSGC; https://www.nsgc.org)

15 Crawford St., STE 100
Needham, MA 02494
(p) 626-350-0537
(f) 626-454-1667
Lab Director: Arash Radfar M.D.
CLIA: 22D0957540

GENES TESTED:

Custom Beacon Preconception Carrier Screening Panel - 2 Genes

This analysis was run using the Custom Beacon Preconception Carrier Screening Panel gene list. 2 genes were tested with 100.0% of targets sequenced at >20x coverage. For more gene-specific information and assistance with residual risk calculation, see the SUPPLEMENTAL TABLE.

CDH23, SBDS

METHODS:

Genomic DNA was isolated from the submitted specimen indicated above (if cellular material was submitted). DNA was barcoded, and enriched for the coding exons of targeted genes using hybrid capture technology. Prepared DNA libraries were then sequenced using a Next Generation Sequencing technology. Following alignment to the human genome reference sequence (assembly GRCh37), variants were detected in regions of at least 10x coverage. For this specimen, 100.00% and 100.00% of coding regions and splicing junctions of genes listed had been sequenced with coverage of at least 10x and 20x, respectively, by NGS or by Sanger sequencing. The remaining regions did not have 10x coverage, and were not evaluated. Variants were interpreted manually using locus specific databases, literature searches, and other molecular biological principles. To minimize false positive results, any variants that do not meet internal quality standards are confirmed by Sanger sequencing. Variants classified as pathogenic, likely pathogenic, or risk allele which are located in the coding regions and nearby intronic regions (+/- 20bp) of the genes listed above are reported. Variants outside these intervals may be reported but are typically not guaranteed. When a single pathogenic or likely pathogenic variant is identified in a clinically relevant gene with autosomal recessive inheritance, the laboratory will attempt to ensure 100% coverage of coding sequences either through NGS or Sanger sequencing technologies ("fill-in"). All genes listed were evaluated for large deletions and/or duplications. However, single exon deletions or duplications will not be detected in this assay, nor will copy number alterations in regions of genes with significant pseudogenes. Putative deletions or duplications are analyzed using Fulgent Germline proprietary pipeline for this specimen. Bioinformatics: The Fulgent Germline v2019.2 pipeline was used to analyze this specimen.

LIMITATIONS:

General Limitations

These test results and variant interpretation are based on the proper identification of the submitted specimen, accuracy of any stated familial relationships, and use of the correct human reference sequences at the queried loci. In very rare instances, errors may result due to mix-up or co-mingling of specimens. Positive results do not imply that there are no other contributors, genetic or otherwise, to future pregnancies, and negative results do not rule out the genetic risk to a pregnancy. Official gene names change over time. Fulgent uses the most up to date gene names based on HUGO Gene Nomenclature Committee (https://www.genenames.org) recommendations. If the gene name on report does not match that of ordered gene, please contact the laboratory and details can be provided. Result interpretation is based on the available clinical and family history information for this individual, collected published information, and Alamut annotation available at the time of reporting. This assay is not designed or validated for the detection of low-level mosaicism or somatic mutations. This assay will not detect certain types of genomic aberrations such as translocations, inversions, or repeat expansions other than specified genes. DNA alterations in regulatory regions or deep intronic regions (greater than 20bp from an exon) may not be detected by this test. Unless otherwise indicated, no additional assays have been performed to evaluate genetic changes in this specimen. There are technical limitations on the ability of DNA sequencing to detect small insertions and deletions. Our laboratory uses a sensitive detection algorithm, however these types of alterations are not detected as reliably as single nucleotide variants. Rarely, due to systematic chemical, computational, or human error, DNA variants may be missed. Although next generation sequencing technologies and our bioinformatics analysis significantly reduce the confounding contribution of pseudogene sequences or other highly-homologous sequences, sometimes these may still interfere with the technical ability of the assay to identify pathogenic alterations in both sequencing and deletion/duplication analyses. Deletion/duplication analysis can identify alterations of genomic regions which include one whole gene (buccal swab specimens and whole blood specimens) and are two or more contiguous exons in size (whole blood specimens only); single exon deletions or duplications may occasionally be identified, but are not routinely detected by this test. When novel DNA duplications are identified, it is not possible to discern the genomic location or orientation of the duplicated segment, hence the effect of the duplication cannot be predicted. Where deletions are detected, it is not always possible to determine whether the predicted product will remain in-frame or not. Unless otherwise indicated, deletion/duplication analysis has not been performed in regions that have been sequenced by Sanger.

15 Crawford St., STE 100
Needham, MA 02494
(p) 626-350-0537
(f) 626-454-1667
Lab Director: Arash Radfar M.D.
CLIA: 22D0957540

fulgent | INFORM DIAGNOSTICS™ A FULGENT GENETICS COMPANY
Metroplex Pathology Associates
Lakewood Pathology Associates
Cohen Dermatopathology
Therapath Partners

Gene Specific Notes and Limitations

No gene specific limitations apply to the genes on the tested panel.

SIGNATURE:

Dr. Harry Gao, DABMG, FACMG on **7/10/2024**
Laboratory Director, Fulgent

DISCLAIMER:

This test was developed and its performance characteristics determined by Fulgent Therapeutics LLC CAP #8042697 CLIA #05D2043189; 4399 Santa Anita Ave., El Monte, CA, 91731. It has not been cleared or approved by the FDA. The laboratory is regulated under CLIA as qualified to perform high-complexity testing. This test is used for clinical purposes. It should not be regarded as investigational or for research. Since genetic variation, as well as systematic and technical factors, can affect the accuracy of testing, the results of testing should always be interpreted in the context of clinical and familial data. For assistance with interpretation of these results, healthcare professionals may contact us directly at **626-350-0537** or by email at **info@fulgentgenetics.com**. It is recommended that patients receive appropriate genetic counseling to explain the implications of the test result, including its residual risks, uncertainties and reproductive or medical options.

To view the supplemental table describing the carrier frequencies, detection rates, and residual risks associated with the genes on this test please visit the following link:
Beacon Expanded Carrier Screening Supplemental Table

LabCorp
Laboratory Corporation of America

Client/Sending Facility:
Bellevue Sperm Bank

1220 116th Ave NE Ste 201
Bellevue, WA 98004
Ph: (206)588-1484
WAB-55

LCLS Specimen Number:	**314-129-2611-0**	Account Number:	46071420
Patient Name:	**16010, DONOR**	Ordering Physician:	**J OLLIFFE**
Date of Birth:	[redacted]	Specimen Type:	**BLOOD**
Gender:	M	Client Reference:	
Patient ID:		Date Collected:	11/10/2021
Lab Number:	DCGWA21-4489 L	Date Received:	11/11/2021
Indications:	DONOR	Date Reported:	**12/15/2021**
Test:	**Chromosome, Blood, Routine**		
Cells Counted:	20	Cells Karyotyped:	4
Cells Analyzed:	5	Band Resolution:	550

CYTOGENETIC RESULT: 46,XY

INTERPRETATION: NORMAL MALE KARYOTYPE

Cytogenetic analysis of PHA stimulated cultures has revealed a MALE karyotype with an apparently normal GTW banding pattern in all cells observed.


This result does not exclude the possibility of subtle rearrangements below the resolution of cytogenetics or congenital anomalies due to other etiologies.

LabCorp
Laboratory Corporation of America

Client/Sending Facility:
Bellevue Sperm Bank

1220 116th Ave NE Ste 201
Bellevue, WA 98004
Ph: (206)588-1484
WAB-55

LCLS Specimen Number:	**314-129-2611-0**	Account Number:	46071420
Patient Name:	**16010, DONOR**	Ordering Physician:	**J OLLIFFE**
Date of Birth:	[redacted]	Specimen Type:	**BLOOD**
Gender:	M	Client Reference:	
Patient ID:		Date Collected:	11/10/2021
Lab Number:	DCGWA21-4489 L	Date Received:	11/11/2021



LabCorp
Laboratory Corporation of America

Client/Sending Facility:
Bellevue Sperm Bank

1220 116th Ave NE Ste 201
Bellevue, WA 98004
Ph: (206)588-1484
WAB-55

LCLS Specimen Number:	**314-129-2611-0**	Account Number:	46071420
Patient Name:	**16010, DONOR**	Ordering Physician:	**J OLLIFFE**
Date of Birth:	[redacted]	Specimen Type:	**BLOOD**
Gender:	M	Client Reference:	
Patient ID:		Date Collected:	11/10/2021
Lab Number:	DCGWA21-4489 L	Date Received:	11/11/2021

Hiba Risheg, PhD., FACMG

Daniel Toweill, MD
Medical Director
Stuart Schwartz, PhD
National Director of Cytogenetics

Technical component performed by Laboratory Corporation of America Holdings, 550 17th Ave Ste 300 , Seattle , WA , 98122-5789 (206) 861-7000
Professional Component performed by LabCorp CLIA 50D0630157, 550 17th Ave. Suite 200, Seattle WA 98122-5789. Medical Director, Daniel Toweill, MD
This document contains private and confidential health information **protected by state and federal law.**

16010, Donor
Patient ID:
Specimen ID: **314 129 2613 0**

DOB [redacted]
Age: [redacted]
Sex: **Male**

Patient Report
Account Number: **46071420**
Ordering Physician: **J OLLIFFE**

labcorp

Ordered Items: **LP+12AC+CBC/D/Plt+UA+Rh+ABO...**

Date Collected: **11/10/2021** Date Received: **11/11/2021** Date Reported: **11/15/2021** Fasting: **No**

LP+12AC+CBC/D/Plt+UA+Rh+ABO...

	Test	Current Result and Flag		Previous Result and Date	Units	Reference Interval
	Glucose [01]	95			mg/dL	65-99
	Uric Acid [01]	7.0			mg/dL	3.8-8.4
				Therapeutic target for gout patients: <6.0		
	BUN [01]	14			mg/dL	6-20
	Creatinine [01]	0.85			mg/dL	0.76-1.27
	eGFR If NonAfricn Am	123			mL/min/1.73	>59
	eGFR If Africn Am	142			mL/min/1.73	>59
		In accordance with recommendations from the NKF-ASN Task force, Labcorp is in the process of updating its eGFR calculation to the 2021 CKD-EPI creatinine equation that estimates kidney function without a race variable.				
	Calcium [01]	9.9			mg/dL	8.7-10.2
	Protein, Total [01]	7.5			g/dL	6.0-8.5
	Albumin [01]	4.8			g/dL	4.1-5.2
	Bilirubin, Total [01]	0.2			mg/dL	0.0-1.2
	Alkaline Phosphatase [01]	78			IU/L	44-121
				Please note reference interval change		
	LDH [01]	147			IU/L	121-224
	AST (SGOT) [01]	21			IU/L	0-40
	ALT (SGPT) [01]	24			IU/L	0-44
	Cholesterol, Total [01]	190			mg/dL	100-199
▲	**Triglycerides** [01]	**305**	**High**		mg/dL	0-149
▼	**HDL Cholesterol** [01]	**36**	**Low**		mg/dL	>39
▲	**LDL Chol Calc (NIH)**	**102**	**High**		mg/dL	0-99
	LDL/HDL Ratio	2.8			ratio	0.0-3.6
	Please Note: [01]					

LDL/HDL Ratio

	Men	Women
1/2 Avg.Risk	1.0	1.5
Avg.Risk	3.6	3.2
2X Avg.Risk	6.2	5.0
3X Avg.Risk	8.0	6.1

	Hgb Fractionation by CE: [02]					
	Hgb F [02]	0.0			%	0.0-2.0
	Hgb A [02]	97.5			%	96.4-98.8
	Hgb A2 [02]	2.5			%	1.8-3.2
	Hgb S [02]	0.0			%	0.0
	Interpretation: [02]					
		Normal hemoglobin present; no hemoglobin variant or thalassemia observed.				
	ABO Grouping [01]	O				

©2021 Laboratory Corporation of America® Holdings
All Rights Reserved Enterprise Report Version 2.00

This document contains private and confidential health information protected by state and federal law.
If you have received this document in error please call 206 861 7000

LP+12AC+CBC/D/Plt+UA+Rh+ABO... (Cont.)

Rh Factor[01]	Positive			
	Please note: Prior records for this patient's ABO / Rh type are not available for additional verification.			
.[01]				
CBC, Platelet Ct, and Diff[01]				
WBC[01]	8.1		x10E3/uL	3.4-10.8
RBC[01]	5.05		x10E6/uL	4.14-5.80
Hemoglobin[01]	15.0		g/dL	13.0-17.7
Hematocrit[01]	43.2		%	37.5-51.0
MCV[01]	86		fL	79-97
MCH[01]	29.7		pg	26.6-33.0
MCHC[01]	34.7		g/dL	31.5-35.7
RDW[01]	12.5		%	11.6-15.4
Platelets[01]	378		x10E3/uL	150-450
Neutrophils[01]	52		%	Not Estab.
Lymphs[01]	33		%	Not Estab.
Monocytes[01]	13		%	Not Estab.
Eos[01]	1		%	Not Estab.
Basos[01]	0		%	Not Estab.
Neutrophils (Absolute)[01]	4.2		x10E3/uL	1.4-7.0
Lymphs (Absolute)[01]	2.7		x10E3/uL	0.7-3.1
▲ **Monocytes(Absolute)**[01]	**1.0**	**High**	x10E3/uL	0.1-0.9
Eos (Absolute)[01]	0.1		x10E3/uL	0.0-0.4
Baso (Absolute)[01]	0.0		x10E3/uL	0.0-0.2
Immature Granulocytes[01]	1		%	Not Estab.
Immature Grans (Abs)[01]	0.1		x10E3/uL	0.0-0.1
.[01]				
Urinalysis Gross Exam[01]				
Specific Gravity[01]	1.014			1.005-1.030
pH[01]	6.5			5.0-7.5
Urine-Color[01]	Yellow			Yellow
Appearance[01]	Clear			Clear
WBC Esterase[01]	Negative			Negative
Protein[01]	Negative			Negative/Trace
Glucose[01]	Negative			Negative
Ketones[01]	Negative			Negative
Occult Blood[01]	Negative			Negative
Bilirubin[01]	Negative			Negative
Urobilinogen,Semi-Qn[01]	0.2		mg/dL	0.2-1.0
Nitrite, Urine[01]	Negative			Negative
Microscopic Examination[01]				
	Microscopic not indicated and not performed.			

©2021 Laboratory Corporation of America® Holdings
All Rights Reserved Enterprise Report Version 2.00

This document contains private and confidential health information protected by state and federal law.
If you have received this document in error please call 206 861 7000