

Patient name:	DONOR 13057	Sample type:	Blood	Report date:	02-JUL-2023
DOB:	██████████	Sample collection date:	07-JUN-2023	Invitae #:	RQ5180029
Sex assigned at birth:	Male	Sample accession date:	08-JUN-2023	Clinical team:	Sarah O'Brien Jeffrey Olliffe
Gender:					
Patient ID (MRN):					

Reason for testing

Gamete donor

Test performed

Invitae Carrier Screen


RESULT: POSITIVE

This carrier test evaluated 514 gene(s) for genetic changes (variants) that are associated with an increased risk of having a child with a genetic condition. Knowledge of carrier status for one of these conditions may provide information that can be used to assist with family planning and/or preparation. Carrier screening is not intended for diagnostic purposes. To identify a potential genetic basis for a condition in the individual being tested, diagnostic testing for the gene(s) of interest is recommended.

This test shows the presence of clinically significant genetic change(s) in this individual in the gene(s) indicated below. No other clinically significant changes were identified in the remaining genes evaluated with this test.

RESULTS	GENE	VARIANT(S)	INHERITANCE	PARTNER TESTING RECOMMENDED
Carrier: Congenital adrenal hyperplasia due to 21-hydroxylase deficiency	CYP21A2	c.844G>T (p.Val282Leu)	Autosomal recessive	Yes
Carrier: Congenital disorder of glycosylation type Ik	ALG1	c.773C>T (p.Ser258Leu)	Autosomal recessive	Yes
Carrier: Gitelman syndrome	SLC12A3	c.2186G>T (p.Gly729Val)	Autosomal recessive	Yes



Next steps

- See the table above for recommendations regarding testing of this individual's reproductive partner.
- Even for genes that have a negative test result, there is always a small risk that an individual could still be a carrier. This is called “residual risk.” See the Carrier detection rates and residual risks document.
- Discussion with a physician and/or genetic counselor is recommended to further review the implications of this test result and to understand these results in the context of any family history of a genetic condition.
- All patients, regardless of result, may wish to consider additional screening for hemoglobinopathies by complete blood count (CBC) and hemoglobin electrophoresis, if this has not already been completed.
- Individuals can register their tests at <https://www.invitae.com/patients/> to access online results, educational resources, and next steps.

Clinical summary

RESULT: CARRIER

Congenital adrenal hyperplasia due to 21-hydroxylase deficiency

A single Pathogenic variant, c.844G>T (p.Val282Leu), was identified in CYP21A2. This variant is primarily associated with non-classic congenital adrenal hyperplasia due to 21-hydroxylase deficiency. See "What is congenital adrenal hyperplasia due to 21-hydroxylase deficiency?" and Variant details for additional information.

What is congenital adrenal hyperplasia due to 21-hydroxylase deficiency?

21-hydroxylase deficiency (21-OHD) is one of a group of conditions called congenital adrenal hyperplasia (CAH), which impair hormone production by the adrenal glands. The adrenal glands produce hormones that regulate many essential functions in the body, including sexual development and maturation. There are several types of CAH, which are caused by changes in different genes.


Symptoms of 21-OHD CAH range in severity, and are caused by the adrenal glands producing excess male sex hormones (androgens). There are three types of 21-OHD which include two classic forms, known as the salt-wasting and simple virilizing types, and the third is called the non-classic type. The salt-wasting type is the most severe, the simple virilizing type is less severe, and the non-classic type is the mildest form. Individuals with the salt-wasting type of 21-OHD lose large amounts of sodium in the urine, which can be life-threatening in early infancy. Infants with the simple virilizing type of 21-OHD do not experience salt-wasting. Female infants with classic 21-OHD usually have external genitalia that do not look clearly male or female (ambiguous genitalia). Male infants with classic 21-OHD usually have normal genitalia, although the testes may be smaller than typical. Individuals with a classic form of 21-OHD may have decreased fertility. Females with non-classic 21-OHD are born with typical external genitalia. They may experience irregular menstruation, decreased fertility, excess hair growth on the face and body (hirsutism), and male-pattern baldness. Males with non-classic 21-OHD may experience early beard growth and have small testes. Some individuals with non-classic 21-OHD may not have signs or symptoms of the condition (asymptomatic). The form(s) of 21-OHD CAH for which an individual would be at risk depends on the specific CYP21A2 variants inherited from the reproductive parents. Follow-up depends on each affected individual's specific situation, and discussion with a healthcare provider should be considered.

Next steps

Carrier testing for the reproductive partner is recommended.

If your partner tests positive:

The various forms of congenital adrenal hyperplasia due to 21-hydroxylase deficiency are inherited in an autosomal recessive fashion. In autosomal recessive inheritance, an individual must have disease-causing genetic changes in each copy of the CYP21A2 gene to be affected. Carriers, who have a disease-causing genetic change in only one copy of the gene, typically do not have symptoms. When both reproductive partners are carriers of an autosomal recessive condition, there is a 25% chance for each child to have the condition. The form(s) of 21-OHD CAH for which an individual's offspring would be at risk depends on the specific CYP21A2 variants inherited from the reproductive parents. When an individual has a CYP21A2 variant on each chromosome (in trans), and at least one of the variants is most commonly associated with the non-classic form of the condition, then the individual is most likely to be at risk to have non-classic 21-OHD.



If your partner tests negative:

A negative carrier test result reduces, but does not eliminate, the chance that a person may be a carrier. The risk that a person could still be a carrier, even after a negative test result, is called a residual risk. See the table below for your partner's hypothetical residual risk after testing negative for congenital adrenal hyperplasia due to 21-hydroxylase deficiency. These values are provided only as a guide, are based on the


Patient name: DONOR 13057 **DOB:** [REDACTED]

Invitae #: RQ5180029

detection rate for the condition as tested at Invitae, and assume a negative family history, the absence of symptoms, and vary based on the ethnic background of an individual. For genes associated with both dominant and recessive inheritance, the numbers provided apply to the recessive condition(s) associated with the gene.

DISORDER (INHERITANCE)	GENE	ETHNICITY	CARRIER FREQUENCY BEFORE SCREENING	CARRIER RESIDUAL RISK AFTER NEGATIVE RESULT
Congenital adrenal hyperplasia due to 21-hydroxylase deficiency (AR) NM_000500.7	CYP21A2 *	Pan-ethnic	1 in 61	1 in 751


RESULT: CARRIER

Congenital disorder of glycosylation type 1k

A single Pathogenic variant, c.773C>T (p.Ser258Leu), was identified in ALG1.

What is congenital disorder of glycosylation type 1k?

Congenital disorders of glycosylation (CDGs) are a group of conditions in which individuals have difficulty adding or removing a sugar group to make glycoproteins. This process, called “glycosylation,” is a necessary step to modify proteins for their intended purpose. CDGs affect several body systems. Symptoms of CDG type 1k (CDG-1k) typically present during the first year of life and commonly include a small head (microcephaly), abnormal fat distribution, eye problems such as eyes that do not look in the same direction (strabismus) and involuntary eye movements (nystagmus), problems with blood clotting such as excessive bleeding (hemorrhage) or a tendency to form abnormal clots in blood vessels (thrombosis), difficulty coordinating movements (ataxia), seizures, developmental delay, and intellectual disability. A few patients have been observed with a milder clinical course, but affected individuals often do not survive past the neonatal period. Follow-up depends on each affected individual’s specific situation, and discussion with a healthcare provider should be considered.

Next steps

Carrier testing for the reproductive partner is recommended.




If your partner tests positive:

In autosomal recessive inheritance, an individual must have disease-causing genetic changes in each copy of the ALG1 gene to be affected. Carriers, who have a disease-causing genetic change in only one copy of the gene, typically do not have symptoms. When both reproductive partners are carriers of an autosomal recessive condition, there is a 25% chance for each child to have the condition.



If your partner tests negative:

A negative carrier test result reduces, but does not eliminate, the chance that a person may be a carrier. The risk that a person could still be a carrier, even after a negative test result, is called a residual risk. See the table below for your partner’s hypothetical residual risk after testing negative for congenital disorder of glycosylation type 1k. These values are provided only as a guide, are based on the detection rate for the condition as tested at Invitae, and assume a negative family history, the absence of symptoms, and vary based on the ethnic background of an individual. For genes associated with both dominant and recessive inheritance, the numbers provided apply to the recessive condition(s) associated with the gene.



DISORDER (INHERITANCE)	GENE	ETHNICITY	CARRIER FREQUENCY BEFORE SCREENING	CARRIER RESIDUAL RISK AFTER NEGATIVE RESULT
Congenital disorder of glycosylation type 1k (AR) NM_019109.4	ALG1	Pan-ethnic	≤1 in 500	Reduced


RESULT: CARRIER

Gitelman syndrome

A single Pathogenic variant, c.2186G>T (p.Gly729Val), was identified in SLC12A3.

What is Gitelman syndrome?

Gitelman syndrome is a condition that affects kidney function. Affected individuals have difficulty reabsorbing salt from urine back into the blood, causing low levels of potassium (hypokalemia), magnesium (hypomagnesemia), and calcium (hypocalciuria). Symptoms usually appear in late childhood or adolescence and may include intermittent muscle spasms (tetany), muscle weakness or cramps, joint pain caused by calcium crystals forming in the cartilage (chondrocalcinosis), fatigue, excessive thirst (polydipsia), excessive urination (polyuria), tingling or prickly sensation in the skin caused by pressure on, or damage to, peripheral nerves (paresthesia) and a sensation of rapid heart beats (palpitations). Less commonly, affected individuals have short stature, seizures, abnormally high or low thyroid hormone levels (hyperthyroidism or hypothyroidism, respectively), or an abnormal heart rhythm (arrhythmia). Gitelman syndrome is a highly variable condition; symptoms can vary widely, even among affected members of the same family, and in some cases there may only be mild symptoms or no overt symptoms of the condition (asymptomatic). With treatment and dietary management, affected individuals have a normal life expectancy.

Next steps

Carrier testing for the reproductive partner is recommended.




If your partner tests positive:

In autosomal recessive inheritance, an individual must have disease-causing genetic changes in each copy of the SLC12A3 gene to be affected. Carriers, who have a disease-causing genetic change in only one copy of the gene, typically do not have symptoms. When both reproductive partners are carriers of an autosomal recessive condition, there is a 25% chance for each child to have the condition.



If your partner tests negative:

A negative carrier test result reduces, but does not eliminate, the chance that a person may be a carrier. The risk that a person could still be a carrier, even after a negative test result, is called a residual risk. See the table below for your partner's hypothetical residual risk after testing negative for Gitelman syndrome. These values are provided only as a guide, are based on the detection rate for the condition as tested at Invitae, and assume a negative family history, the absence of symptoms, and vary based on the ethnic background of an individual. For genes associated with both dominant and recessive inheritance, the numbers provided apply to the recessive condition(s) associated with the gene.



DISORDER (INHERITANCE)	GENE	ETHNICITY	CARRIER FREQUENCY BEFORE SCREENING	CARRIER RESIDUAL RISK AFTER NEGATIVE RESULT
Gitelman syndrome (AR) NM_000339.2	SLC12A3	Pan-ethnic	1 in 100	1 in 9900

Results to note

SMN1

- Negative result. SMN1: 2 copies; c.*3+80T>G not detected.

Pseudodeficiency allele(s)

- Benign change, c.1685T>C (p.Ile562Thr), known to be a pseudodeficiency allele, identified in the GALC gene. Pseudodeficiency alleles are not known to be associated with disease, including Krabbe disease.
- The presence of a pseudodeficiency allele does not impact this individual's risk to be a carrier. Individuals with pseudodeficiency alleles may exhibit false positive results on related biochemical tests, including newborn screening. However, pseudodeficiency alleles are not known to cause disease, even when there are two copies of the variant (homozygous) or when in combination with another disease-causing variant (compound heterozygous). Carrier testing for the reproductive partner is not indicated based on this result.

Variant details

ALG1, Exon 7, c.773C>T (p.Ser258Leu), heterozygous, PATHOGENIC

- This sequence change replaces serine, which is neutral and polar, with leucine, which is neutral and non-polar, at codon 258 of the ALG1 protein (p.Ser258Leu).
- This variant is present in population databases (rs28939378, gnomAD 0.05%).
- This missense change has been observed in individuals with congenital disorder of glycosylation (PMID: 14709599, 14973778, 14973782, 20679665, 22966035, 26931382, 27172925, 27325525, 28554332).
- ClinVar contains an entry for this variant (Variation ID: 4724).
- Advanced modeling of protein sequence and biophysical properties (such as structural, functional, and spatial information, amino acid conservation, physicochemical variation, residue mobility, and thermodynamic stability) performed at Invitae indicates that this missense variant is not expected to disrupt ALG1 protein function.
- Experimental studies have shown that this missense change affects ALG1 function (PMID: 14973778, 22966035, 26931382).
- For these reasons, this variant has been classified as Pathogenic.

CYP21A2, Exon 7, c.844G>T (p.Val282Leu), heterozygous, PATHOGENIC

- This sequence change replaces valine, which is neutral and non-polar, with leucine, which is neutral and non-polar, at codon 282 of the CYP21A2 protein (p.Val282Leu).
- The frequency data for this variant in the population databases (gnomAD) is considered unreliable due to the presence of homologous sequence, such as pseudogenes or paralogs, in the genome.
- This missense change has been observed in individual(s) with primarily non-classic, and less frequently, classic salt-wasting or simple virilizing, congenital adrenal hyperplasia due to 21-hydroxylase deficiency (PMID: 1644925, 23359698, 24953648, 26804566, 31344365). In at least one individual the data is consistent with being in trans (on the opposite chromosome) from a pathogenic variant. It has also been observed to segregate with disease in related individuals.
- This variant is also known as V281L.
- ClinVar contains an entry for this variant (Variation ID: 12151).
- Advanced modeling of protein sequence and biophysical properties (such as structural, functional, and spatial information, amino acid conservation, physicochemical variation, residue mobility, and thermodynamic stability) performed at Invitae indicates that this missense variant is not expected to disrupt CYP21A2 protein function.
- Experimental studies have shown that this missense change affects CYP21A2 function (PMID: 1864962, 2249999, 31344365).
- For these reasons, this variant has been classified as Pathogenic.



SLC12A3, Exon 18, c.2186G>T (p.Gly729Val), heterozygous, PATHOGENIC

- This sequence change replaces glycine, which is neutral and non-polar, with valine, which is neutral and non-polar, at codon 729 of the SLC12A3 protein (p.Gly729Val).
- This variant is present in population databases (rs373901523, gnomAD 0.02%).
- This missense change has been observed in individuals with Gitelman syndrome (PMID: 11168953, 17654016, 31672324).
- ClinVar contains an entry for this variant (Variation ID: 994770).
- Advanced modeling of protein sequence and biophysical properties (such as structural, functional, and spatial information, amino acid conservation, physicochemical variation, residue mobility, and thermodynamic stability) performed at Invitae indicates that this missense variant is expected to disrupt SLC12A3 protein function.
- This variant disrupts the p.Gly729 amino acid residue in SLC12A3. Other variant(s) that disrupt this residue have been observed in individuals with SLC12A3-related conditions (PMID: 21415153, 31672324), which suggests that this may be a clinically significant amino acid residue.
- For these reasons, this variant has been classified as Pathogenic.

Residual risk

No carrier test can detect 100% of carriers. There still remains a small risk of being a carrier after a negative test (residual risk). Residual risk values assume a negative family history and are inferred from published carrier frequencies and estimated detection rates based on testing technologies used at Invitae. You can view Invitae's complete Carrier detection rates and residual risks document (containing all carrier genes) online at <https://www.invitae.com/carrier-residual-risks/>. Additionally, the order-specific information for this report is available to download in the portal (under this order's documents) or can be requested by contacting Invitae Client Services. The complete Carrier detection rates and residual risks document will not be applicable for any genes with specimen-specific limitations in sequencing and/or deletion/duplication coverage. Please see the final bullet point in the Limitations section of this report to view if this specimen had any gene-specific coverage gaps.

Genes analyzed

This table represents a complete list of genes analyzed for this individual, including the relevant gene transcript(s). If more than one transcript is listed for a single gene, variants were reported using the first transcript listed unless otherwise indicated in the report. An asterisk (*) indicates that this gene has a limitation. Please see the Limitations section for details. Results are negative, unless otherwise indicated in the report.

GENE	TRANSCRIPT	GENE	TRANSCRIPT	GENE	TRANSCRIPT
AAAS	NM_015665.5	AP1S1	NM_001283.3	CBS	NM_000071.2
ABCA12	NM_173076.2	AQP2	NM_000486.5	CC2D1A	NM_017721.5
ABCA3	NM_001089.2	ARG1	NM_000045.3	CC2D2A	NM_001080522.2
ABCA4	NM_000350.2	ARL6	NM_177976.2	CCDC103	NM_213607.2
ABCB11	NM_003742.2	ARSA	NM_000487.5	CCDC39	NM_181426.1
ABCB4	NM_000443.3	ARSB	NM_000046.3	CCDC88C	NM_001080414.3
ABCC2*	NM_000392.4	ASL	NM_000048.3	CD3D	NM_000732.4
ABCC8	NM_000352.4	ASNS	NM_133436.3	CD3E	NM_000733.3
ACAD9	NM_014049.4	ASPA	NM_000049.2	CD40	NM_001250.5
ACADM	NM_000016.5	ASS1	NM_000050.4	CD59	NM_203330.2
ACADVL	NM_000018.3	ATM*	NM_000051.3	CDH23	NM_022124.5
ACAT1	NM_000019.3	ATP6V1B1	NM_001692.3	CEP152	NM_014985.3
ACOX1	NM_004035.6	ATP7B	NM_000053.3	CEP290	NM_025114.3
ACSF3	NM_174917.4	ATP8B1*	NM_005603.4	CERKL	NM_001030311.2
ADA	NM_000022.2	BBS1	NM_024649.4	CFTR*	NM_000492.3
ADAMTS2	NM_014244.4	BBS10	NM_024685.3	CHAT	NM_020549.4
ADAMTS4	NM_019032.5	BBS12	NM_152618.2	CHRNE	NM_000080.3
ADGRG1	NM_005682.6	BBS2	NM_031885.3	CHRNA	NM_005199.4
ADGRV1	NM_032119.3	BBS4	NM_033028.4	CIITA	NM_000246.3
AGA	NM_000027.3	BBS5	NM_152384.2	CLCN1	NM_000083.2
AGL	NM_000642.2	BBS7	NM_176824.2	CLN3	NM_001042432.1
AGPS	NM_003659.3	BBS9*	NM_198428.2	CLN5	NM_006493.2
AGXT	NM_000030.2	BCKDHA	NM_000709.3	CLN6	NM_017882.2
AHI1	NM_017651.4	BCKDHB	NM_183050.2	CLN8	NM_018941.3
AIPL1*	NM_014336.4	BCL1L	NM_004328.4	CLRN1	NM_174878.2
AIRE	NM_000383.3	BLM	NM_000057.3	CNGB3	NM_019098.4
ALDH3A2	NM_000382.2	BLOC1S3	NM_212550.4	COL11A2*	NM_080680.2
ALDH7A1	NM_001182.4	BLOC1S6	NM_012388.3	COL17A1	NM_000494.3
ALDOB	NM_000035.3	BMP1	NM_006129.4;NM_001199.3	COL27A1	NM_032888.3
ALG1	NM_019109.4	BRIP1	NM_032043.2	COL4A3	NM_000091.4
ALG6	NM_013339.3	BSND	NM_057176.2	COL4A4	NM_000092.4
ALMS1	NM_015120.4	BTBD	NM_000060.3	COL7A1	NM_000094.3
ALPL	NM_000478.5	CAD	NM_004341.4	COX15	NM_004376.6
AMN*	NM_030943.3	CANT1	NM_138793.3	CPS1	NM_001875.4
AMT	NM_000481.3	CAPN3	NM_000070.2	CPT1A	NM_001876.3
ANO10*	NM_018075.3	CASQ2	NM_001232.3	CPT2	NM_000098.2

GENE	TRANSCRIPT
CRB1	NM_201253.2
CRTAP	NM_006371.4
CTNS	NM_004937.2
CTSA	NM_000308.3
CTSC	NM_001814.5
CTSD	NM_001909.4
CTSK	NM_000396.3
CYBA	NM_000101.3
CYP11A1	NM_000781.2
CYP11B1	NM_000497.3
CYP11B2	NM_000498.3
CYP17A1	NM_000102.3
CYP19A1	NM_031226.2
CYP1B1	NM_000104.3
CYP21A2*	NM_000500.7
CYP27A1	NM_000784.3
CYP27B1	NM_000785.3
CYP7B1	NM_004820.3
DBT	NM_001918.3
DCAF17	NM_025000.3
DCLRE1C	NM_001033855.2
DDX11*	NM_030653.3
DFNB59	NM_001042702.3
DGAT1	NM_012079.5
DGUOK	NM_080916.2
DHCR7	NM_001360.2
DHDDS	NM_024887.3
DLD	NM_000108.4
DLL3	NM_016941.3
DNAH11	NM_001277115.1
DNAH5	NM_001369.2
DNAI1	NM_012144.3
DNAI2	NM_023036.4
DNMT3B	NM_006892.3
DOK7	NM_173660.4
DUOX2*	NM_014080.4
DYNC2H1	NM_001080463.1
DYSF	NM_003494.3
EIF2AK3	NM_004836.6

GENE	TRANSCRIPT
EIF2B1	NM_001414.3
EIF2B2	NM_014239.3
EIF2B3	NM_020365.4
EIF2B4	NM_015636.3
EIF2B5	NM_003907.2
ELP1	NM_003640.3
EPG5	NM_020964.2
ERCC2	NM_000400.3
ERCC6	NM_000124.3
ERCC8	NM_000082.3
ESCO2	NM_001017420.2
ETFA	NM_000126.3
ETFB	NM_001985.2
ETFDH	NM_004453.3
ETHE1	NM_014297.3
EVC	NM_153717.2
EVC2	NM_147127.4
EXOSC3	NM_016042.3
EYS*	NM_001142800.1
FAH*	NM_000137.2
FAM161A	NM_001201543.1
FANCA	NM_000135.2
FANCC	NM_000136.2
FANCD2*	NM_033084.3
FANCE	NM_021922.2
FANCG	NM_004629.1
FANCI	NM_001113378.1
FANCL*	NM_018062.3
FBP1	NM_000507.3
FBXO7	NM_012179.3
FH*	NM_000143.3
FKBP10	NM_021939.3
FKRP	NM_024301.4
FKTN	NM_001079802.1
FMO3	NM_006894.6
FOXN1	NM_003593.2
FOXRED1	NM_017547.3
FRAS1	NM_025074.6
FREM2	NM_207361.5

GENE	TRANSCRIPT
FUCA1	NM_000147.4
G6PC	NM_000151.3
G6PC3	NM_138387.3
GAA	NM_000152.3
GALC*	NM_000153.3
GALE*	NM_000403.3
GALK1	NM_000154.1
GALNS	NM_000512.4
GALNT3	NM_004482.3
GALT	NM_000155.3
GAMT	NM_000156.5
GATM	NM_001482.2
GBA*	NM_001005741.2
GBE1	NM_000158.3
GCDH	NM_000159.3
GCH1	NM_000161.2
GDF5	NM_000557.4
GFM1	NM_024996.5
GHR*	NM_000163.4
GJB2	NM_004004.5
GLB1	NM_000404.2
GLDC	NM_000170.2
GLE1	NM_001003722.1
GNE*	NM_001128227.2
GNPAT	NM_014236.3
GNPTAB	NM_024312.4
GNPTG	NM_032520.4
GNS	NM_002076.3
GORAB	NM_152281.2
GRHPR	NM_012203.1
GRIP1	NM_021150.3
GSS	NM_000178.2
GUCY2D	NM_000180.3
GUSB	NM_000181.3
HADH	NM_005327.4
HADHA	NM_000182.4
HADHB	NM_000183.2
HAMP	NM_021175.2
HAX1	NM_006118.3



Patient name: DONOR 13057 DOB: [REDACTED]

Invitae #: RQ5180029

GENE	TRANSCRIPT
HBA1*	NM_000558.4
HBA2	NM_000517.4
HBB	NM_000518.4
HEXA	NM_000520.4
HEXB	NM_000521.3
HGSNAT	NM_152419.2
HJV	NM_213653.3
HLCS	NM_000411.6
HMGCL	NM_000191.2
HMOX1	NM_002133.2
HOGA1	NM_138413.3
HPD	NM_002150.2
HPS1	NM_000195.4
HPS3	NM_032383.4
HPS4	NM_022081.5
HPS5	NM_181507.1
HPS6	NM_024747.5
HSD17B3	NM_000197.1
HSD17B4	NM_000414.3
HSD3B2	NM_000198.3
HYAL1	NM_153281.1
HYLS1	NM_145014.2
IDUA	NM_000203.4
IGHMBP2	NM_002180.2
IKBB	NM_001556.2
IL7R	NM_002185.3
INVS	NM_014425.3
ITGA6	NM_000210.3
ITGB3	NM_000212.2
ITGB4	NM_001005731.2
IVD	NM_002225.3
JAK3	NM_000215.3
KCNJ1	NM_000220.4
KCNJ11	NM_000525.3
LAMA2	NM_000426.3
LAMA3	NM_000227.4
LAMB3	NM_000228.2
LAMC2	NM_005562.2
LARGE1	NM_004737.4

GENE	TRANSCRIPT
LCA5	NM_181714.3
LDLR	NM_000527.4
LDLRAP1	NM_015627.2
LHX3	NM_014564.4
LIFR*	NM_002310.5
LIG4	NM_002312.3
LIPA	NM_000235.3
LMBRD1	NM_018368.3
LOXHD1	NM_144612.6
LPL	NM_000237.2
LRAT	NM_004744.4
LRP2	NM_004525.2
LRPPRC	NM_133259.3
LYST	NM_000081.3
MAK	NM_001242957.2
MAN2B1	NM_000528.3
MANBA	NM_005908.3
MCEE	NM_032601.3
MCOLN1	NM_020533.2
MCPH1	NM_024596.4
MECR	NM_016011.3
MED17	NM_004268.4
MESP2	NM_001039958.1
MFSD8	NM_152778.2
MKKS	NM_018848.3
MKS1	NM_017777.3
MLC1*	NM_015166.3
MLYCD	NM_012213.2
MMAA	NM_172250.2
MMAB	NM_052845.3
MMACHC	NM_015506.2
MMADHC	NM_015702.2
MOCS1	NM_001358530.2
MOCS2A	NM_176806.3
MOCS2B	NM_004531.4
MPI	NM_002435.2
MPL	NM_005373.2
MPV17	NM_002437.4
MRE11	NM_005591.3

GENE	TRANSCRIPT
MTHFR*	NM_005957.4
MTR	NM_000254.2
MTRR	NM_002454.2
MTTP	NM_000253.3
MUSK	NM_005592.3
MUT	NM_000255.3
MVK	NM_000431.3
MYO15A	NM_016239.3
MYO7A	NM_000260.3
NAGA	NM_000262.2
NAGLU	NM_000263.3
NAGS	NM_153006.2
NBN	NM_002485.4
NCF2	NM_000433.3
NDRG1	NM_006096.3
NDUFAF2	NM_174889.4
NDUFAF5	NM_024120.4
NDUFS4	NM_002495.3
NDUFS6	NM_004553.4
NDUFS7	NM_024407.4
NDUFV1	NM_007103.3
NEB*	NM_001271208.1
NEU1	NM_000434.3
NGLY1	NM_018297.3
NPC1	NM_000271.4
NPC2	NM_006432.3
NPHP1	NM_000272.3
NPHS1	NM_004646.3
NPHS2	NM_014625.3
NR2E3	NM_014249.3
NSMCE3	NM_138704.3
NTRK1	NM_001012331.1
OAT*	NM_000274.3
OCA2	NM_000275.2
OPA3	NM_025136.3
OSTM1	NM_014028.3
OTOA*	NM_144672.3
OTOF	NM_194248.2;NM_194323.2
P3H1	NM_022356.3


Patient name: DONOR 13057 **DOB:** [REDACTED]

Invitae #: RQ5180029

GENE	TRANSCRIPT
PAH	NM_000277.1
PANK2	NM_153638.2
PC	NM_000920.3
PCBD1	NM_000281.3
PCCA	NM_000282.3
PCCB	NM_000532.4
PCDH15	NM_033056.3
PCNT	NM_006031.5
PDHB	NM_000925.3
PEPD	NM_000285.3
PET100	NM_001171155.1
PEX1*	NM_000466.2
PEX10	NM_153818.1
PEX12	NM_000286.2
PEX13	NM_002618.3
PEX16	NM_004813.2
PEX2	NM_000318.2
PEX26	NM_017929.5
PEX5	NM_001131025.1
PEX6	NM_000287.3
PEX7	NM_000288.3
PFKM	NM_000289.5
PGM3	NM_001199917.1
PHGDH	NM_006623.3
PHKB	NM_000293.2;NM_00103183.5.2
PHKG2	NM_000294.2
PHYH	NM_006214.3
PIGN	NM_176787.4
PKHD1*	NM_138694.3
PLA2G6	NM_003560.2
PLEKHG5	NM_020631.4
PLOD1	NM_000302.3
PMM2	NM_000303.2
PNPO	NM_018129.3
POLG	NM_002693.2
POLH	NM_006502.2
POMGNT1	NM_017739.3
POMT1	NM_007171.3
POMT2	NM_013382.5

GENE	TRANSCRIPT
POR	NM_000941.2
POU1F1	NM_000306.3
PPT1	NM_000310.3
PRCD	NM_001077620.2
PRDM5	NM_018699.3
PRF1	NM_001083116.1
PROP1	NM_006261.4
PSAP	NM_002778.3
PTPRC*	NM_002838.4
PTS	NM_000317.2
PUS1	NM_025215.5
PYGM	NM_005609.3
QDPR	NM_000320.2
RAB23	NM_183227.2
RAG1	NM_000448.2
RAG2	NM_000536.3
RAPSN	NM_005055.4
RARS2	NM_020320.3
RDH12	NM_152443.2
RLBP1	NM_000326.4
RMRP	NR_003051.3
RNASEH2A	NM_006397.2
RNASEH2B	NM_024570.3
RNASEH2C	NM_032193.3
RPE65	NM_000329.2
RPGRIP1L	NM_015272.2
RTEL1	NM_001283009.1
RXYLT1	NM_014254.2
RYR1	NM_000540.2
SACS	NM_014363.5
SAMD9	NM_017654.3
SAMHD1	NM_015474.3
SCO2	NM_005138.2
SEC23B	NM_006363.4
SEPSECS	NM_016955.3
SGCA	NM_000023.2
SGCB	NM_000232.4
SGCD	NM_000337.5
SGCG	NM_000231.2

GENE	TRANSCRIPT
SGSH	NM_000199.3
SKIV2L	NM_006929.4
SLC12A1	NM_000338.2
SLC12A3	NM_000339.2
SLC12A6	NM_133647.1
SLC17A5	NM_012434.4
SLC19A2	NM_006996.2
SLC19A3	NM_025243.3
SLC1A4	NM_003038.4
SLC22A5	NM_003060.3
SLC25A13	NM_014251.2
SLC25A15	NM_014252.3
SLC25A20	NM_000387.5
SLC26A2	NM_000112.3
SLC26A3	NM_000111.2
SLC26A4	NM_000441.1
SLC27A4	NM_005094.3
SLC35A3	NM_012243.2
SLC37A4	NM_001164277.1
SLC38A8	NM_001080442.2
SLC39A4	NM_130849.3
SLC45A2	NM_016180.4
SLC4A11	NM_032034.3
SLC5A5	NM_000453.2
SLC7A7	NM_001126106.2
SMARCA1	NM_014140.3
SMN1*	NM_000344.3
SMPD1	NM_000543.4
SNAP29	NM_004782.3
SPG11	NM_025137.3
SPR	NM_003124.4
SRD5A2	NM_000348.3
ST3GAL5	NM_003896.3
STAR	NM_000349.2
STX11	NM_003764.3
STXBP2	NM_006949.3
SUMF1	NM_182760.3
SUOX	NM_000456.2
SURF1	NM_003172.3


Patient name: DONOR 13057 **DOB:** [REDACTED]

Invitae #: RQ5180029

GENE	TRANSCRIPT
SYNE4	NM_001039876.2
TANGO2	NM_152906.6
TAT	NM_000353.2
TBCD	NM_005993.4
TBCE*	NM_003193.4
TCIRG1	NM_006019.3
TCN2	NM_000355.3
TECPR2	NM_014844.3
TERT	NM_198253.2
TF	NM_001063.3
TFR2	NM_003227.3
TG*	NM_003235.4
TGM1	NM_000359.2
TH	NM_199292.2
TK2	NM_004614.4
TMC1	NM_138691.2
TMEM216	NM_001173990.2
TMEM67	NM_153704.5
TMPRSS3	NM_024022.2
TPO	NM_000547.5
TPP1	NM_000391.3
TREX1	NM_033629.4
TRIM32	NM_012210.3
TRIM37	NM_015294.4
TRMU	NM_018006.4
TSEN54	NM_207346.2
TSFM*	NM_001172696.1
TSHB	NM_000549.4
TSHR	NM_000369.2
TTC37	NM_014639.3
TTPA	NM_000370.3
TULP1	NM_003322.4
TYMP	NM_001953.4
TYR*	NM_000372.4
TYRP1	NM_000550.2
UBR1	NM_174916.2
UNC13D	NM_199242.2
USH1C*	NM_005709.3
USH2A	NM_206933.2

GENE	TRANSCRIPT
VDR	NM_001017535.1
VLDLR	NM_003383.4
VPS11	NM_021729.5
VPS13A*	NM_033305.2
VPS13B	NM_017890.4
VPS45	NM_007259.4
VPS53*	NM_001128159.2
VRK1	NM_003384.2
VSX2	NM_182894.2
WISP3	NM_003880.3
WNT10A	NM_025216.2
WRN*	NM_000553.4
XPA	NM_000380.3
XPC	NM_004628.4
ZBTB24	NM_014797.2
ZFYVE26	NM_015346.3
ZNF469	NM_001127464.2

Methods

- Genomic DNA obtained from the submitted sample is enriched for targeted regions using a hybridization-based protocol, and sequenced using Illumina technology. Unless otherwise indicated, all targeted regions are sequenced with $\geq 50\times$ depth or are supplemented with additional analysis. Reads are aligned to a reference sequence (GRCh37), and sequence changes are identified and interpreted in the context of a single clinically relevant transcript, indicated in the Genes Analyzed table. Enrichment and analysis focus on the coding sequence of the indicated transcripts, 20bp of flanking intronic sequence, and other specific genomic regions demonstrated to be causative of disease at the time of assay design. Promoters, untranslated regions, and other non-coding regions are not otherwise interrogated. Exonic deletions and duplications are called using an in-house algorithm that determines copy number at each target by comparing the read depth for each target in the proband sequence with both mean read-depth and read-depth distribution, obtained from a set of clinical samples. Markers across the X and Y chromosomes are analyzed for quality control purposes and may detect deviations from the expected sex chromosome complement. Such deviations may be included in the report in accordance with internal guidelines. Invitae utilizes a classification methodology to identify next-generation sequencing (NGS)-detected variants that require orthogonal confirmation (Lincoln, et al. J Mol Diagn. 2019 Mar;21(2):318-329). Confirmation of the presence and location of reportable variants is performed as needed based on stringent criteria using one of several validated orthogonal approaches (PubMed ID 30610921). Sequencing is performed by Invitae Corporation (1400 16th Street, San Francisco, CA 94103, #05D2040778). Confirmatory sequencing is performed by Invitae Corporation (1400 16th Street, San Francisco, CA 94103, #05D2040778).

The following additional analyses are performed if relevant to the requisition. For GBA the reference genome has been modified to mask the sites of polymorphic paralog sequence variants (PSVs) in both the gene and pseudogene. For CYP21A2 and GBA, if one or more reportable variants, gene conversion, or fusion event is identified via our NGS pipeline (see Limitations), these variants are confirmed by PacBio sequencing of an amplicon generated by long-range PCR and subsequent short-range PCR. In some cases, it may not be possible to disambiguate between the gene and pseudogene. For GJB2, the reportable range includes large upstream deletions overlapping GJB6. For HBA1/2, the reference genome has been modified to force some sequencing reads derived from HBA1 to align to HBA2, and variant calling algorithms are modified to support an expectation of 4 alleles in these regions. HBA1/2 copy number calling is performed by a custom hypothesis testing algorithm which generates diplotype calls. If sequence data for a sample does not support a unique high confidence match from among hypotheses tested, that sample is flagged for manual review. Copy number variation is only reported for coding sequence of HBA1 and HBA2 and the HS-40 region. This assay does not distinguish among the $\alpha 3.7$ subtypes, and all $\alpha 3.7$ variants are called as HBA1 deletions. This assay may not detect overlapping copy gain and copy loss events when the breakpoints of those events are similar. For FMR1, cytosine-guanine-guanine (CGG) triplet repeats in the 5' untranslated region (5' UTR) of the FMR1 gene are detected by triplet repeat-primed PCR (RP-PCR) with fluorescently labeled primers followed by capillary electrophoresis. Reference ranges: Normal: <45 CGG repeats, intermediate: 45-54 CGG repeats, premutation: 55-200 CGG repeats, full mutation: >200 CGG repeats. For alleles with 55-90 triplet repeats, the region surrounding the FMR1 repeat is amplified by PCR. The PCR amplicons are then processed through PacBio SMRTBell library prep and sequenced using PacBio long read technology. The number of AGG interruptions within the 55-90 triplet repeat is read directly from the resulting DNA sequences.

- This report only includes variants that have a clinically significant association with the conditions tested as of the report date. Variants of uncertain significance, benign variants, and likely benign variants are not included in this report. However, if additional evidence becomes available to indicate that the clinical significance of a variant has changed, Invitae may update this report and provide notification.
- A PMID is a unique identifier referring to a published, scientific paper. Search by PMID at <http://www.ncbi.nlm.nih.gov/pubmed>.
- An rsID is a unique identifier referring to a single genomic position, and is used to associate population frequency information with sequence changes at that position. Reported population frequencies are derived from a number of public sites that aggregate data from large-scale population sequencing projects, including ExAC (<http://exac.broadinstitute.org>), gnomAD (<http://gnomad.broadinstitute.org>), and dbSNP (<http://ncbi.nlm.nih.gov/SNP>).

Disclaimer

DNA studies do not constitute a definitive test for the selected condition(s) in all individuals. It should be realized that there are possible sources of error. Errors can result from trace contamination, rare technical errors, rare genetic variants that interfere with analysis, recent scientific developments, and alternative classification systems. This test should be one of many aspects used by the healthcare provider to help with a diagnosis and treatment plan, but it is not a diagnosis itself. This test was developed and its performance characteristics determined by Invitae. It has not been cleared or approved by

the FDA. The laboratory is regulated under the Clinical Laboratory Improvement Act (CLIA) as qualified to perform high-complexity clinical tests (CLIA ID: 05D2040778). This test is used for clinical purposes. It should not be regarded as investigational or for research.

Limitations

- Based on validation study results, this assay achieves >99% analytical sensitivity and specificity for single nucleotide variants, insertions and deletions <15bp in length, and exon-level deletions and duplications. Invitae's methods also detect insertions and deletions larger than 15bp but smaller than a full exon but sensitivity for these may be marginally reduced. Invitae's deletion/duplication analysis determines copy number at a single exon resolution at virtually all targeted exons. However, in rare situations, single-exon copy number events may not be analyzed due to inherent sequence properties or isolated reduction in data quality. Certain types of variants, such as structural rearrangements (e.g. inversions, gene conversion events, translocations, etc.) or variants embedded in sequence with complex architecture (e.g. short tandem repeats or segmental duplications), may not be detected. Additionally, it may not be possible to fully resolve certain details about variants, such as mosaicism, phasing, or mapping ambiguity. Unless explicitly guaranteed, sequence changes in the promoter, non-coding exons, and other non-coding regions are not covered by this assay. Please consult the test definition on our website for details regarding regions or types of variants that are covered or excluded for this test. This report reflects the analysis of an extracted genomic DNA sample. While this test is intended to reflect the analysis of extracted genomic DNA from a referred patient, in very rare cases the analyzed DNA may not represent that individual's constitutional genome, such as in the case of a circulating hematolymphoid neoplasm, bone marrow transplant, blood transfusion, chimerism, culture artifact or maternal cell contamination.
- GALE: Sequencing analysis for exons 10 includes only cds +/- 5 bp. DDX11: NM_030653.3:c.1763-1G>C variant only. PKHD1: Deletion/duplication analysis is not offered for exon 13. VPS13A: Deletion/duplication analysis is not offered for exons 2-3, 27-28. PTPRC: Sequencing analysis is not offered for exons 3, 15. ABCC2: Deletion/duplication analysis is not offered for exons 24-25. OTOA: Deletion/duplication and sequencing analysis is not offered for exons 20-28. DUOX2: Deletion/duplication and sequencing analysis is not offered for exons 6-7. GNE: Sequencing analysis for exons 8 includes only cds +/- 10 bp. NEB: Deletion/duplication analysis is not offered for exons 82-105. NEB variants in this region with no evidence towards pathogenicity are not included in this report, but are available upon request. TBCE: Sequencing analysis for exons 2 includes only cds +/- 10 bp. BBS9: Deletion/duplication analysis is not offered for exon 4. EYS: Sequencing analysis for exons 30 includes only cds +/- 0 bp. FH: Sequencing analysis for exons 9 includes only cds +/- 10 bp. CFTR: Sequencing analysis for exons 7 includes only cds +/- 10 bp. GHR: Deletion/duplication and sequencing analysis is not offered for exon 3. WRN: Deletion/duplication analysis is not offered for exons 10-11. Sequencing analysis for exons 8, 10-11 includes only cds +/- 10 bp. OAT: Deletion/duplication analysis is not offered for exon 2. ANO10: Sequencing analysis for exons 8 includes only cds +/- 0 bp. ATP8B1: Sequencing analysis for exons 19 includes only cds +/- 10 bp. FANCD2: Deletion/duplication analysis is not offered for exons 14-17, 22 and sequencing analysis is not offered for exons 15-17. Sequencing analysis for exons 6, 14, 18, 20, 23, 25, 34 includes only cds +/- 10 bp. GBA: c.84dupG (p.Leu29Alafs*18), c.115+1G>A (Splice donor), c.222_224delTAC (p.Thr75del), c.475C>T (p.Arg159Trp), c.595_596delCT (p.Leu199Aspfs*62), c.680A>G (p.Asn227Ser), c.721G>A (p.Gly241Arg), c.754T>A (p.Phe252Ile), c.1226A>G (p.Asn409Ser), c.1246G>A (p.Gly416Ser), c.1263_1317del (p.Leu422Profs*4), c.1297G>T (p.Val433Leu), c.1342G>C (p.Asp448His), c.1343A>T (p.Asp448Val), c.1448T>C (p.Leu483Pro), c.1504C>T (p.Arg502Cys), c.1505G>A (p.Arg502His), c.1603C>T (p.Arg535Cys), c.1604G>A (p.Arg535His) variants only. Rarely, sensitivity to detect these variants may be reduced. When sensitivity is reduced, zygosity may be reported as "unknown". HBA1/2: This assay is designed to detect deletions and duplications of HBA1 and/or HBA2, resulting from the -alpha20.5, --MED, --SEA, --FIL/--THAI, -alpha3.7, -alpha4.2, anti3.7 and anti4.2. Sensitivity to detect other copy number variants may be reduced. Detection of overlapping deletion and duplication events will be limited to combinations of events with significantly differing boundaries. In addition, deletion of the enhancer element HS-40 and the sequence variant, Constant Spring (NM_000517.4:c.427T>C), can be identified by this assay. MTHFR: The NM_005957.4:c.665C>T (p.Ala222Val) (aka 677C>T) and c.1286A>C (p.Glu429Ala) (aka 1298A>C) variants are not reported in our primary report. TSFM: Sequencing analysis is not offered for exon 5. VPS53: Sequencing analysis for exons 14 includes only cds +/- 5 bp. COL11A2: Deletion/duplication analysis is not offered for exon 36. CYP21A2: Analysis includes the most common variants (c.92C>T(p.Pro31Leu), c.293-13C>G (intronic), c.332_339delGAGACTAC (p.Gly111Valfs*21), c.518T>A (p.Ile173Asn), c.710T>A (p.Ile237Asn), c.713T>A (p.Val238Glu), c.719T>A (p.Met240Lys), c.844G>T (p.Val282Leu), c.923dupT (p.Leu308Phefs*6), c.955C>T (p.Gln319*), c.1069C>T(p.Arg357Trp), c.1360C>T (p.Pro454Ser) and the 30Kb deletion) as well as select rare HGMD variants only (list available upon request). Full gene duplications are reported only in the presence of a pathogenic variant(s). When a duplication and a pathogenic variant(s) is identified, phase (cis/trans) cannot be determined. Full gene deletion analysis is not offered. Sensitivity to detect these variants, if they result from complex gene conversion/fusion events, may be reduced. AIPL1: Sequencing analysis for exons 2 includes only cds +/- 10 bp. LIFR: Sequencing analysis for exons 3 includes only cds +/- 5 bp. SMN1: Systematic exon numbering is used for all genes, including SMN1, and for this reason the exon typically referred to as exon 7 in the literature (PMID: 8838816) is referred to as exon 8 in this report. This assay unambiguously detects SMN1 exon 8 copy number. The presence of the g.27134T>G variant (also known as c.*3+80T>G) is reported if SMN1 copy number = 2. SMN1 or



Patient name: DONOR 13057 DOB: [REDACTED]

Invitae #: RQ5180029

SMN2: NM_000344.3:c.*3+80T>G variant only. TYR: Deletion/duplication and sequencing analysis is not offered for exon 5. TG: Deletion/duplication analysis is not offered for exon 18. Sequencing analysis for exons 44 includes only cds +/- 0 bp. FANCL: Sequencing analysis for exons 4, 10 includes only cds +/- 10 bp. USH1C: Deletion/duplication analysis is not offered for exons 5-6. AMN: Deletion/duplication analysis is not offered for exon 1. FAH: Deletion/duplication analysis is not offered for exon 14. GALC: Deletion/duplication analysis is not offered for exon 6. ATM: Sequencing analysis for exons 6, 24, 43 includes only cds +/- 10 bp. MLC1: Sequencing analysis for exons 11 includes only cds +/- 10 bp. PEX1: Sequencing analysis for exons 16 includes only cds +/- 0 bp.

This report has been reviewed and approved by:

Christina Y. Hung, MD, FACMG
Clinical Molecular and Biochemical Geneticist

This table displays residual risks after a negative result for each of the genes and corresponding disorders. The values provided assume a negative family history and the absence of symptoms for each disorder. For genes associated with both dominant and recessive inheritance, the numbers in this table apply to the recessive condition(s) associated with the gene, unless otherwise noted. Residual risk values are provided for disorders when carrier frequency is greater than 1 in 500. For disorders with carrier frequency equal to, or less than, 1 in 500, residual risk is considered to be reduced substantially. When provided, residual risk values are inferred from published carrier frequencies, and estimated detection rates are based on testing technologies used at Invitae. Residual risks are provided only as a guide for assessing approximate risk given a negative result; values may vary based on the ethnic background(s) of an individual. For any genes marked with an asterisk*, refer to the Limitations section of the patient report for detailed coverage information. In the case of a sample-specific limitation, "N/A" indicates that a residual risk value could not be calculated. AR = autosomal recessive, XL = X-linked, AD = autosomal dominant.

DISORDER (INHERITANCE)	GENE	ETHNICITY	CARRIER FREQUENCY	DETECTION RATE	RISK TO BE A CARRIER AFTER NEGATIVE RESULT
3-hydroxy-3-methylglutaryl-CoA lyase deficiency (AR) NM_000191.2	HMGCL	Pan-ethnic	≤1 in 500	99%	Reduced
17-beta hydroxysteroid dehydrogenase 3 deficiency (AR) NM_000197.1	HSD17B3	Pan-ethnic	≤1 in 500	99%	Reduced
ABCA3-related conditions (AR) NM_001089.2	ABCA3	Pan-ethnic	1 in 277	99%	1 in 27600
ABCA4-related conditions (AR) NM_000350.2	ABCA4	Pan-ethnic	1 in 45	90%	1 in 441
ABCB4-related conditions (AR) NM_000443.3	ABCB4	Pan-ethnic	1 in 204	99%	1 in 20300
ABCB11-related conditions (AR) NM_003742.2	ABCB11	Pan-ethnic	1 in 100	99%	1 in 9900
ABCC8-related conditions (AR) NM_000352.4 When the mother is a noncarrier, but the father is a carrier, there is a residual risk for focal disease (1 in 540 for the Ashkenazi Jewish population; undetermined in other ethnic groups)	ABCC8	Pan-ethnic	1 in 177	99%	1 in 17600
Abetalipoproteinemia (AR) NM_000253.3	MTTP	Pan-ethnic	≤1 in 500	99%	Reduced
Achromatopsia (CNGB3-related) (AR) NM_019098.4	CNGB3	Pan-ethnic	1 in 93	99%	1 in 9200
ACOX1-related conditions (AR) NM_004035.6	ACOX1	Pan-ethnic	≤1 in 500	99%	Reduced
Acrodermatitis enteropathica (AR) NM_130849.3	SLC39A4	Pan-ethnic	1 in 354	99%	1 in 35300
Adenosine deaminase deficiency (AR) NM_000022.2	ADA	Pan-ethnic	1 in 224	92%	1 in 2788
ADGRV1-related conditions (AR) NM_032119.3	ADGRV1	Pan-ethnic	1 in 223	99%	1 in 22200
AHI1-related conditions (AR) NM_017651.4	AHI1	Pan-ethnic	1 in 447	99%	1 in 44600
Aicardi-Goutieres syndrome 2 (AR) NM_024570.3	RNASEH2B	Pan-ethnic	≤1 in 500	99%	Reduced
Aicardi-Goutieres syndrome 3 (AR) NM_032193.3	RNASEH2C	Pan-ethnic	≤1 in 500	99%	Reduced
Aicardi-Goutieres syndrome 4 (AR) NM_006397.2	RNASEH2A	Pan-ethnic	≤1 in 500	99%	Reduced
Aicardi-Goutieres syndrome 5 (AR) NM_015474.3	SAMHD1	Pan-ethnic	≤1 in 500	99%	Reduced
AIPL1-related conditions (AR) NM_014336.4	AIPL1 *	Pan-ethnic	1 in 408	99%	1 in 40700
Aldosterone synthase deficiency (AR) NM_000498.3	CYP11B2	Pan-ethnic	≤1 in 500	99%	Reduced
Alpha-mannosidosis (AR) NM_000528.3	MAN2B1	Pan-ethnic	1 in 354	99%	1 in 35300
Alpha-N-acetylgalactosaminidase deficiency (AR) NM_000262.2	NAGA	Pan-ethnic	≤1 in 500	99%	Reduced
Alpha-thalassemia (AR) NM_000558.4, NM_000517.4	HBA1/ HBA2 *	African-American	1 in 30	90%	1 in 291
		Asian	1 in 20	90%	1 in 191
		Caucasian	≤1 in 500	90%	Reduced

DISORDER (INHERITANCE)	GENE	ETHNICITY	CARRIER FREQUENCY	DETECTION RATE	RISK TO BE A CARRIER AFTER NEGATIVE RESULT
		Pan-ethnic	1 in 25	90%	1 in 241
Alport syndrome (COL4A3-related) (AR) NM_000091.4	COL4A3	Pan-ethnic	1 in 354	99%	1 in 35300
Alport syndrome (COL4A4-related) (AR) NM_000092.4	COL4A4	Pan-ethnic	1 in 353	99%	1 in 35200
Alström syndrome (AR) NM_015120.4	ALMS1	Pan-ethnic	≤1 in 500	99%	Reduced
Arginase deficiency (AR) NM_000045.3	ARG1	Pan-ethnic	1 in 274	99%	1 in 27300
Argininosuccinate lyase deficiency (AR) NM_000048.3	ASL	Pan-ethnic	1 in 133	90%	1 in 1321
ARL6-related conditions (AR) NM_177976.2	ARL6	Pan-ethnic	≤1 in 500	99%	Reduced
Aromatase deficiency (AR) NM_031226.2	CYP19A1	Pan-ethnic	≤1 in 500	99%	Reduced
Asparagine synthetase deficiency (AR) NM_133436.3	ASNS	Pan-ethnic	≤1 in 500	99%	Reduced
Aspartylglucosaminuria (AR) NM_000027.3	AGA	Pan-ethnic	≤1 in 500	99%	Reduced
Ataxia with vitamin E deficiency (AR) NM_000370.3	TTPA	Pan-ethnic	≤1 in 500	90%	Reduced
Ataxia-telangiectasia-like disorder (AR) NM_005591.3	MRE11	Pan-ethnic	≤1 in 500	99%	Reduced
ATM-related conditions (AR) NM_000051.3	ATM *	Pan-ethnic	1 in 100	99%	1 in 9900
ATP8B1-related conditions (AR) NM_005603.4	ATP8B1 *	Pan-ethnic	1 in 112	99%	1 in 11100
Atransferrinemia (AR) NM_001063.3	TF	Pan-ethnic	≤1 in 500	99%	Reduced
Autoimmune polyendocrinopathy with candidiasis and ectodermal dysplasia (AR) NM_000383.3	AIRE	Pan-ethnic	1 in 150	99%	1 in 14900
Autosomal recessive congenital ichthyosis (ABCA12-related) (AR) NM_173076.2	ABCA12	Pan-ethnic	≤1 in 500	99%	Reduced
Autosomal recessive congenital ichthyosis (TGM1-related) (AR) NM_000359.2	TGM1	Pan-ethnic	1 in 224	95%	1 in 4460
Autosomal recessive spastic ataxia of Charlevoix-Saguenay (AR) NM_014363.5	SACS	Pan-ethnic	≤1 in 500	99%	Reduced
Bardet-Biedl syndrome (BBS7-related) (AR) NM_176824.2	BBS7	Pan-ethnic	≤1 in 500	99%	Reduced
Bardet-Biedl syndrome (BBS9-related) (AR) NM_198428.2	BBS9 *	Pan-ethnic	≤1 in 500	99%	Reduced
Bardet-Biedl syndrome (BBS10-related) (AR) NM_024685.3	BBS10	Pan-ethnic	1 in 354	99%	1 in 35300
Bardet-Biedl syndrome (BBS12-related) (AR) NM_152618.2	BBS12	Pan-ethnic	≤1 in 500	99%	Reduced
Bartter syndrome type 1 (AR) NM_000338.2	SLC12A1	Pan-ethnic	1 in 224	99%	1 in 22300
Bartter syndrome type 2 (AR) NM_000220.4	KCNJ1	Pan-ethnic	≤1 in 500	99%	Reduced
BBS1-related conditions (AR) NM_024649.4	BBS1	Pan-ethnic	1 in 330	99%	1 in 32900
BBS2-related conditions (AR) NM_031885.3	BBS2	Pan-ethnic	≤1 in 500	99%	Reduced
BBS4-related conditions (AR) NM_033028.4	BBS4	Pan-ethnic	≤1 in 500	99%	Reduced
BBS5-related conditions (AR) NM_152384.2	BBS5	Pan-ethnic	≤1 in 500	99%	Reduced
BCS1L-related conditions (AR) NM_004328.4	BCS1L	Pan-ethnic	≤1 in 500	99%	Reduced

DISORDER (INHERITANCE)	GENE	ETHNICITY	CARRIER FREQUENCY	DETECTION RATE	RISK TO BE A CARRIER AFTER NEGATIVE RESULT
Beta-ketothiolase deficiency (AR) NM_000019.3	ACAT1	Pan-ethnic	≤1 in 500	99%	Reduced
Beta-mannosidosis (AR) NM_005908.3	MANBA	Pan-ethnic	≤1 in 500	99%	Reduced
Biopterin-deficient hyperphenylalaninemia (PCBD1-related) (AR) NM_000281.3	PCBD1	Pan-ethnic	≤1 in 500	99%	Reduced
Biopterin-deficient hyperphenylalaninemia (PTS-related) (AR) NM_000317.2	PTS	Pan-ethnic	1 in 433	99%	1 in 43200
Biopterin-deficient hyperphenylalaninemia (QDPR-related) (AR) NM_000320.2	QDPR	Pan-ethnic	≤1 in 500	99%	Reduced
Biotin-responsive basal ganglia disease (AR) NM_025243.3	SLC19A3	Pan-ethnic	≤1 in 500	99%	Reduced
Biotinidase deficiency (AR) NM_000060.3	BTD	Pan-ethnic	1 in 125	99%	1 in 12400
Bloom syndrome (AR) NM_000057.3	BLM	Pan-ethnic	≤1 in 500	99%	Reduced
BRIP1-related conditions (AR) NM_032043.2	BRIP1	Pan-ethnic	≤1 in 500	99%	Reduced
Brittle cornea syndrome (PRDM5-related) (AR) NM_018699.3	PRDM5	Pan-ethnic	≤1 in 500	99%	Reduced
Brittle cornea syndrome (ZNF469-related) (AR) NM_001127464.2	ZNF469	Pan-ethnic	≤1 in 500	99%	Reduced
BSND-related conditions (AR) NM_057176.2	BSND	Pan-ethnic	≤1 in 500	99%	Reduced
Canavan disease (AR) NM_000049.2	ASPA	Pan-ethnic	1 in 159	99%	1 in 15800
Carbamoyl phosphate synthetase I deficiency (AR) NM_001875.4	CPS1	Pan-ethnic	≤1 in 500	99%	Reduced
Cardioencephalomyopathy (AR) NM_005138.2	SCO2	Pan-ethnic	1 in 387	99%	1 in 38600
Carnitine palmitoyltransferase I deficiency (AR) NM_001876.3	CPT1A	Pan-ethnic	≤1 in 500	99%	Reduced
Carnitine palmitoyltransferase II deficiency (AR) NM_000098.2	CPT2	Pan-ethnic	1 in 182	99%	1 in 18100
Carnitine-acylcarnitine translocase deficiency (AR) NM_000387.5	SLC25A20	Pan-ethnic	≤1 in 500	99%	Reduced
Carpenter syndrome (RAB23-related) (AR) NM_183227.2	RAB23	Pan-ethnic	≤1 in 500	99%	Reduced
Cartilage-hair hypoplasia-anauxetic dysplasia spectrum disorders (AR) NR_003051.3	RMRP	Pan-ethnic	≤1 in 500	99%	Reduced
Catecholaminergic polymorphic ventricular tachycardia (CASQ2-related) (AR) NM_001232.3	CASQ2	Pan-ethnic	1 in 224	99%	1 in 22300
CC2D2A-related conditions (AR) NM_001080522.2	CC2D2A	Pan-ethnic	1 in 426	99%	1 in 42500
CDH23-related conditions (AR) NM_022124.5	CDH23	Pan-ethnic	1 in 202	95%	1 in 4020
CEP290-related conditions (AR) NM_025114.3	CEP290	Pan-ethnic	1 in 185	99%	1 in 18400
Cerebellar ataxia, intellectual disability, and dysequilibrium syndrome 1 (AR) NM_003383.4	VLDLR	Pan-ethnic	≤1 in 500	99%	Reduced
Cerebral dysgenesis, neuropathy, ichthyosis, and keratoderma (AR) NM_004782.3	SNAP29	Pan-ethnic	≤1 in 500	99%	Reduced
Cerebrotendinous xanthomatosis (AR) NM_000784.3	CYP27A1	Pan-ethnic	1 in 112	98%	1 in 5550
CERKL-related conditions (AR) NM_001030311.2	CERKL	Pan-ethnic	1 in 137	99%	1 in 13600

DISORDER (INHERITANCE)	GENE	ETHNICITY	CARRIER FREQUENCY	DETECTION RATE	RISK TO BE A CARRIER AFTER NEGATIVE RESULT
CFTR-related conditions (AR) NM_000492.3	CFTR *	Pan-ethnic - classic CF	1 in 45	99%	1 in 4400
		Pan-ethnic - classic CF and CFTR-related disorders	1 in 9	99%	1 in 800
Charcot-Marie-Tooth disease type 4D (AR) NM_006096.3	NDRG1	Pan-ethnic	≤1 in 500	99%	Reduced
Chediak-Higashi syndrome (AR) NM_000081.3	LYST	Pan-ethnic	≤1 in 500	99%	Reduced
Childhood-onset dystonia with optic atrophy and basal ganglia abnormalities (AR) NM_016011.3	MECR	Pan-ethnic	≤1 in 500	99%	Reduced
Chorea-acanthocytosis (AR) NM_033305.2	VPS13A *	Pan-ethnic	≤1 in 500	97%	Reduced
Chronic granulomatous disease (CYBA-related) (AR) NM_000101.3	CYBA	Pan-ethnic	≤1 in 500	99%	Reduced
Chronic granulomatous disease (NCF2-related) (AR) NM_000433.3	NCF2	Pan-ethnic	≤1 in 500	99%	Reduced
Citrin deficiency (AR) NM_014251.2	SLC25A13	Pan-ethnic	1 in 313	99%	1 in 31200
Citrullinemia type 1 (AR) NM_000050.4	ASS1	Pan-ethnic	1 in 120	96%	1 in 2975
CLN3-related conditions (AR) NM_001042432.1	CLN3	Pan-ethnic	1 in 230	99%	1 in 22900
CLRN1-related conditions (AR) NM_174878.2	CLRN1	Pan-ethnic	≤1 in 500	99%	Reduced
Cobalamin C deficiency (AR) NM_015506.2	MMACHC	Pan-ethnic	1 in 123	99%	1 in 12200
Cobalamin D deficiency (AR) NM_015702.2	MMADHC	Pan-ethnic	≤1 in 500	99%	Reduced
Cobalamin F deficiency (AR) NM_018368.3	LMBRD1	Pan-ethnic	≤1 in 500	99%	Reduced
Cockayne syndrome A (AR) NM_000082.3	ERCC8	Pan-ethnic	≤1 in 500	99%	Reduced
Cockayne syndrome B (AR) NM_000124.3	ERCC6	Pan-ethnic	1 in 377	99%	1 in 37600
Cohen syndrome (AR) NM_017890.4	VPS13B	Pan-ethnic	≤1 in 500	99%	Reduced
COL11A2-related conditions (AR) NM_080680.2	COL11A2 *	Pan-ethnic	≤1 in 500	99%	Reduced
COL17A1-related conditions (AR) NM_000494.3	COL17A1	Pan-ethnic	≤1 in 500	99%	Reduced
Combined malonic and methylmalonic aciduria (AR) NM_174917.4	ACSF3	Pan-ethnic	1 in 87	99%	1 in 8600
Combined oxidative phosphorylation deficiency 1 (AR) NM_024996.5	GFM1	Pan-ethnic	≤1 in 500	99%	Reduced
Combined oxidative phosphorylation deficiency 3 (AR) NM_001172696.1	TSFM *	Pan-ethnic	≤1 in 500	93%	Reduced
Combined pituitary hormone deficiency (LHX3-related) (AR) NM_014564.4	LHX3	Pan-ethnic	≤1 in 500	99%	Reduced
Combined pituitary hormone deficiency (POU1F1-related) (AR) NM_000306.3	POU1F1	Pan-ethnic	≤1 in 500	99%	Reduced
Combined pituitary hormone deficiency (PROP1-related) (AR) NM_006261.4	PROP1	Pan-ethnic	1 in 45	98%	1 in 2200
Congenital adrenal hyperplasia due to 3-beta-hydroxysteroid dehydrogenase deficiency (AR) NM_000198.3	HSD3B2	Pan-ethnic	≤1 in 500	99%	Reduced
Congenital adrenal insufficiency (AR) NM_000781.2	CYP11A1	Pan-ethnic	≤1 in 500	99%	Reduced
Congenital chronic diarrhea (DGAT1-related) (AR) NM_012079.5	DGAT1	Pan-ethnic	≤1 in 500	99%	Reduced

DISORDER (INHERITANCE)	GENE	ETHNICITY	CARRIER FREQUENCY	DETECTION RATE	RISK TO BE A CARRIER AFTER NEGATIVE RESULT
Congenital disorder of glycosylation (SLC35A3-related) (AR) NM_012243.2	SLC35A3	Pan-ethnic	≤1 in 500	99%	Reduced
Congenital disorder of glycosylation type Ia (AR) NM_000303.2	PMM2	Pan-ethnic	1 in 190	99%	1 in 18900
Congenital disorder of glycosylation type Ib (AR) NM_002435.2	MPI	Pan-ethnic	≤1 in 500	99%	Reduced
Congenital disorder of glycosylation type Ic (AR) NM_013339.3	ALG6	Pan-ethnic	≤1 in 500	99%	Reduced
Congenital disorder of glycosylation type Iv (AR) NM_018297.3	NGLY1	Pan-ethnic	≤1 in 500	99%	Reduced
Congenital dyserythropoietic anemia type II (AR) NM_006363.4	SEC23B	Pan-ethnic	≤1 in 500	99%	Reduced
Congenital hydrocephalus-1 (AR) NM_001080414.3	CCDC88C	Pan-ethnic	≤1 in 500	99%	Reduced
Congenital hypothyroidism (TSHB-related) (AR) NM_000549.4	TSHB	Pan-ethnic	≤1 in 500	99%	Reduced
Congenital insensitivity to pain with anhidrosis (AR) NM_001012331.1	NTRK1	Pan-ethnic	≤1 in 500	99%	Reduced
Congenital myasthenic syndrome (CHAT-related) (AR) NM_020549.4	CHAT	Pan-ethnic	≤1 in 500	99%	Reduced
Congenital myasthenic syndrome (CHRNE-related) (AR) NM_000080.3	CHRNE	Pan-ethnic	1 in 200	99%	1 in 19900
Congenital nephrotic syndrome type 1 (AR) NM_004646.3	NPHS1	Pan-ethnic	≤1 in 500	99%	Reduced
Congenital nephrotic syndrome type 2 (AR) NM_014625.3	NPHS2	Pan-ethnic	≤1 in 500	99%	Reduced
Congenital secretory chloride diarrhea (AR) NM_000111.2	SLC26A3	Pan-ethnic	≤1 in 500	99%	Reduced
Corneal dystrophy and perceptive deafness (AR) NM_032034.3	SLC4A11	Pan-ethnic	≤1 in 500	99%	Reduced
CRB1-related conditions (AR) NM_201253.2	CRB1	Pan-ethnic	1 in 112	99%	1 in 11100
CTSC-related conditions (AR) NM_001814.5	CTSC	Pan-ethnic	1 in 250	99%	1 in 24900
CYP1B1-related conditions (AR) NM_000104.3	CYP1B1	Pan-ethnic	1 in 79	99%	1 in 7800
CYP7B1-related conditions (AR) NM_004820.3	CYP7B1	Pan-ethnic	≤1 in 500	99%	Reduced
CYP11B1-related conditions (AR) NM_000497.3	CYP11B1	Pan-ethnic	1 in 194	99%	1 in 19300
CYP17A1-related conditions (AR) NM_000102.3	CYP17A1	Pan-ethnic	≤1 in 500	99%	Reduced
Cystinosis (AR) NM_004937.2	CTNS	Pan-ethnic	1 in 158	99%	1 in 15700
Cytochrome P450 oxidoreductase deficiency (AR) NM_000941.2	POR	Pan-ethnic	1 in 158	99%	1 in 15700
Desbuquois dysplasia type 1 (AR) NM_138793.3	CANT1	Pan-ethnic	≤1 in 500	99%	Reduced
Developmental and epileptic encephalopathy (CAD-related) (AR) NM_004341.4	CAD	Pan-ethnic	≤1 in 500	99%	Reduced
DGUOK-related conditions (AR) NM_080916.2	DGUOK	Pan-ethnic	≤1 in 500	99%	Reduced
DHDDS-related conditions (AR) NM_024887.3	DHDDS	Pan-ethnic	≤1 in 500	99%	Reduced
Dihydrolipoamide dehydrogenase deficiency (AR) NM_000108.4	DLD	Pan-ethnic	≤1 in 500	99%	Reduced
Distal renal tubular acidosis with deafness (ATP6V1B1-related) (AR) NM_001692.3	ATP6V1B1	Pan-ethnic	≤1 in 500	99%	Reduced
DOK7-related conditions (AR) NM_173660.4	DOK7	Pan-ethnic	1 in 115	99%	1 in 11400

DISORDER (INHERITANCE)	GENE	ETHNICITY	CARRIER FREQUENCY	DETECTION RATE	RISK TO BE A CARRIER AFTER NEGATIVE RESULT
Donnai-Barrow syndrome (AR) NM_004525.2	LRP2	Pan-ethnic	≤1 in 500	99%	Reduced
Dubin-Johnson syndrome (AR) NM_000392.4	ABCC2 *	Pan-ethnic	≤1 in 500	99%	Reduced
DUOX2-related conditions (AR) NM_014080.4	DUOX2 *	Pan-ethnic	1 in 58	91%	1 in 634
DYNC2H1-related conditions (AR) NM_001080463.1	DYNC2H1	Pan-ethnic	1 in 224	99%	1 in 22300
DYSF-related conditions (AR) NM_003494.3	DYSF	Pan-ethnic	1 in 311	99%	1 in 31000
Dyskeratosis congenita spectrum disorders (RTEL1-related) (AR) NM_001283009.1	RTEL1	Pan-ethnic	≤1 in 500	99%	Reduced
Dyskeratosis congenita spectrum disorders (TERT-related) (AR) NM_198253.2	TERT	Pan-ethnic	≤1 in 500	99%	Reduced
Dystrophic epidermolysis bullosa (AR) NM_000094.3	COL7A1	Pan-ethnic	1 in 370	97%	1 in 12300
Ehlers-Danlos syndrome, dermatosparaxis type (AR) NM_014244.4	ADAMTS2	Pan-ethnic	≤1 in 500	99%	Reduced
Ehlers-Danlos syndrome, kyphoscoliotic type (AR) NM_000302.3	PLOD1	Pan-ethnic	1 in 150	99%	1 in 14900
Ellis-van Creveld syndrome (EVC-related) (AR) NM_153717.2	EVC	Pan-ethnic	1 in 220	99%	1 in 21900
Epidermolysis bullosa with pyloric atresia (ITGB4-related) (AR) NM_001005731.2	ITGB4	Pan-ethnic	1 in 393	99%	1 in 39200
Epimerase deficiency galactosemia (AR) NM_000403.3	GALE *	Pan-ethnic	≤1 in 500	99%	Reduced
ERCC2-related conditions (AR) NM_000400.3	ERCC2	Pan-ethnic	≤1 in 500	99%	Reduced
Ethylmalonic encephalopathy (AR) NM_014297.3	ETHE1	Pan-ethnic	≤1 in 500	99%	Reduced
EVC2-related conditions (AR) NM_147127.4	EVC2	Pan-ethnic	1 in 199	99%	1 in 19800
Familial chylomicronemia syndrome (AR) NM_000237.2	LPL	Pan-ethnic	≤1 in 500	99%	Reduced
Familial dysautonomia (AR) NM_003640.3	ELP1	Pan-ethnic	≤1 in 500	99%	Reduced
Familial hemophagocytic lymphohistiocytosis type 2 (AR) NM_001083116.1	PRF1	Pan-ethnic	1 in 177	99%	1 in 17600
Familial hemophagocytic lymphohistiocytosis type 3 (AR) NM_199242.2	UNC13D	Pan-ethnic	1 in 177	93%	1 in 2515
Familial hemophagocytic lymphohistiocytosis type 4 (AR) NM_003764.3	STX11	Pan-ethnic	1 in 224	99%	1 in 22300
Familial hemophagocytic lymphohistiocytosis type 5 (AR) NM_006949.3	STXBP2	Pan-ethnic	1 in 224	99%	1 in 22300
Familial hypercholesterolemia (LDLR-related) (AD) NM_000527.4	LDLR	Pan-ethnic	1 in 250	99%	1 in 24900
Familial hypercholesterolemia (LDLRAP1-related) (AR) NM_015627.2	LDLRAP1	Pan-ethnic	≤1 in 500	99%	Reduced
Fanconi anemia type A (AR) NM_000135.2	FANCA	Pan-ethnic	1 in 345	99%	1 in 34400
Fanconi anemia type C (AR) NM_000136.2	FANCC	Pan-ethnic	1 in 417	99%	1 in 41600
Fanconi anemia type D2 (AR) NM_033084.3	FANCD2 *	Pan-ethnic	≤1 in 500	94%	Reduced
Fanconi anemia type E (AR) NM_021922.2	FANCE	Pan-ethnic	≤1 in 500	99%	Reduced
Fanconi anemia type G (AR) NM_004629.1	FANCG	Pan-ethnic	≤1 in 500	99%	Reduced
Fanconi anemia type I (AR) NM_001113378.1	FANCI	Pan-ethnic	≤1 in 500	99%	Reduced

DISORDER (INHERITANCE)	GENE	ETHNICITY	CARRIER FREQUENCY	DETECTION RATE	RISK TO BE A CARRIER AFTER NEGATIVE RESULT
Fanconi anemia type L (AR) NM_018062.3	FANCL *	Pan-ethnic	≤1 in 500	99%	Reduced
FH-related conditions (AR) NM_000143.3	FH *	Pan-ethnic	≤1 in 500	99%	Reduced
FKBP10-related conditions (AR) NM_021939.3	FKBP10	Pan-ethnic	≤1 in 500	99%	Reduced
Foveal hypoplasia (SLC38A8-related) (AR) NM_001080442.2	SLC38A8	Pan-ethnic	≤1 in 500	99%	Reduced
FOXN1-related conditions (AR) NM_003593.2	FOXN1	Pan-ethnic	≤1 in 500	99%	Reduced
Fraser syndrome (FRAS1-related) (AR) NM_025074.6	FRAS1	Pan-ethnic	1 in 316	99%	1 in 31500
Fraser syndrome (FREM2-related) (AR) NM_207361.5	FREM2	Pan-ethnic	≤1 in 500	99%	Reduced
Fraser syndrome (GRIP1-related) (AR) NM_021150.3	GRIP1	Pan-ethnic	1 in 447	99%	1 in 44600
Fructose-1,6-bisphosphatase deficiency (AR) NM_000507.3	FBP1	Pan-ethnic	≤1 in 500	99%	Reduced
Fucosidosis (AR) NM_000147.4	FUCA1	Pan-ethnic	≤1 in 500	99%	Reduced
Galactokinase deficiency galactosemia (AR) NM_000154.1	GALK1	Pan-ethnic	1 in 122	99%	1 in 12100
Galactosemia (GALT-related) (AR) NM_000155.3	GALT	Pan-ethnic	1 in 100	99%	1 in 9900
Galactosialidosis (AR) NM_000308.3	CTSA	Pan-ethnic	≤1 in 500	99%	Reduced
GATM-related conditions (AR) NM_001482.2	GATM	Pan-ethnic	≤1 in 500	99%	Reduced
GBA-related conditions including Gaucher disease (AR) NM_001005741.2	GBA *	Ashkenazi Jewish	1 in 15	94%	1 in 234
		Pan-ethnic	1 in 158	72%	1 in 561
GBE1-related conditions (AR) NM_000158.3	GBE1	Pan-ethnic	1 in 387	99%	1 in 38600
GCH1-related conditions (AR) NM_000161.2	GCH1	Pan-ethnic	≤1 in 500	99%	Reduced
GDF5-related conditions (AR) NM_000557.4	GDF5	Pan-ethnic	≤1 in 500	99%	Reduced
Geroderma osteodysplastica (AR) NM_152281.2	GORAB	Pan-ethnic	≤1 in 500	99%	Reduced
GHR-related conditions (AR) NM_000163.4	GHR *	Pan-ethnic	≤1 in 500	98%	Reduced
GJB2-related conditions (AR) NM_004004.5	GJB2	Pan-ethnic	1 in 50	99%	1 in 4900
GLB1-related conditions (AR) NM_000404.2	GLB1	Pan-ethnic	1 in 158	99%	1 in 15700
GLE1-related conditions (AR) NM_001003722.1	GLE1	Pan-ethnic	≤1 in 500	99%	Reduced
Glutaric acidemia type I (AR) NM_000159.3	GCDH	Pan-ethnic	1 in 87	99%	1 in 8600
Glutaric acidemia type IIA (AR) NM_000126.3	ETFA	Pan-ethnic	≤1 in 500	99%	Reduced
Glutaric acidemia type IIB (AR) NM_001985.2	ETFB	Pan-ethnic	≤1 in 500	99%	Reduced
Glutaric acidemia type IIC (AR) NM_004453.3	ETFDH	Pan-ethnic	1 in 250	99%	1 in 24900
Glutathione synthetase deficiency (AR) NM_000178.2	GSS	Pan-ethnic	≤1 in 500	99%	Reduced
Glycine encephalopathy (AMT-related) (AR) NM_000481.3	AMT	Pan-ethnic	1 in 325	99%	1 in 32400
Glycine encephalopathy (GLDC-related) (AR) NM_000170.2	GLDC	Pan-ethnic	1 in 165	99%	1 in 16400
Glycogen storage disease type Ia (AR) NM_000151.3	G6PC	Pan-ethnic	1 in 177	95%	1 in 3520

DISORDER (INHERITANCE)	GENE	ETHNICITY	CARRIER FREQUENCY	DETECTION RATE	RISK TO BE A CARRIER AFTER NEGATIVE RESULT
Glycogen storage disease type II (Pompe disease) (AR) NM_000152.3	GAA	Pan-ethnic	1 in 100	99%	1 in 9900
Glycogen storage disease type III (AR) NM_000642.2	AGL	Pan-ethnic	1 in 159	95%	1 in 3160
Glycogen storage disease type IXb (AR) NM_000293.2	PHKB	Pan-ethnic	≤1 in 500	99%	Reduced
Glycogen storage disease type IXc (AR) NM_000294.2	PHKG2	Pan-ethnic	≤1 in 500	99%	Reduced
Glycogen storage disease type V (AR) NM_005609.3	PYGM	Pan-ethnic	1 in 171	99%	1 in 17000
Glycogen storage disease type VII (AR) NM_000289.5	PFKM	Pan-ethnic	≤1 in 500	99%	Reduced
GM3 synthase deficiency (AR) NM_003896.3	ST3GAL5	Pan-ethnic	≤1 in 500	99%	Reduced
GNE-related conditions (AR) NM_001128227.2	GNE *	Pan-ethnic	1 in 179	99%	1 in 17800
GNPTAB-related conditions (AR) NM_024312.4	GNPTAB	Pan-ethnic	1 in 200	99%	1 in 19900
Guanidinoacetate methyltransferase deficiency (AR) NM_000156.5	GAMT	Pan-ethnic	≤1 in 500	99%	Reduced
GUCY2D-related conditions (AR) NM_000180.3	GUCY2D	Pan-ethnic	1 in 204	99%	1 in 20300
Gyrate atrophy of the choroid and retina (AR) NM_000274.3	OAT *	Pan-ethnic	≤1 in 500	99%	Reduced
HADHA-related conditions (AR) NM_000182.4	HADHA	Pan-ethnic	1 in 350	99%	1 in 34900
HBB-related hemoglobinopathies (AR) NM_000518.4	HBB	Pan-ethnic	1 in 49	99%	1 in 4800
Heme oxygenase 1 deficiency (AR) NM_002133.2	HMOX1	Pan-ethnic	≤1 in 500	99%	Reduced
Hemolytic anemia, CD59-mediated (AR) NM_203330.2	CD59	Pan-ethnic	≤1 in 500	99%	Reduced
Hereditary fructose intolerance (AR) NM_000035.3	ALDOB	Pan-ethnic	1 in 122	99%	1 in 12100
Hereditary hemochromatosis type 2 (HAMP-related) (AR) NM_021175.2	HAMP	Pan-ethnic	≤1 in 500	99%	Reduced
Hereditary hemochromatosis type 2 (HJV-related) (AR) NM_213653.3	HJV	Pan-ethnic	≤1 in 500	99%	Reduced
Hereditary hemochromatosis type 3 (AR) NM_003227.3	TFR2	Pan-ethnic	≤1 in 500	99%	Reduced
Hermansky-Pudlak syndrome type 1 (AR) NM_000195.4	HPS1	Pan-ethnic	≤1 in 500	99%	Reduced
Hermansky-Pudlak syndrome type 3 (AR) NM_032383.4	HPS3	Pan-ethnic	≤1 in 500	99%	Reduced
Hermansky-Pudlak syndrome type 4 (AR) NM_022081.5	HPS4	Pan-ethnic	≤1 in 500	99%	Reduced
Hermansky-Pudlak syndrome type 5 (AR) NM_181507.1	HPS5	Pan-ethnic	≤1 in 500	99%	Reduced
Hermansky-Pudlak syndrome type 6 (AR) NM_024747.5	HPS6	Pan-ethnic	≤1 in 500	99%	Reduced
Hermansky-Pudlak syndrome type 8 (AR) NM_212550.4	BLOC1S3	Pan-ethnic	≤1 in 500	99%	Reduced
Hermansky-Pudlak syndrome type 9 (AR) NM_012388.3	BLOC1S6	Pan-ethnic	≤1 in 500	99%	Reduced
HGSNAT-related conditions (AR) NM_152419.2	HGSNAT	Pan-ethnic	≤1 in 500	99%	Reduced
Holocarboxylase synthetase deficiency (AR) NM_000411.6	HLCS	Pan-ethnic	1 in 224	99%	1 in 22300
Homocystinuria due to cobalamin E deficiency (AR) NM_002454.2	MTRR	Pan-ethnic	≤1 in 500	99%	Reduced
Homocystinuria due to cobalamin G deficiency (AR) NM_000254.2	MTR	Pan-ethnic	≤1 in 500	99%	Reduced

DISORDER (INHERITANCE)	GENE	ETHNICITY	CARRIER FREQUENCY	DETECTION RATE	RISK TO BE A CARRIER AFTER NEGATIVE RESULT
Homocystinuria due to cystathionine beta-synthase deficiency (AR) NM_000071.2	CBS	Pan-ethnic	1 in 224	99%	1 in 22300
Homocystinuria due to MTHFR deficiency (AR) NM_005957.4	MTHFR *	Pan-ethnic	≤1 in 500	99%	Reduced
HSD17B4-related conditions (AR) NM_000414.3	HSD17B4	Pan-ethnic	1 in 158	99%	1 in 15700
Hydrolethals syndrome type 1 (AR) NM_145014.2	HYLS1	Pan-ethnic	≤1 in 500	99%	Reduced
Hyper-IgM immunodeficiency (CD40-related) (AR) NM_001250.5	CD40	Pan-ethnic	≤1 in 500	99%	Reduced
Hyperornithinemia-hyperammonemia-homocitrullinuria syndrome (AR) NM_014252.3	SLC25A15	Pan-ethnic	≤1 in 500	99%	Reduced
Hyperphosphatemic familial tumoral calcinosis (GALNT3-related) (AR) NM_004482.3	GALNT3	Pan-ethnic	≤1 in 500	99%	Reduced
Hypomyelinating leukodystrophy-12 (AR) NM_021729.5	VPS11	Pan-ethnic	≤1 in 500	99%	Reduced
Hypophosphatasia (AR) NM_000478.5	ALPL	Pan-ethnic	1 in 150	95%	1 in 2980
Ichthyosis prematurity syndrome (AR) NM_005094.3	SLC27A4	Pan-ethnic	≤1 in 500	99%	Reduced
IGHMBP2-related conditions (AR) NM_002180.2	IGHMBP2	Pan-ethnic	≤1 in 500	99%	Reduced
IKBKB-related conditions (AR) NM_001556.2	IKBKB	Pan-ethnic	≤1 in 500	99%	Reduced
Imerslund-Gräsbeck syndrome (AR) NM_030943.3	AMN *	Pan-ethnic	≤1 in 500	99%	Reduced
Immunodeficiency-centromeric instability-facial anomalies syndrome 1 (AR) NM_006892.3	DNMT3B	Pan-ethnic	≤1 in 500	99%	Reduced
Immunodeficiency-centromeric instability-facial anomalies syndrome 2 (AR) NM_014797.2	ZBTB24	Pan-ethnic	≤1 in 500	99%	Reduced
Isolated ectopia lentis (AR) NM_019032.5	ADAMTSL4	Pan-ethnic	≤1 in 500	99%	Reduced
Isovaleric acidemia (AR) NM_002225.3	IVD	Pan-ethnic	1 in 250	99%	1 in 24900
ITGB3-related conditions (AR) NM_000212.2	ITGB3	Pan-ethnic	≤1 in 500	99%	Reduced
Johanson-Blizzard syndrome (AR) NM_174916.2	UBR1	Pan-ethnic	1 in 250	99%	1 in 24900
Joubert syndrome and related disorders (MKS1-related) (AR) NM_017777.3	MKS1	Pan-ethnic	1 in 260	95%	1 in 5180
Joubert syndrome and related disorders (RPGRIP1L-related) (AR) NM_015272.2	RPGRIP1L	Pan-ethnic	1 in 259	95%	1 in 5160
Joubert syndrome and related disorders (TMEM216-related) (AR) NM_001173990.2	TMEM216	Pan-ethnic	≤1 in 500	99%	Reduced
Junctional epidermolysis bullosa (LAMC2-related) (AR) NM_005562.2	LAMC2	Pan-ethnic	≤1 in 500	99%	Reduced
Junctional epidermolysis bullosa with pyloric atresia (ITGA6-related) (AR) NM_000210.3	ITGA6	Pan-ethnic	≤1 in 500	99%	Reduced
KCNJ11-related conditions (AR) NM_000525.3	KCNJ11	Pan-ethnic	≤1 in 500	99%	Reduced
Krabbe disease (AR) NM_000153.3	GALC *	Pan-ethnic	1 in 158	99%	1 in 15700
LAMA2-related muscular dystrophy (AR) NM_000426.3	LAMA2	Pan-ethnic	1 in 87	99%	1 in 8600

DISORDER (INHERITANCE)	GENE	ETHNICITY	CARRIER FREQUENCY	DETECTION RATE	RISK TO BE A CARRIER AFTER NEGATIVE RESULT
LAMA3-related conditions (AR) NM_000227.4	LAMA3	Pan-ethnic	≤1 in 500	99%	Reduced
LAMB3-related conditions (AR) NM_000228.2	LAMB3	Pan-ethnic	1 in 317	99%	1 in 31600
Leber congenital amaurosis 5 (AR) NM_181714.3	LCA5	Pan-ethnic	≤1 in 500	97%	Reduced
Leukoencephalopathy with vanishing white matter (EIF2B1-related) (AR) NM_001414.3	EIF2B1	Pan-ethnic	≤1 in 500	99%	Reduced
Leukoencephalopathy with vanishing white matter (EIF2B2-related) (AR) NM_014239.3	EIF2B2	Pan-ethnic	≤1 in 500	99%	Reduced
Leukoencephalopathy with vanishing white matter (EIF2B3-related) (AR) NM_020365.4	EIF2B3	Pan-ethnic	≤1 in 500	99%	Reduced
Leukoencephalopathy with vanishing white matter (EIF2B4-related) (AR) NM_015636.3	EIF2B4	Pan-ethnic	≤1 in 500	99%	Reduced
Leukoencephalopathy with vanishing white matter (EIF2B5-related) (AR) NM_003907.2	EIF2B5	Pan-ethnic	≤1 in 500	99%	Reduced
LIG4 syndrome (AR) NM_002312.3	LIG4	Pan-ethnic	≤1 in 500	99%	Reduced
Limb-girdle muscular dystrophy (CAPN3-related) (AR) NM_000070.2	CAPN3	Pan-ethnic	1 in 134	99%	1 in 13300
Limb-girdle muscular dystrophy type 2C (AR) NM_000231.2	SGCG	Pan-ethnic	≤1 in 500	99%	Reduced
Limb-girdle muscular dystrophy type 2D (AR) NM_000023.2	SGCA	Pan-ethnic	≤1 in 500	99%	Reduced
Limb-girdle muscular dystrophy type 2E (AR) NM_000232.4	SGCB	Pan-ethnic	≤1 in 500	92%	Reduced
Limb-girdle muscular dystrophy type 2F (AR) NM_000337.5	SGCD	Pan-ethnic	≤1 in 500	99%	Reduced
Lipoid congenital adrenal hyperplasia (AR) NM_000349.2	STAR	Pan-ethnic	≤1 in 500	99%	Reduced
LRAT-related conditions (AR) NM_004744.4	LRAT	Pan-ethnic	1 in 296	99%	1 in 29500
Lysinuric protein intolerance (AR) NM_001126106.2	SLC7A7	Pan-ethnic	≤1 in 500	99%	Reduced
Lysosomal acid lipase deficiency (AR) NM_000235.3	LIPA	Pan-ethnic	1 in 359	94%	1 in 5967
Major histocompatibility complex class II deficiency (CIITA-related) (AR) NM_000246.3	CIITA	Pan-ethnic	≤1 in 500	99%	Reduced
Malonyl-CoA decarboxylase deficiency (AR) NM_012213.2	MLYCD	Pan-ethnic	≤1 in 500	99%	Reduced
Maple syrup urine disease type 1A (AR) NM_000709.3	BCKDHA	Pan-ethnic	1 in 373	99%	1 in 37200
Maple syrup urine disease type 1B (AR) NM_183050.2	BCKDHB	Pan-ethnic	1 in 346	99%	1 in 34500
Maple syrup urine disease type 2 (AR) NM_001918.3	DBT	Pan-ethnic	≤1 in 500	99%	Reduced
Medium-chain acyl-CoA dehydrogenase deficiency (AR) NM_000016.5	ACADM	Pan-ethnic	1 in 66	99%	1 in 6500
Medium/short-chain 3-hydroxyacyl-CoA dehydrogenase deficiency (AR) NM_005327.4	HADH	Pan-ethnic	≤1 in 500	99%	Reduced
MEDNIK syndrome (AR) NM_001283.3	AP1S1	Pan-ethnic	≤1 in 500	99%	Reduced
Megalencephalic leukoencephalopathy with subcortical cysts 1 (AR) NM_015166.3	MLC1 *	Pan-ethnic	≤1 in 500	99%	Reduced

DISORDER (INHERITANCE)	GENE	ETHNICITY	CARRIER FREQUENCY	DETECTION RATE	RISK TO BE A CARRIER AFTER NEGATIVE RESULT
Metabolic crises with rhabdomyolysis, cardiac arrhythmias and neurodegeneration (AR) NM_152906.6	TANGO2	Pan-ethnic	≤1 in 500	99%	Reduced
Metachromatic leukodystrophy (ARSA-related) (AR) NM_000487.5	ARSA	Pan-ethnic	1 in 100	95%	1 in 1980
Methylmalonic acidemia (MCEE-related) (AR) NM_032601.3	MCEE	Pan-ethnic	≤1 in 500	99%	Reduced
Methylmalonic acidemia (MMAA-related) (AR) NM_172250.2	MMAA	Pan-ethnic	1 in 316	97%	1 in 10500
Methylmalonic acidemia (MMAB-related) (AR) NM_052845.3	MMAB	Pan-ethnic	1 in 456	98%	1 in 22750
Methylmalonic acidemia (MUT-related) (AR) NM_000255.3	MUT	Pan-ethnic	1 in 204	96%	1 in 5075
MFSD8-related conditions (AR) NM_152778.2	MFSD8	Pan-ethnic	≤1 in 500	99%	Reduced
Microcephalic osteodysplastic primordial dwarfism type II (AR) NM_006031.5	PCNT	Pan-ethnic	≤1 in 500	99%	Reduced
Microcephaly, postnatal progressive, with seizures and brain atrophy (AR) NM_004268.4	MED17	Pan-ethnic	≤1 in 500	99%	Reduced
Mitochondrial complex I deficiency 1 (AR) NM_002495.3	NDUFS4	Pan-ethnic	≤1 in 500	99%	Reduced
Mitochondrial complex I deficiency 3 (AR) NM_024407.4	NDUFS7	Pan-ethnic	1 in 387	99%	1 in 38600
Mitochondrial complex I deficiency 4 (AR) NM_007103.3	NDUFV1	Pan-ethnic	1 in 387	99%	1 in 38600
Mitochondrial complex I deficiency 9 (AR) NM_004553.4	NDUFS6	Pan-ethnic	≤1 in 500	99%	Reduced
Mitochondrial complex I deficiency 10 (AR) NM_174889.4	NDUFAF2	Pan-ethnic	1 in 387	99%	1 in 38600
Mitochondrial complex I deficiency 16 (AR) NM_024120.4	NDUFAF5	Pan-ethnic	≤1 in 500	99%	Reduced
Mitochondrial complex I deficiency 19 (AR) NM_017547.3	FOXRED1	Pan-ethnic	1 in 376	99%	1 in 37500
Mitochondrial complex I deficiency 20/ACAD9 deficiency (AR) NM_014049.4	ACAD9	Pan-ethnic	≤1 in 500	99%	Reduced
Mitochondrial complex IV deficiency 6 (AR) NM_004376.6	COX15	Pan-ethnic	≤1 in 500	99%	Reduced
Mitochondrial complex IV deficiency 12 (AR) NM_001171155.1	PET100	Pan-ethnic	1 in 387	99%	1 in 38600
Mitochondrial complex IV deficiency / Leigh syndrome, French Canadian type (AR) NM_133259.3	LRPPRC	Pan-ethnic	≤1 in 500	99%	Reduced
Mitochondrial DNA depletion syndrome-2 (AR) NM_004614.4	TK2	Pan-ethnic	≤1 in 500	99%	Reduced
Mitochondrial neurogastrointestinal encephalomyopathy (AR) NM_001953.4	TYMP	Pan-ethnic	≤1 in 500	99%	Reduced
Mitochondrial trifunctional protein deficiency (HADHB-related) (AR) NM_000183.2	HADHB	Pan-ethnic	≤1 in 500	99%	Reduced
MKKS-related conditions (AR) NM_018848.3	MKKS	Pan-ethnic	≤1 in 500	99%	Reduced
Molybdenum cofactor deficiency (MOCS1-related) (AR) NM_001358530.2	MOCS1	Pan-ethnic	1 in 226	99%	1 in 22500
Molybdenum cofactor deficiency (MOCS2-related) (AR) NM_004531.4	MOCS2B	Pan-ethnic	≤1 in 500	99%	Reduced
Molybdenum cofactor deficiency (MOCS2-related) (AR) NM_176806.3	MOCS2A	Pan-ethnic	≤1 in 500	99%	Reduced
MPL-related conditions (AR) NM_005373.2	MPL	Pan-ethnic	≤1 in 500	99%	Reduced

DISORDER (INHERITANCE)	GENE	ETHNICITY	CARRIER FREQUENCY	DETECTION RATE	RISK TO BE A CARRIER AFTER NEGATIVE RESULT
MPV17-related conditions (AR) NM_002437.4	MPV17	Pan-ethnic	≤1 in 500	99%	Reduced
Mucopolidosis type III gamma (AR) NM_032520.4	GNPTG	Pan-ethnic	≤1 in 500	99%	Reduced
Mucopolidosis type IV (AR) NM_020533.2	MCOLN1	Pan-ethnic	≤1 in 500	99%	Reduced
Mucopolysaccharidosis type I (AR) NM_000203.4	IDUA	Pan-ethnic	1 in 148	97%	1 in 4900
Mucopolysaccharidosis type IIIA (AR) NM_000199.3	SGSH	Pan-ethnic	1 in 215	99%	1 in 21400
Mucopolysaccharidosis type IIIB (AR) NM_000263.3	NAGLU	Pan-ethnic	1 in 224	99%	1 in 22300
Mucopolysaccharidosis type IIID (AR) NM_002076.3	GNS	Pan-ethnic	≤1 in 500	99%	Reduced
Mucopolysaccharidosis type IVA (AR) NM_000512.4	GALNS	Pan-ethnic	1 in 224	99%	1 in 22300
Mucopolysaccharidosis type IX (AR) NM_153281.1	HYAL1	Pan-ethnic	≤1 in 500	99%	Reduced
Mucopolysaccharidosis type VI (AR) NM_000046.3	ARSB	Pan-ethnic	1 in 250	99%	1 in 24900
Mucopolysaccharidosis type VII (AR) NM_000181.3	GUSB	Pan-ethnic	1 in 250	99%	1 in 24900
Mulibrey nanism (AR) NM_015294.4	TRIM37	Pan-ethnic	≤1 in 500	99%	Reduced
Multiple pterygium syndrome (AR) NM_005199.4	CHRNA3	Pan-ethnic	≤1 in 500	99%	Reduced
Multiple sulfatase deficiency (AR) NM_182760.3	SUMF1	Pan-ethnic	≤1 in 500	99%	Reduced
Muscular dystrophy-dystroglycanopathy (FKRP-related) (AR) NM_024301.4	FKRP	Pan-ethnic	1 in 158	99%	1 in 15700
Muscular dystrophy-dystroglycanopathy (FKTN-related) (AR) NM_001079802.1	FKTN	Pan-ethnic	≤1 in 500	99%	Reduced
Muscular dystrophy-dystroglycanopathy (LARGE1-related) (AR) NM_004737.4	LARGE1	Pan-ethnic	≤1 in 500	99%	Reduced
Muscular dystrophy-dystroglycanopathy (POMT1-related) (AR) NM_007171.3	POMT1	Pan-ethnic	1 in 268	99%	1 in 26700
Muscular dystrophy-dystroglycanopathy (POMT2-related) (AR) NM_013382.5	POMT2	Pan-ethnic	1 in 371	99%	1 in 37000
Muscular dystrophy-dystroglycanopathy (RXYLT1-related) (AR) NM_014254.2	RXYLT1	Pan-ethnic	≤1 in 500	99%	Reduced
MUSK-related conditions (AR) NM_005592.3	MUSK	Pan-ethnic	1 in 447	99%	1 in 44600
MVK-related conditions (AR) NM_000431.3	MVK	Pan-ethnic	≤1 in 500	99%	Reduced
MYO7A-related conditions (AR) NM_000260.3	MYO7A	Pan-ethnic	1 in 200	95%	1 in 3980
Myopathy, lactic acidosis, and sideroblastic anemia 1 (AR) NM_025215.5	PUS1	Pan-ethnic	≤1 in 500	99%	Reduced
Myotonia congenita (AR) NM_000083.2	CLCN1	Pan-ethnic	1 in 112	99%	1 in 11100
N-acetylglutamate synthase deficiency (AR) NM_153006.2	NAGS	Pan-ethnic	≤1 in 500	99%	Reduced
Nemaline myopathy 2 (AR) NM_001271208.1	NEB *	Pan-ethnic	1 in 158	95%	1 in 3140
Nephrogenic diabetes insipidus (AQP2-related) (AR) NM_000486.5	AQP2	Pan-ethnic	≤1 in 500	99%	Reduced

DISORDER (INHERITANCE)	GENE	ETHNICITY	CARRIER FREQUENCY	DETECTION RATE	RISK TO BE A CARRIER AFTER NEGATIVE RESULT
Nephronophthisis (INVS-related) (AR) NM_014425.3	INVS	Pan-ethnic	≤1 in 500	99%	Reduced
Nephronophthisis (NPHP1-related) (AR) NM_000272.3	NPHP1	Pan-ethnic	≤1 in 500	99%	Reduced
Neuronal ceroid lipofuscinosis type 1 (AR) NM_000310.3	PPT1	Pan-ethnic	1 in 199	98%	1 in 9900
Neuronal ceroid lipofuscinosis type 2 (AR) NM_000391.3	TPP1	Pan-ethnic	1 in 250	97%	1 in 8300
Neuronal ceroid lipofuscinosis type 5 (AR) NM_006493.2	CLN5	Pan-ethnic	≤1 in 500	99%	Reduced
Neuronal ceroid lipofuscinosis type 6 (AR) NM_017882.2	CLN6	Pan-ethnic	≤1 in 500	99%	Reduced
Neuronal ceroid lipofuscinosis type 8 (AR) NM_018941.3	CLN8	Pan-ethnic	≤1 in 500	99%	Reduced
Neuronal ceroid lipofuscinosis type 10 (AR) NM_001909.4	CTSD	Pan-ethnic	≤1 in 500	99%	Reduced
Niemann-Pick disease type C (NPC1-related) (AR) NM_000271.4	NPC1	Pan-ethnic	1 in 183	99%	1 in 18200
Niemann-Pick disease type C (NPC2-related) (AR) NM_006432.3	NPC2	Pan-ethnic	≤1 in 500	99%	Reduced
Niemann-Pick disease types A and B (AR) NM_000543.4	SMPD1	Pan-ethnic	1 in 250	95%	1 in 4980
Nijmegen breakage syndrome (AR) NM_002485.4	NBN	Pan-ethnic	≤1 in 500	99%	Reduced
Nonsyndromic deafness (LOXHD1-related) (AR) NM_144612.6	LOXHD1	Pan-ethnic	≤1 in 500	99%	Reduced
Nonsyndromic deafness (MYO15A-related) (AR) NM_016239.3	MYO15A	Pan-ethnic	≤1 in 500	99%	Reduced
Nonsyndromic deafness (OTOA-related) (AR) NM_144672.3	OTOA *	Pan-ethnic	≤1 in 500	88%	Reduced
Nonsyndromic deafness (SYNE4-related) (AR) NM_001039876.2	SYNE4	Pan-ethnic	≤1 in 500	99%	Reduced
Nonsyndromic deafness (TMC1-related) (AR) NM_138691.2	TMC1	Pan-ethnic	≤1 in 500	99%	Reduced
Nonsyndromic deafness (TMPRSS3-related) (AR) NM_024022.2	TMPRSS3	Pan-ethnic	≤1 in 500	99%	Reduced
Nonsyndromic intellectual disability (CC2D1A-related) (AR) NM_017721.5	CC2D1A	Pan-ethnic	≤1 in 500	99%	Reduced
NR2E3-related conditions (AR) NM_014249.3	NR2E3	Pan-ethnic	≤1 in 500	99%	Reduced
NSMCE3 deficiency (AR) NM_138704.3	NSMCE3	Pan-ethnic	≤1 in 500	99%	Reduced
Oculocutaneous albinism type 2 (AR) NM_000275.2	OCA2	Pan-ethnic	1 in 95	99%	1 in 9400
Oculocutaneous albinism type 3 (AR) NM_000550.2	TYRP1	Pan-ethnic	≤1 in 500	99%	Reduced
Oculocutaneous albinism type 4 (AR) NM_016180.4	SLC45A2	Pan-ethnic	1 in 158	99%	1 in 15700
Oculocutaneous albinism types 1A and 1B (AR) NM_000372.4	TYR *	Pan-ethnic	1 in 100	97%	1 in 3300
OPA3-related conditions (AR) NM_025136.3	OPA3	Pan-ethnic	≤1 in 500	99%	Reduced
Osteogenesis imperfecta (BMP1-related) (AR) NM_006129.4	BMP1	Pan-ethnic	≤1 in 500	99%	Reduced
Osteogenesis imperfecta (CRTAP-related) (AR) NM_006371.4	CRTAP	Pan-ethnic	≤1 in 500	99%	Reduced
Osteogenesis imperfecta (P3H1-related) (AR) NM_022356.3	P3H1	Pan-ethnic	≤1 in 500	99%	Reduced
Osteopetrosis (TCIRG1-related) (AR) NM_006019.3	TCIRG1	Pan-ethnic	1 in 317	99%	1 in 31600
OSTM1 deficiency associated osteopetrosis (AR) NM_014028.3	OSTM1	Pan-ethnic	≤1 in 500	99%	Reduced

DISORDER (INHERITANCE)	GENE	ETHNICITY	CARRIER FREQUENCY	DETECTION RATE	RISK TO BE A CARRIER AFTER NEGATIVE RESULT
OTOF-related conditions (AR) NM_194248.2	OTOF	Pan-ethnic	≤1 in 500	99%	Reduced
Pantothenate kinase-associated neurodegeneration (AR) NM_153638.2	PANK2	Pan-ethnic	1 in 289	99%	1 in 28800
Parkinson disease 15 (AR) NM_012179.3	FBXO7	Pan-ethnic	≤1 in 500	99%	Reduced
PCDH15-related conditions (AR) NM_033056.3	PCDH15	Pan-ethnic	1 in 400	99%	1 in 39900
PEX5-related conditions (AR) NM_001131025.1	PEX5	Pan-ethnic	≤1 in 500	99%	Reduced
PEX7-related conditions (AR) NM_000288.3	PEX7	Pan-ethnic	1 in 157	99%	1 in 15600
PGM3-congenital disorder of glycosylation (AR) NM_001199917.1	PGM3	Pan-ethnic	≤1 in 500	99%	Reduced
Phenylalanine hydroxylase deficiency (AR) NM_000277.1	PAH	Pan-ethnic	1 in 58	99%	1 in 5700
Phosphoglycerate dehydrogenase deficiency (AR) NM_006623.3	PHGDH	Pan-ethnic	≤1 in 500	99%	Reduced
PIGN-congenital disorder of glycosylation (AR) NM_176787.4	PIGN	Pan-ethnic	≤1 in 500	99%	Reduced
PJVK-related conditions (AR) NM_001042702.3	DFNB59	Pan-ethnic	≤1 in 500	99%	Reduced
PLA2G6-related conditions (AR) NM_003560.2	PLA2G6	Pan-ethnic	≤1 in 500	99%	Reduced
PLEKHG5-related conditions (AR) NM_020631.4	PLEKHG5	Pan-ethnic	≤1 in 500	99%	Reduced
POLG-related conditions (AR) NM_002693.2	POLG	Pan-ethnic	1 in 113	95%	1 in 2240
Polycystic kidney disease (PKHD1-related) (AR) NM_138694.3	PKHD1 *	Pan-ethnic	1 in 70	99%	1 in 6900
Polymicrogyria (ADGRG1-related) (AR) NM_005682.6	ADGRG1	Pan-ethnic	≤1 in 500	99%	Reduced
POMGNT1-related conditions (AR) NM_017739.3	POMGNT1	Pan-ethnic	≤1 in 500	99%	Reduced
Pontocerebellar hypoplasia (TSEN54-related) (AR) NM_207346.2	TSEN54	Pan-ethnic	≤1 in 500	99%	Reduced
Pontocerebellar hypoplasia type 1B (AR) NM_016042.3	EXOSC3	Pan-ethnic	≤1 in 500	99%	Reduced
Pontocerebellar hypoplasia type 2D (AR) NM_016955.3	SEPSECS	Pan-ethnic	≤1 in 500	99%	Reduced
Pontocerebellar hypoplasia type 6 (AR) NM_020320.3	RARS2	Pan-ethnic	≤1 in 500	99%	Reduced
Primary carnitine deficiency (AR) NM_003060.3	SLC22A5	Pan-ethnic	1 in 71	99%	1 in 7000
Primary ciliary dyskinesia (CCDC39-related) (AR) NM_181426.1	CCDC39	Pan-ethnic	1 in 211	99%	1 in 21000
Primary ciliary dyskinesia (CCDC103-related) (AR) NM_213607.2	CCDC103	Pan-ethnic	1 in 316	99%	1 in 31500
Primary ciliary dyskinesia (DNAH5-related) (AR) NM_001369.2	DNAH5	Pan-ethnic	1 in 109	99%	1 in 10800
Primary ciliary dyskinesia (DNAH11-related) (AR) NM_001277115.1	DNAH11	Pan-ethnic	1 in 211	99%	1 in 21000
Primary ciliary dyskinesia (DNAI1-related) (AR) NM_012144.3	DNAI1	Pan-ethnic	1 in 250	99%	1 in 24900
Primary ciliary dyskinesia (DNAI2-related) (AR) NM_023036.4	DNAI2	Pan-ethnic	1 in 354	99%	1 in 35300
Primary hyperoxaluria type 1 (AR) NM_000030.2	AGXT	Pan-ethnic	1 in 135	99%	1 in 13400
Primary hyperoxaluria type 2 (AR) NM_012203.1	GRHPR	Pan-ethnic	≤1 in 500	99%	Reduced
Primary hyperoxaluria type 3 (AR) NM_138413.3	HOGA1	Pan-ethnic	1 in 354	99%	1 in 35300

DISORDER (INHERITANCE)	GENE	ETHNICITY	CARRIER FREQUENCY	DETECTION RATE	RISK TO BE A CARRIER AFTER NEGATIVE RESULT
Primary microcephaly (MCPH1-related) (AR) NM_024596.4	MCPH1	Pan-ethnic	≤1 in 500	99%	Reduced
Progressive early-onset encephalopathy with brain atrophy and thin corpus callosum (PEBAT) (AR) NM_005993.4	TBCD	Pan-ethnic	≤1 in 500	99%	Reduced
Progressive pseudorheumatoid dysplasia (AR) NM_003880.3	WISP3	Pan-ethnic	≤1 in 500	99%	Reduced
Prolidase deficiency (AR) NM_000285.3	PEPD	Pan-ethnic	≤1 in 500	99%	Reduced
Propionic acidemia (PCCA-related) (AR) NM_000282.3	PCCA	Pan-ethnic	1 in 224	96%	1 in 5575
Propionic acidemia (PCCB-related) (AR) NM_000532.4	PCCB	Pan-ethnic	1 in 224	99%	1 in 22300
PSAP-related conditions (AR) NM_002778.3	PSAP	Pan-ethnic	≤1 in 500	99%	Reduced
Pycnodysostosis (AR) NM_000396.3	CTSK	Pan-ethnic	1 in 438	99%	1 in 43700
Pyridoxal 5'-phosphate-dependent epilepsy (AR) NM_018129.3	PNPO	Pan-ethnic	≤1 in 500	99%	Reduced
Pyridoxine-dependent epilepsy (ALDH7A1-related) (AR) NM_001182.4	ALDH7A1	Pan-ethnic	1 in 127	99%	1 in 12600
Pyruvate carboxylase deficiency (AR) NM_000920.3	PC	Pan-ethnic	1 in 250	95%	1 in 4980
Pyruvate dehydrogenase complex deficiency (PDHB-related) (AR) NM_000925.3	PDHB	Pan-ethnic	≤1 in 500	99%	Reduced
RAPSN-related conditions (AR) NM_005055.4	RAPSN	Pan-ethnic	1 in 283	99%	1 in 28200
RDH12-related conditions (AR) NM_152443.2	RDH12	Pan-ethnic	1 in 460	99%	1 in 45900
Refsum disease (PHYH-related) (AR) NM_006214.3	PHYH	Pan-ethnic	≤1 in 500	99%	Reduced
Retinitis pigmentosa 25 (AR) NM_001142800.1	EYS *	Pan-ethnic	1 in 129	99%	1 in 12800
Retinitis pigmentosa 28 (AR) NM_001201543.1	FAM161A	Pan-ethnic	1 in 289	99%	1 in 28800
Retinitis pigmentosa 36 (AR) NM_001077620.2	PRCD	Pan-ethnic	1 in 296	99%	1 in 29500
Retinitis pigmentosa 62 (AR) NM_001242957.2	MAK	Pan-ethnic	1 in 274	99%	1 in 27300
Rhizomelic chondrodysplasia punctata type 2 (AR) NM_014236.3	GNPAT	Pan-ethnic	≤1 in 500	99%	Reduced
Rhizomelic chondrodysplasia punctata type 3 (AR) NM_003659.3	AGPS	Pan-ethnic	≤1 in 500	99%	Reduced
RLBP1-related conditions (AR) NM_000326.4	RLBP1	Pan-ethnic	1 in 296	99%	1 in 29500
Roberts syndrome (AR) NM_001017420.2	ESCO2	Pan-ethnic	≤1 in 500	99%	Reduced
RPE65-related conditions (AR) NM_000329.2	RPE65	Pan-ethnic	1 in 228	99%	1 in 22700
RYR1-related conditions (AR) NM_000540.2	RYR1	Pan-ethnic	≤1 in 500	99%	Reduced
SAMD9-related conditions (AR) NM_017654.3	SAMD9	Pan-ethnic	≤1 in 500	99%	Reduced
Sandhoff disease (AR) NM_000521.3	HEXB	Pan-ethnic	1 in 180	99%	1 in 17900
Schimke immuno-osseous dysplasia (AR) NM_014140.3	SMARCA1	Pan-ethnic	≤1 in 500	99%	Reduced
Seckel syndrome (CEP152-related) (AR) NM_014985.3	CEP152	Pan-ethnic	≤1 in 500	99%	Reduced
Sepiapterin reductase deficiency (AR) NM_003124.4	SPR	Pan-ethnic	≤1 in 500	99%	Reduced

DISORDER (INHERITANCE)	GENE	ETHNICITY	CARRIER FREQUENCY	DETECTION RATE	RISK TO BE A CARRIER AFTER NEGATIVE RESULT
Severe combined immunodeficiency due to CD3-delta deficiency (AR) NM_000732.4	CD3D	Pan-ethnic	≤1 in 500	99%	Reduced
Severe combined immunodeficiency due to CD3-epsilon deficiency (AR) NM_000733.3	CD3E	Pan-ethnic	≤1 in 500	99%	Reduced
Severe combined immunodeficiency due to CD45 deficiency (AR) NM_002838.4	PTPRC *	Pan-ethnic	≤1 in 500	99%	Reduced
Severe combined immunodeficiency due to DCLRE1C (Artemis) deficiency (AR) NM_001033855.2	DCLRE1C	Pan-ethnic	≤1 in 500	99%	Reduced
Severe combined immunodeficiency due to IL7R-alpha deficiency (AR) NM_002185.3	IL7R	Pan-ethnic	1 in 348	99%	1 in 34700
Severe combined immunodeficiency due to JAK3 deficiency (AR) NM_000215.3	JAK3	Pan-ethnic	1 in 455	99%	1 in 45400
Severe combined immunodeficiency due to RAG1 deficiency (AR) NM_000448.2	RAG1	Pan-ethnic	1 in 301	99%	1 in 30000
Severe combined immunodeficiency due to RAG2 deficiency (AR) NM_000536.3	RAG2	Pan-ethnic	≤1 in 500	99%	Reduced
Severe congenital neutropenia due to G6PC3 deficiency (AR) NM_138387.3	G6PC3	Pan-ethnic	≤1 in 500	99%	Reduced
Severe congenital neutropenia due to HAX1 deficiency (AR) NM_006118.3	HAX1	Pan-ethnic	≤1 in 500	99%	Reduced
Severe congenital neutropenia due to VPS45 deficiency (AR) NM_007259.4	VPS45	Pan-ethnic	≤1 in 500	99%	Reduced
Sialic acid storage diseases (AR) NM_012434.4	SLC17A5	Pan-ethnic	≤1 in 500	99%	Reduced
Sialidosis (AR) NM_000434.3	NEU1	Pan-ethnic	≤1 in 500	99%	Reduced
Sjögren-Larsson syndrome (AR) NM_000382.2	ALDH3A2	Pan-ethnic	≤1 in 500	99%	Reduced
SLC12A6-related conditions (AR) NM_133647.1	SLC12A6	Pan-ethnic	≤1 in 500	99%	Reduced
SLC26A2-related conditions (AR) NM_000112.3	SLC26A2	Pan-ethnic	1 in 158	95%	1 in 3140
SLC26A4-related conditions (AR) NM_000441.1	SLC26A4	Pan-ethnic	1 in 80	99%	1 in 7900
SLC37A4-related conditions (AR) NM_001164277.1	SLC37A4	Pan-ethnic	1 in 354	95%	1 in 7060
Smith-Lemli-Opitz syndrome (AR) NM_001360.2	DHCR7	Pan-ethnic	1 in 71	99%	1 in 7000
Spastic paraplegia type 15 (AR) NM_015346.3	ZFYVE26	Pan-ethnic	≤1 in 500	99%	Reduced
Spastic paraplegia type 49 (AR) NM_014844.3	TECPR2	Pan-ethnic	≤1 in 500	99%	Reduced
Spastic tetraplegia, thin corpus callosum, and progressive microcephaly (AR) NM_003038.4	SLC1A4	Pan-ethnic	≤1 in 500	99%	Reduced
SPG11-related conditions (AR) NM_025137.3	SPG11	Pan-ethnic	1 in 141	99%	1 in 14000
Spinal muscular atrophy (AR) NM_000344.3 Carrier residual risks listed are for 2 copy SMN1 results. Carrier residual risk for >2 copies are 5- to 10-fold lower.	SMN1 *	African-American	1 in 59	83%	1 in 342
		Ashkenazi Jewish	1 in 62	94%	1 in 1017
		Asian	1 in 50	93%	1 in 701
		Caucasian	1 in 45	95%	1 in 880
		Hispanic	1 in 48	94%	1 in 784
		Pan-ethnic	1 in 49	94%	1 in 800

DISORDER (INHERITANCE)	GENE	ETHNICITY	CARRIER FREQUENCY	DETECTION RATE	RISK TO BE A CARRIER AFTER NEGATIVE RESULT
Spinocerebellar ataxia (ANO10-related) (AR) NM_018075.3	ANO10 *	Pan-ethnic	≤1 in 500	99%	Reduced
Spondylocostal dysostosis (DLL3-related) (AR) NM_016941.3	DLL3	Pan-ethnic	1 in 350	99%	1 in 34900
Spondylocostal dysostosis (MESP2-related) (AR) NM_001039958.1	MESP2	Pan-ethnic	1 in 224	99%	1 in 22300
Steel syndrome (AR) NM_032888.3	COL27A1	Pan-ethnic	≤1 in 500	99%	Reduced
Steroid 5-alpha-reductase deficiency (AR) NM_000348.3	SRD5A2	Pan-ethnic	≤1 in 500	99%	Reduced
Stüve-Wiedemann syndrome (AR) NM_002310.5	LIFR *	Pan-ethnic	≤1 in 500	99%	Reduced
Sulfite oxidase deficiency (AR) NM_000456.2	SUOX	Pan-ethnic	≤1 in 500	99%	Reduced
SURF1-related conditions (AR) NM_003172.3	SURF1	Pan-ethnic	1 in 128	99%	1 in 12700
Tay-Sachs disease (AR) NM_000520.4	HEXA	Pan-ethnic	1 in 250	99%	1 in 24900
TBCE-related conditions (AR) NM_003193.4	TBCE *	Pan-ethnic	≤1 in 500	99%	Reduced
Thiamine-responsive megaloblastic anemia (AR) NM_006996.2	SLC19A2	Pan-ethnic	≤1 in 500	99%	Reduced
Thyroid dysharmonogenesis (SLC5A5-related) (AR) NM_000453.2	SLC5A5	Pan-ethnic	≤1 in 500	99%	Reduced
Thyroid dysharmonogenesis (TG-related) (AR) NM_003235.4	TG *	Pan-ethnic	≤1 in 500	99%	Reduced
Thyroid dysharmonogenesis (TPO-related) (AR) NM_000547.5	TPO	Pan-ethnic	1 in 129	99%	1 in 12800
TMEM67-related conditions (AR) NM_153704.5	TMEM67	Pan-ethnic	1 in 316	99%	1 in 31500
Transcobalamin II deficiency (AR) NM_000355.3	TCN2	Pan-ethnic	≤1 in 500	99%	Reduced
Transient infantile liver failure (AR) NM_018006.4	TRMU	Pan-ethnic	≤1 in 500	99%	Reduced
TREX1-related conditions (AR) NM_033629.4	TREX1	Pan-ethnic	≤1 in 500	99%	Reduced
Trichohepatoenteric syndrome (SKIV2L-related) (AR) NM_006929.4	SKIV2L	Pan-ethnic	≤1 in 500	99%	Reduced
Trichohepatoenteric syndrome (TTC37-related) (AR) NM_014639.3	TTC37	Pan-ethnic	≤1 in 500	99%	Reduced
TRIM32-related conditions (AR) NM_012210.3	TRIM32	Pan-ethnic	1 in 408	99%	1 in 40700
Trimethylaminuria (AR) NM_006894.6	FMO3	Pan-ethnic	≤1 in 500	99%	Reduced
Triple A syndrome (AR) NM_015665.5	AAAS	Pan-ethnic	≤1 in 500	99%	Reduced
TSHR-related conditions (AR) NM_000369.2	TSHR	Pan-ethnic	1 in 158	99%	1 in 15700
TULP1-related conditions (AR) NM_003322.4	TULP1	Pan-ethnic	1 in 296	99%	1 in 29500
Tyrosine hydroxylase deficiency (AR) NM_199292.2	TH	Pan-ethnic	≤1 in 500	99%	Reduced
Tyrosinemia type I (AR) NM_000137.2	FAH *	Pan-ethnic	1 in 125	95%	1 in 2480
Tyrosinemia type II (AR) NM_000353.2	TAT	Pan-ethnic	1 in 250	99%	1 in 24900
Tyrosinemia type III (AR) NM_002150.2	HPD	Pan-ethnic	≤1 in 500	99%	Reduced
USH1C-related conditions (AR) NM_005709.3	USH1C *	Pan-ethnic	1 in 353	90%	1 in 3521
USH2A-related conditions (AR) NM_206933.2	USH2A	Pan-ethnic	1 in 112	99%	1 in 11100

DISORDER (INHERITANCE)	GENE	ETHNICITY	CARRIER FREQUENCY	DETECTION RATE	RISK TO BE A CARRIER AFTER NEGATIVE RESULT
Very long-chain acyl-CoA dehydrogenase deficiency (AR) NM_000018.3	ACADVL	Pan-ethnic	1 in 100	99%	1 in 9900
Vici syndrome (AR) NM_020964.2	EPG5	Pan-ethnic	≤1 in 500	99%	Reduced
Vitamin D-dependent rickets type 1A (AR) NM_000785.3	CYP27B1	Pan-ethnic	≤1 in 500	99%	Reduced
Vitamin D-dependent rickets type 2A (AR) NM_001017535.1	VDR	Pan-ethnic	≤1 in 500	99%	Reduced
VPS53-related conditions (AR) NM_001128159.2	VPS53 *	Pan-ethnic	≤1 in 500	99%	Reduced
VRK1-related conditions (AR) NM_003384.2	VRK1	Pan-ethnic	≤1 in 500	99%	Reduced
VSX2-related conditions (AR) NM_182894.2	VSX2	Pan-ethnic	≤1 in 500	99%	Reduced
Warsaw syndrome (AR) NM_030653.3	DDX11 *	Pan-ethnic	≤1 in 500	15%	Reduced
Werner syndrome (AR) NM_000553.4	WRN *	Pan-ethnic	1 in 224	99%	1 in 22300
Wilson disease (AR) NM_000053.3	ATP7B	Pan-ethnic	1 in 90	98%	1 in 4450
WNT10A-related conditions (AR) NM_025216.2	WNT10A	Pan-ethnic	1 in 305	99%	1 in 30400
Wolcott-Rallison syndrome (AR) NM_004836.6	EIF2AK3	Pan-ethnic	≤1 in 500	99%	Reduced
Woodhouse-Sakati syndrome (AR) NM_025000.3	DCAF17	Pan-ethnic	≤1 in 500	99%	Reduced
Xeroderma pigmentosum complementation group A (AR) NM_000380.3	XPA	Pan-ethnic	≤1 in 500	99%	Reduced
Xeroderma pigmentosum complementation group C (AR) NM_004628.4	XPC	Pan-ethnic	≤1 in 500	99%	Reduced
Xeroderma pigmentosum, variant type (AR) NM_006502.2	POLH	Pan-ethnic	≤1 in 500	99%	Reduced
Zellweger spectrum disorder (PEX1-related) (AR) NM_000466.2	PEX1 *	Pan-ethnic	1 in 144	99%	1 in 14300
Zellweger spectrum disorder (PEX2-related) (AR) NM_000318.2	PEX2	Pan-ethnic	≤1 in 500	99%	Reduced
Zellweger spectrum disorder (PEX6-related) (AR) NM_000287.3	PEX6	Pan-ethnic	1 in 294	99%	1 in 29300
Zellweger spectrum disorder (PEX10-related) (AR) NM_153818.1	PEX10	Pan-ethnic	≤1 in 500	94%	Reduced
Zellweger spectrum disorder (PEX12-related) (AR) NM_000286.2	PEX12	Pan-ethnic	1 in 409	99%	1 in 40800
Zellweger spectrum disorder (PEX13-related) (AR) NM_002618.3	PEX13	Pan-ethnic	≤1 in 500	99%	Reduced
Zellweger spectrum disorder (PEX16-related) (AR) NM_004813.2	PEX16	Pan-ethnic	≤1 in 500	99%	Reduced
Zellweger spectrum disorder (PEX26-related) (AR) NM_017929.5	PEX26	Pan-ethnic	≤1 in 500	99%	Reduced

15 Crawford St., STE 100
Needham, MA 02494
(p) 626-350-0537
(f) 626-454-1667
Lab Director: Arash Radfar M.D.
CLIA: 22D0957540



Patient Information:

13057, Donor

DOB: [REDACTED]

Sex: M

MR#:

Patient#: FT-PT9113855

Partner Information:

Not Tested

Physician:

Kuan, James

ATTN: Martinez, Lupe

Seattle Sperm Bank

4915 25th Avenue NE, Ste 204W

Seattle, WA 98105

Phone: (206) 588-1484

Laboratory:

Fulgent Therapeutics LLC

CAP#: 8042697

CLIA#: 05D2043189

Laboratory Director:

Dr. Amar Jariwala

Report Date: **Jun 17, 2025**

Accession:

FT-7352242

Test#: FT-TS15227152

Specimen Type: Blood (EDTA)

Collected: Jun 07, 2025

Accession:

N/A

FINAL RESULTS



No carrier mutations identified

TEST PERFORMED

Single Gene Carrier Screening: FTCD

(1 Gene Panel: *FTCD*; gene sequencing with deletion and duplication analysis)

INTERPRETATION:

Notes and Recommendations:

- No carrier mutations were identified in the submitted specimen. A negative result does not rule out the possibility of a genetic predisposition nor does it rule out any pathogenic mutations in areas not assessed by this test or in regions that were covered at a level too low to reliably assess. Also, it does not rule out mutations that are of the sort not queried by this test; see Methods and Limitations for more information. A negative result reduces, but does not eliminate, the chance to be a carrier for any condition included in this screen. Please see the supplemental table for details.
- This carrier screening test does not screen for all possible genetic conditions, nor for all possible mutations in every gene tested. This report does not include variants of uncertain significance; only variants classified as pathogenic or likely pathogenic at the time of testing, and considered relevant for reproductive carrier screening, are reported. Please see the gene specific notes for details. Please note that the classification of variants can change over time.
- Patients may wish to discuss any carrier results with blood relatives, as there is an increased chance that they are also carriers. These results should be interpreted in the context of this individual's clinical findings, biochemical profile, and family history.
- Gene specific notes and limitations may be present. See below.
- Genetic counseling is recommended. Available genetic counselors and additional resources can be found at the National Society of Genetic Counselors (NSGC; <https://www.nsgc.org>)



GENES TESTED:

Custom Beacon Preconception Carrier Screening Panel - Gene

This analysis was run using the Custom Beacon Preconception Carrier Screening Panel gene list. 1 genes were tested with 100.0% of targets sequenced at >20x coverage. For more gene-specific information and assistance with residual risk calculation, see the SUPPLEMENTAL TABLE.

FTCD

METHODS:

Genomic DNA was isolated from the submitted specimen indicated above (if cellular material was submitted). DNA was barcoded, and enriched for the coding exons of targeted genes using hybrid capture technology. Prepared DNA libraries were then sequenced using a Next Generation Sequencing technology. Following alignment to the human genome reference sequence (assembly GRCh37), variants were detected in regions of at least 10x coverage. For this specimen, 100.00% and 100.00% of coding regions and splicing junctions of genes listed had been sequenced with coverage of at least 10x and 20x, respectively, by NGS or by Sanger sequencing. The remaining regions did not have 10x coverage, and were not evaluated. Variants were interpreted manually using locus specific databases, literature searches, and other molecular biological principles. To minimize false positive results, any variants that do not meet internal quality standards are confirmed by Sanger sequencing. Variants classified as pathogenic, likely pathogenic, or risk allele which are located in the coding regions and nearby intronic regions (+/- 20bp) of the genes listed above are reported. Variants outside these intervals may be reported but are typically not guaranteed. When a single pathogenic or likely pathogenic variant is identified in a clinically relevant gene with autosomal recessive inheritance, the laboratory will attempt to ensure 100% coverage of coding sequences either through NGS or Sanger sequencing technologies ("fill-in"). All genes listed were evaluated for large deletions and/or duplications. However, single exon deletions or duplications will not be detected in this assay, nor will copy number alterations in regions of genes with significant pseudogenes. Putative deletions or duplications are analyzed using Fulgent Germline proprietary pipeline for this specimen. Bioinformatics: The FPLMv2.0 pipeline was used to analyze this specimen.

LIMITATIONS:

General Limitations

These test results and variant interpretation are based on the proper identification of the submitted specimen, accuracy of any stated familial relationships, and use of the correct human reference sequences at the queried loci. In very rare instances, errors may result due to mix-up or co-mingling of specimens. Positive results do not imply that there are no other contributors, genetic or otherwise, to future pregnancies, and negative results do not rule out the genetic risk to a pregnancy. Official gene names change over time. Fulgent uses the most up to date gene names based on HUGO Gene Nomenclature Committee (<https://www.genenames.org>) recommendations. If the gene name on report does not match that of ordered gene, please contact the laboratory and details can be provided. Result interpretation is based on the available clinical and family history information for this individual, collected published information, and Alamut annotation available at the time of reporting. This assay is not designed or validated for the detection of low-level mosaicism or somatic mutations. This assay will not detect certain types of genomic aberrations such as translocations, inversions, or repeat expansions other than specified genes. DNA alterations in regulatory regions or deep intronic regions (greater than 20bp from an exon) may not be detected by this test. Unless otherwise indicated, no additional assays have been performed to evaluate genetic changes in this specimen. There are technical limitations on the ability of DNA sequencing to detect small insertions and deletions. Our laboratory uses a sensitive detection algorithm, however these types of alterations are not detected as reliably as single nucleotide variants. Rarely, due to systematic chemical, computational, or human error, DNA variants may be missed. Although next generation sequencing technologies and our bioinformatics analysis significantly reduce the confounding contribution of pseudogene sequences or other highly-homologous sequences, sometimes these may still interfere with the technical ability of the assay to identify pathogenic alterations in both sequencing and deletion/duplication analyses. Deletion/duplication analysis can identify alterations of genomic regions which include one whole gene (buccal swab specimens and whole blood specimens) and are two or more contiguous exons in size (whole blood specimens only); single exon deletions or duplications may occasionally be identified, but are not routinely detected by this test. When novel DNA duplications are identified, it is not possible to discern the genomic location or orientation of the duplicated segment, hence the effect of the duplication cannot be predicted. Where deletions are detected, it is not always possible to determine whether the predicted product will remain in-frame or not. Unless otherwise indicated, deletion/duplication analysis has not been performed in regions that have been sequenced by Sanger.



Gene Specific Notes and Limitations

FTCD: The current testing method may not reliably detect copy number variants in exons 8-10 (NM_006657.2) of this gene due to the high GC content.

SIGNATURE:



Dr. Harry Gao, DABMG, FACMG on 6/17/2025
Laboratory Director, Fulgent

DISCLAIMER:

This test was developed, performed, and its performance characteristics determined by **Fulgent Therapeutics LLC** (CAP# 8042697, CLIA# 05D2043189), 4399 Santa Anita Ave., El Monte, CA 91731. It has not been cleared or approved by the FDA. The laboratory is regulated under CLIA as qualified to perform high-complexity testing. This test is used for clinical purposes. It should not be regarded as investigational or for research. Since genetic variation, as well as systematic and technical factors, can affect the accuracy of testing, the results of testing should always be interpreted in the context of clinical and familial data. For assistance with interpretation of these results, healthcare professionals may contact us directly at **626-350-0537** or by email at info@fulgentgenetics.com. It is recommended that patients receive appropriate genetic counseling to explain the implications of the test result, including its residual risks, uncertainties and reproductive or medical options.

To view the supplemental table describing the carrier frequencies, detection rates, and residual risks associated with the genes tested on any Beacon panel, please visit the following link:
[Beacon Expanded Carrier Screening Supplemental Table](#)





Patient Information	Specimen Information	Client Information
13057, DONOR DOB: [REDACTED] AGE: [REDACTED] Gender: M Fasting: U Phone: 206.588.1484 Patient ID: 13057	Specimen: OW916905V Requisition: 0000520 Collected: 06/07/2023 / 13:30 PDT Received: 06/08/2023 / 04:34 PDT Reported: 06/15/2023 / 17:17 PDT	Client #: 98105026 VNLZR00 OLLIFFE, JEFFREY F SEATTLE SPERM BANK 4915 25TH AVE NE STE 204W SEATTLE, WA 98105-5668

COMMENTS: FASTING:UNKNOWN

Cytogenetic Report

CHROMOSOME ANALYSIS, BLOOD - 14596

Lab:EZ

CHROMOSOME ANALYSIS, BLOOD

Order ID: 23-252853
Specimen Type: Blood
Clinical Indication: GAMETE DONOR

RESULT:

NORMAL MALE KARYOTYPE

INTERPRETATION:

Chromosome analysis revealed normal G-band patterns within the limits of standard cytogenetic analysis.

Please expect the results of any other concurrent study in a separate report.

NOMENCLATURE:

46,XY

ASSAY INFORMATION:

Method: G-Band (Digital Analysis: MetaSyst)
Cells Counted: 20
Band Level: 500
Cells Analyzed: 5
Cells Karyotyped: 5


This test does not address genetic disorders that cannot be detected by standard cytogenetic methods or rare events such as low level mosaicism or subtle rearrangements.

Reha M. Toydemir, MD, PhD, FACMG

Electronic Signature: 6/15/2023 7:31 PM



Patient Information	Specimen Information	Client Information
13057, DONOR DOB: ████████ AGE: ██████ Gender: M Fasting: U Patient ID: 13057	Specimen: OW916905V Collected: 06/07/2023 / 13:30 PDT Received: 06/08/2023 / 04:34 PDT Reported: 06/15/2023 / 17:17 PDT	Client #: 98105026 OLLIFFE, JEFFREY F



PERFORMING SITE:

EZ QUEST DIAGNOSTICS/NICHOLS SJC, 33608 ORTEGA HWY, SAN JUAN CAPISTRANO, CA 92675-2042 Laboratory Director: IRINA MARAMICA,MD,PHD,MBA, CLIA: 05D0643352

Date Collected: 06/07/2023

Date Received: 06/08/2023

Date Reported: 06/12/2023

Fasting: Not Given

Ordered Items: LP+12AC+CBC/D/Plt+UA+Rh+ABO...

General Comments & Additional Information

A courtesy copy of this report has been sent to 2064664696.

Date Collected: 06/07/2023

LP+12AC+CBC/D/Plt+UA+Rh+ABO...

Test	Current Result and Flag		Previous Result and Date	Units	Reference Interval
Glucose ⁰¹	89			mg/dL	70-99
Uric Acid ⁰¹	4.7			mg/dL	3.8-8.4
Therapeutic target for gout patients: <6.0					
BUN ⁰¹	14			mg/dL	6-20
Creatinine ⁰¹	0.89			mg/dL	0.76-1.27
eGFR	119			mL/min/1.73	>59
Calcium ⁰¹	9.5			mg/dL	8.7-10.2
Protein, Total ⁰¹	7.4			g/dL	6.0-8.5
Albumin ⁰¹	4.9			g/dL	4.1-5.2
Bilirubin, Total ⁰¹	0.5			mg/dL	0.0-1.2
Alkaline Phosphatase ⁰¹	74			IU/L	44-121
LDH ⁰¹	192			IU/L	121-224
AST (SGOT) ⁰¹	29			IU/L	0-40
ALT (SGPT) ⁰¹	35			IU/L	0-44
Cholesterol, Total ⁰¹	162			mg/dL	100-199
Triglycerides ⁰¹	62			mg/dL	0-149
HDL Cholesterol ⁰¹	46			mg/dL	>39
▲ LDL Chol Calc (NIH)	104	High		mg/dL	0-99
LDL/HDL Ratio	2.3			ratio	0.0-3.6
Please Note: ⁰¹					

LDL/HDL Ratio			
		Men	Women
1/2 Avg.Risk		1.0	1.5
Avg.Risk		3.6	3.2
2X Avg.Risk		6.2	5.0
3X Avg.Risk		8.0	6.1

Hgb Fractionation by CE: ⁰²				
Hgb F ⁰²	0.0		%	0.0-2.0
Hgb A ⁰²	97.2		%	96.4-98.8
Hgb A2 ⁰²	2.8		%	1.8-3.2
Hgb S ⁰²	0.0		%	0.0

Interpretation:⁰²

Normal hemoglobin present; no hemoglobin variant or beta thalassemia identified.

Note: Alpha thalassemia may not be detected by the Hgb Fractionation Cascade panel. If alpha thalassemia is suspected, Labcorp offers

LP+12AC+CBC/D/Plt+UA+Rh+ABO... (Cont.)

Alpha-Thalassemia DNA Analysis (#511172).

ABO Grouping ⁰¹	O		
Rh Factor ⁰¹	Positive	Please note: Prior records for this patient's ABO / Rh type are not available for additional verification.	
. ⁰¹			
CBC, Platelet Ct, and Diff ⁰¹			
WBC ⁰¹	8.6	x10E3/uL	3.4-10.8
RBC ⁰¹	4.21	x10E6/uL	4.14-5.80
Hemoglobin ⁰¹	13.0	g/dL	13.0-17.7
Hematocrit ⁰¹	38.2	%	37.5-51.0
MCV ⁰¹	91	fL	79-97
MCH ⁰¹	30.9	pg	26.6-33.0
MCHC ⁰¹	34.0	g/dL	31.5-35.7
RDW ⁰¹	11.9	%	11.6-15.4
Platelets ⁰¹	259	x10E3/uL	150-450
Neutrophils ⁰¹	71	%	Not Estab.
Lymphs ⁰¹	22	%	Not Estab.
Monocytes ⁰¹	6	%	Not Estab.
Eos ⁰¹	0	%	Not Estab.
Basos ⁰¹	1	%	Not Estab.
Neutrophils (Absolute) ⁰¹	6.1	x10E3/uL	1.4-7.0
Lymphs (Absolute) ⁰¹	1.9	x10E3/uL	0.7-3.1
Monocytes(Absolute) ⁰¹	0.6	x10E3/uL	0.1-0.9
Eos (Absolute) ⁰¹	0.0	x10E3/uL	0.0-0.4
Baso (Absolute) ⁰¹	0.0	x10E3/uL	0.0-0.2
Immature Granulocytes ⁰¹	0	%	Not Estab.
Immature Grans (Abs) ⁰¹	0.0	x10E3/uL	0.0-0.1
. ⁰¹			
Urinalysis Gross Exam ⁰¹			
Specific Gravity ⁰¹	1.007		1.005-1.030
pH ⁰¹	6.5		5.0-7.5
Urine-Color ⁰¹	Yellow		Yellow
Appearance ⁰¹	Clear		Clear
WBC Esterase ⁰¹	Negative		Negative
Protein ⁰¹	Negative		Negative/Trace
Glucose ⁰¹	Negative		Negative
Ketones ⁰¹	Negative		Negative
Occult Blood ⁰¹	Negative		Negative
Bilirubin ⁰¹	Negative		Negative
Urobilinogen,Semi-Qn ⁰¹	0.2	mg/dL	0.2-1.0
Nitrite, Urine ⁰¹	Negative		Negative
Microscopic Examination ⁰¹	Microscopic not indicated and not performed.		

13057, Donor

Patient ID: 13057
Specimen ID: 158-129-1206-0

DOB: [REDACTED]
Age: [REDACTED]
Sex: Male

Patient Report

Account Number: 46857540
Ordering Physician: J OLLIFFE



Disclaimer

The Previous Result is listed for the most recent test performed by Labcorp in the past 5 years where there is sufficient patient demographic data to match the result to the patient. Results from certain tests are excluded from the Previous Result display.

Icon Legend

▲ Out of Reference Range ■ Critical or Alert

Performing Labs

01: SE - Labcorp Seattle 550 17th Avenue Ste 300, Seattle, WA, 98122-5789 Dir: Daniel Toweill, MD
02: BN - Labcorp Burlington 1447 York Court, Burlington, NC, 27215-3361 Dir: Sanjai Nagendra, MD
For Inquiries, the physician may contact Branch: 206-861-7000 Lab: 206-861-7000

Patient Details
13057, Donor

Phone:
Date of Birth: [REDACTED]
Age: [REDACTED]
Sex: Male
Patient ID: 13057
Alternate Patient ID: 13057

Physician Details
J OLLIFFE
Seattle Sperm Bank
4915 25th Ave Ne Ste 204, SEATTLE, WA, 98105

Phone: 206-588-1484
Account Number: 46857540
Physician ID:
NPI: 1306838271

Specimen Details
Specimen ID: 158-129-1206-0
Control ID: L2303813927
Alternate Control Number: L2303813927
Date Collected: 06/07/2023 1350 Local
Date Received: 06/08/2023 0000 ET
Date Entered: 06/08/2023 0044 ET
Date Reported: 06/12/2023 1007 ET