15 Crawford St., STE 100
Needham, MA 02494
(p) 626-350-0537
(f) 626-454-1667
Lab Director: Arash Radfar M.D.
CLIA: 22D0957540

fulgent | INFORM DIAGNOSTICS™ A FULGENT GENETICS COMPANY
Metroplex Pathology Associates
Lakewood Pathology Associates
Cohen Dermatopathology
Therapath Partners

Patient Information:
10912, Donor
DOB: [redacted]
Sex: M
MR#:
Patient#: FT-PT8770582

Partner Information:
Not Tested

Physician:
Kuan, James
ATTN: Kuan, James
Phoenix Sperm Bank
4915 25th Avenue NE, Ste 204W
Seattle, WA 98105
Phone: (206) 588-1484

Laboratory:
Fulgent Therapeutics LLC
CAP#: 8042697
CLIA#: 05D2043189
Laboratory Director:
Lawrence M. Weiss, MD
Report Date: **Jul 17,2024**

Accession:
FT-7017364
Test#: FT-TS14877457
Specimen Type: Blood (EDTA)
Collected: Jun 24,2024

Accession:
N/A

FINAL RESULTS



Carrier for genetic conditions in **multiple** genes.
Genetic counseling is recommended.

TEST PERFORMED

Beacon Preconception Carrier Screening - 515 Genes (without X-linked Disorders)
(515 Gene Panel; gene sequencing with deletion and duplication analysis)

Condition and Gene	Inheritance	10912, Donor	Partner
Congenital adrenal hyperplasia due to 21-hydroxylase deficiency *CYP21A2*	AR	⊕ Carrier c.92C>T (p.Pro31Leu)	N/A
Familial hemophagocytic lymphohistiocytosis type 3 *UNC13D*	AR	⊕ Carrier c.2346_2349del (p.Arg782Serfs*12)	N/A
Hereditary fructose intolerance *ALDOB*	AR	⊕ Carrier c.448G>C (p.Ala150Pro)	N/A

INTERPRETATION:

Notes and Recommendations:

- Based on these results, this individual is positive for carrier mutations in 3 genes. Carrier screening for the reproductive partner is recommended to accurately assess the risk for any autosomal recessive conditions. A negative result reduces, but does not eliminate, the chance to be a carrier for any condition included in this screen. Please see the supplemental table for details.
- Testing for copy number changes in the SMN1 gene was performed to screen for the carrier status of Spinal Muscular Atrophy. The results for this individual are within the normal range for non-carriers. See Limitations section for more information.
- This carrier screening test does not screen for all possible genetic conditions, nor for all possible mutations in every gene tested. This report does not include variants of uncertain significance; only variants classified as pathogenic or likely pathogenic at the time of testing, and considered relevant for reproductive carrier screening, are reported. Please see the gene specific notes for details. Please note that the classification of variants can change over time.
- Patients may wish to discuss any carrier results with blood relatives, as there is an increased chance that they are also carriers. These results should be interpreted in the context of this individual's clinical findings, biochemical profile, and family history.
- X-linked genes are not routinely analyzed for male carrier screening tests. Gene specific notes and limitations may be present. See below.
- Genetic counseling is recommended. Available genetic counselors and additional resources can be found at the National Society of Genetic Counselors (NSGC; https://www.nsgc.org)

15 Crawford St., STE 100
Needham, MA 02494
(p) 626-350-0537
(f) 626-454-1667
Lab Director: Arash Radfar M.D.
CLIA: 22D0957540

fulgent | INFORM DIAGNOSTICS™ A FULGENT GENETICS COMPANY
Metroplex Pathology Associates
Lakewood Pathology Associates
Cohen Dermatopathology
Therapath Partners

CONGENITAL ADRENAL HYPERPLASIA DUE TO 21-HYDROXYLASE DEFICIENCY

Patient	10912, Donor	Partner
Result	Carrier	N/A
Variant Details	***CYP21A2*** (NM_000500.9) c.92C>T (p.Pro31Leu)	N/A

What is Congenital adrenal hyperplasia due to 21-hydroxylase deficiency?

Congenital adrenal hyperplasia (CAH) due to 21-hydroxylase deficiency is an inherited disorder that affects the adrenal glands and hormone production. Approximately 75 percent of individuals with classic 21-hydroxylase deficiency have the salt-wasting type, whereby the body excretes too much salt in urine. Affected infants present with poor feeding, weight loss, dehydration, and vomiting, all of which can be life-threatening. Females with this condition typically have ambiguous genitalia, while males usually have normal genitalia, but with small testes. Individuals with the simple virilizing form and the non-classic form of the disease do not experience salt loss. Males and females with either the classic or non-classic forms of 21-hydroxylase deficiency tend to have an early growth spurt, but their final adult height is usually shorter than others in their family, and affected individuals may have reduced fertility. Additionally, individuals may have excessive body hair growth, hair loss, and irregular menstruation. Some individuals (male or female) with the non-classic form of the disease may have mild, non-life-threatening symptoms, while others may never develop symptoms of the disorder at all.

What is my risk of having an affected child?

CAH due to 21-hydroxylase deficiency is inherited in an autosomal recessive manner. If the patient and the partner are both carriers, the risk for an affected child is 1 in 4 (25%).

What kind of medical management is available?

Treatment consists of early initiation of hormone replacement therapy and/or surgery for females. Prognosis is good for patients with appropriate medical management and psychological support.

What mutation was detected?

The detected heterozygous variant was NM_000500.9:c.92C>T (p.Pro31Leu). This variant, p.Pro31Leu (also described as p.Pro30Leu), is a commonly reported pathogenic variant in the CYP21A2 gene. It has associated with a non-classic and simple virilizing forms of congenital adrenal hyperplasia (PubMed: 20301350, 10720040, 2072928, 23359698). Functional studies have shown that this variant significantly reduces normal protein activity (PubMed: 24953648, 23927611, 28539365)This variant is classified as "Pathogenic" in ClinVar, with multiple submitters in agreement (ClinVar: 12153). The laboratory classifies this variant as pathogenic.

15 Crawford St., STE 100
Needham, MA 02494
(p) 626-350-0537
(f) 626-454-1667
Lab Director: Arash Radfar M.D.
CLIA: 22D0957540

fulgent | INFORM DIAGNOSTICS™ A FULGENT GENETICS COMPANY
Metroplex Pathology Associates
Lakewood Pathology Associates
Cohen Dermatopathology
Therapath Partners

FAMILIAL HEMOPHAGOCYTIC LYMPHOHISTIOCYTOSIS TYPE 3

Patient	10912, Donor	Partner
Result	Carrier	N/A
Variant Details	***UNC13D*** (NM_199242.3) c.2346_2349del (p.Arg782Serfs*12)	N/A

What is Familial hemophagocytic lymphohistiocytosis type 3?

Familial hemophagocytic lymphohistiocytosis is an inherited condition where an individual's immune system overproduces activated immune cells (lymphocytes) and other immune system proteins (cytokines). This condition also causes the blood-producing cells in an individual's bone marrow to be destroyed, resulting in low red blood cells (anemia) and platelets. Primary symptoms of this condition usually appear during infancy and include fever, liver and spleen damage, fatigue, easy bruising, and abnormal bleeding. Brain, heart, kidney, and other organ abnormalities can be seen. Affected individuals also have an increased risk for leukemia and lymphoma. Onset of this subtype is often in infancy.

What is my risk of having an affected child?

Familial hemophagocytic lymphohistiocytosis is inherited in an autosomal recessive manner. This means that if both parents are carriers, their risk of having an affected child is 1 in 4 (25%).

What kind of medical management is available?

Management for familial hemophagocytic lymphohistiocytosis will vary based on the severity and age of onset of symptoms. A stem cell or bone marrow transplant is considered the cure for this condition. Often chemotherapy or immunotherapy will be used for treatment before a transplant is available.

What mutation was detected?

The detected heterozygous variant was NM_199242.3:c.2346_2349del (p.Arg782Serfs*12). This variant is predicted to introduce a premature stop codon at least 50 nucleotides upstream of the canonical donor splice site of the penultimate exon and to result in the loss of function of the protein product due to nonsense-mediated mRNA decay (PubMed: 25741868, 30192042, 27618451, 11532962, 18066079). There's sufficient evidence that loss of function in this gene is a known disease mechanism for hemophagocytic lymphohistiocytosis (PubMed: 33746956, 32542393, 31388699, 14622600). This variant was previously reported in multiple patients with hemophagocytic lymphohistiocytosis (HLH), as homozygous in two unrelated individuals and as compound heterozygous with another likely pathogenic or pathogenic variant with unknown phase in multiple unrelated individuals (PubMed: 16278825, 20823128, 32542393, 28973083, 32500225). This variant is classified as "Pathogenic" in ClinVar, with multiple submitters in agreement (Variation ID: 410246). The laboratory classifies this variant as pathogenic.

15 Crawford St., STE 100
Needham, MA 02494
(p) 626-350-0537
(f) 626-454-1667
Lab Director: Arash Radfar M.D.
CLIA: 22D0957540

fulgent | INFORM DIAGNOSTICS™ A FULGENT GENETICS COMPANY
Metroplex Pathology Associates
Lakewood Pathology Associates
Cohen Dermatopathology
Therapath Partners

HEREDITARY FRUCTOSE INTOLERANCE

Patient	10912, Donor	Partner
Result	⊕ Carrier	N/A
Variant Details	***ALDOB*** (NM_000035.4) c.448G>C (p.Ala150Pro)	N/A

What is Hereditary fructose intolerance?

Hereditary fructose intolerance is a condition that affects a person's ability to digest fructose. Fructose is a simple sugar found primarily in fruits. Affected individuals develop signs and symptoms of the disorder in infancy when fruits, juices, or other foods containing fructose are introduced into the diet. After ingesting fructose, individuals with hereditary fructose intolerance may experience nausea, bloating, abdominal pain, diarrhea, vomiting, and hypoglycemia. Affected infants may have failure to thrive. Repeated ingestion of fructose-containing foods can lead to liver and kidney damage.

What is my risk of having an affected child?

Hereditary fructose intolerance is inherited in an autosomal recessive manner. If the patient and the partner are both carriers, the risk for an affected child is 1 in 4 (25%).

What kind of medical management is available?

If large quantities of fructose are ingested, the infant may acutely develop lethargy, seizures, and/or progressive coma. Untreated hereditary fructose intolerance may result in renal and hepatic failure. If identified and treated before permanent organ injury occurs, individuals with hereditary fructose intolerance can experience a normal quality of life and life expectancy.

What mutation was detected?

The detected heterozygous variant was NM_000035.4:c.448G>C (p.Ala150Pro). This variant, p.Ala150Pro (also known as p.Ala149Pro in the literature), is a common variant found in individuals with hereditary fructose intolerance (PubMed: 3383242, 2623136, 8096362, 11757579, 18541450). This variant is classified as "Pathogenic/Likely pathogenic" in ClinVar, with multiple submitters in agreement (ClinVar: 464).Functional analysis of this variant indicates that this change reduces substrate affinity, enzyme stability and activity (PubMed: 12417303, 12464284, 15733923). The laboratory classifies this variant as pathogenic.

15 Crawford St., STE 100
Needham, MA 02494
(p) 626-350-0537
(f) 626-454-1667
Lab Director: Arash Radfar M.D.
CLIA: 22D0957540

GENES TESTED:

Beacon Preconception Carrier Screening - 515 Genes (without X-linked Disorders) - 515 Genes

This analysis was run using the Beacon Preconception Carrier Screening - 515 Genes (without X-linked Disorders) gene list. 515 genes were tested with 99.4% of targets sequenced at >20x coverage. For more gene-specific information and assistance with residual risk calculation, see the SUPPLEMENTAL TABLE.

AAAS, ABCA12, ABCA3, ABCA4, ABCB11, ABCB4, ABCC2, ABCC8, ACAD9, ACADM, ACADVL, ACAT1, ACOX1, ACSF3, ADA, ADAMTS2, ADAMTSL4, ADGRG1, ADGRV1, AGA, AGL, AGPS, AGXT, AHI1, AIPL1, AIRE, ALDH3A2, ALDH7A1, ALDOB, ALG1, ALG6, ALMS1, ALPL, AMN, AMT, ANO10, AP1S1, AQP2, ARG1, ARL6, ARSA, ARSB, ASL, ASNS, ASPA, ASS1, ATM, ATP6V1B1, ATP7B, ATP8B1, BBS1, BBS10, BBS12, BBS2, BBS4, BBS5, BBS7, BBS9, BCKDHA, BCKDHB, BCS1L, BLM, BLOC1S3, BLOC1S6, BMP1, BRIP1, BSND, CAD, CANT1, CAPN3, CASQ2, CBS, CC2D1A, CC2D2A, CCDC103, CCDC39, CCDC88C, CD3D, CD3E, CD40, CD59, CDH23, CEP152, CEP290, CERKL, CFTR, CHAT, CHRNE, CHRNG, CIITA, CLCN1, CLN3, CLN5, CLN6, CLN8, CLRN1, CNGB3, COL11A2, COL17A1, COL27A1, COL4A3, COL4A4, COL7A1, COX15, CPS1, CPT1A, CPT2, CRB1, CRTAP, CRYL1, CTNS, CTSA, CTSC, CTSD, CTSK, CYBA, CYP11A1, CYP11B1, CYP11B2, CYP17A1, CYP19A1, CYP1B1, CYP21A2, CYP27A1, CYP27B1, CYP7B1, DBT, DCAF17, DCLRE1C, DDX11, DGAT1, DGUOK, DHCR7, DHDDS, DLD, DLL3, DNAH11, DNAH5, DNAI1, DNAI2, DNMT3B, DOK7, DUOX2, DYNC2H1, DYSF, EIF2AK3, EIF2B1, EIF2B2, EIF2B3, EIF2B4, EIF2B5, ELP1, EPG5, ERCC2, ERCC6, ERCC8, ESCO2, ETFA, ETFB, ETFDH, ETHE1, EVC, EVC2, EXOSC3, EYS, FAH, FAM161A, FANCA, FANCC, FANCD2, FANCE, FANCG, FANCI, FANCL, FBP1, FBXO7, FH, FKBP10, FKRP, FKTN, FMO3, FOXN1, FOXRED1, FRAS1, FREM2, FUCA1, G6PC, G6PC3, GAA, GALC, GALE, GALK1, GALNS, GALNT3, GALT, GAMT, GATM, GBA, GBE1, GCDH, GCH1, GDF5, GFM1, GHR, GJB2, GJB6, GLB1, GLDC, GLE1, GNE, GNPAT, GNPTAB, GNPTG, GNS, GORAB, GRHPR, GRIP1, GSS, GUCY2D, GUSB, HADH, HADHA, HADHB, HAMP, HAX1, HBA1, HBA2, HBB, HEXA, HEXB, HGSNAT, HJV, HLCS, HMGCL, HMOX1, HOGA1, HPD, HPS1, HPS3, HPS4, HPS5, HPS6, HSD17B3, HSD17B4, HSD3B2, HYAL1, HYLS1, IDUA, IGHMBP2, IKBKB, IL7R, INVS, ITGA6, ITGB3, ITGB4, IVD, JAK3, KCNJ1, KCNJ11, LAMA2, LAMA3, LAMB3, LAMC2, LARGE1, LCA5, LDLR, LDLRAP1, LHX3, LIFR, LIG4, LIPA, LMBRD1, LOXHD1, LPL, LRAT, LRP2, LRPPRC, LYST, MAK, MAN2B1, MANBA, MCEE, MCOLN1, MCPH1, MECR, MED17, MESP2, MFSD8, MKKS, MKS1, MLC1, MLYCD, MMAA, MMAB, MMACHC, MMADHC, MOCS1, MOCS2, MPI, MPL, MPV17, MRE11, MTHFR, MTR, MTRR, MTTP, MUSK, MUT, MVK, MYO15A, MYO7A, NAGA, NAGLU, NAGS, NBN, NCF2, NDRG1, NDUFAF2, NDUFAF5, NDUFS4, NDUFS6, NDUFS7, NDUFV1, NEB, NEU1, NGLY1, NPC1, NPC2, NPHP1, NPHS1, NPHS2, NR2E3, NSMCE3, NTRK1, OAT, OCA2, OPA3, OSTM1, OTOA, OTOF, P3H1, PAH, PANK2, PC, PCBD1, PCCA, PCCB, PCDH15, PCNT, PDHB, PEPD, PET100, PEX1, PEX10, PEX12, PEX13, PEX16, PEX2, PEX26, PEX5, PEX6, PEX7, PFKM, PGM3, PHGDH, PHKB, PHKG2, PHYH, PIGN, PJVK, PKHD1, PLA2G6, PLEKHG5, PLOD1, PMM2, PNPO, POLG, POLH, POMGNT1, POMT1, POMT2, POR, POU1F1, PPT1, PRCD, PRDM5, PRF1, PROP1, PSAP, PTPRC, PTS, PUS1, PYGM, QDPR, RAB23, RAG1, RAG2, RAPSN, RARS2, RDH12, RLBP1, RMRP, RNASEH2A, RNASEH2B, RNASEH2C, RPE65, RPGRIP1L, RTEL1, RXYLT1, RYR1, SACS, SAMD9, SAMHD1, SCO2, SEC23B, SEPSECS, SGCA, SGCB, SGCD, SGCG, SGSH, SKIV2L, SLC12A1, SLC12A3, SLC12A6, SLC17A5, SLC19A2, SLC19A3, SLC1A4, SLC22A5, SLC25A13, SLC25A15, SLC25A20, SLC26A2, SLC26A3, SLC26A4, SLC27A4, SLC35A3, SLC37A4, SLC38A8, SLC39A4, SLC45A2, SLC4A11, SLC5A5, SLC7A7, SMARCAL1, SMN1, SMPD1, SNAP29, SPG11, SPR, SRD5A2, ST3GAL5, STAR, STX11, STXBP2, SUMF1, SUOX, SURF1, SYNE4, TANGO2, TAT, TBCD, TBCE, TCIRG1, TCN2, TECPR2, TERT, TF, TFR2, TG, TGM1, TH, TK2, TMC1, TMEM216, TMEM67, TMPRSS3, TPO, TPP1, TREX1, TRIM32, TRIM37, TRMU, TSEN54, TSFM, TSHB, TSHR, TTC37, TTPA, TULP1, TYMP, TYR, TYRP1, UBR1, UNC13D, USH1C, USH2A, VDR, VLDLR, VPS11, VPS13A, VPS13B, VPS45, VPS53, VRK1, VSX2, WISP3, WNT10A, WRN, XPA, XPC, ZBTB24, ZFYVE26, ZNF469

METHODS:

Genomic DNA was isolated from the submitted specimen indicated above (if cellular material was submitted). DNA was barcoded, and enriched for the coding exons of targeted genes using hybrid capture technology. Prepared DNA libraries were then sequenced using a Next Generation Sequencing technology. Following alignment to the human genome reference sequence (assembly GRCh37), variants were detected in regions of at least 10x coverage. For this specimen, 99.47% and 99.42% of coding regions and splicing junctions of genes listed had been sequenced with coverage of at least 10x and 20x, respectively, by NGS or by Sanger sequencing. The remaining regions did not have 10x coverage, and were not evaluated. Variants were interpreted manually using locus specific databases, literature searches, and other molecular biological principles. To minimize false positive results, any variants that do not meet internal quality standards are confirmed by Sanger sequencing. Variants classified as pathogenic, likely pathogenic, or risk allele which are located in the coding regions and nearby intronic regions (+/- 20bp) of the genes listed above are reported. Variants outside these intervals may be reported but are typically not guaranteed. When a single pathogenic or likely pathogenic variant is identified in a clinically relevant gene with autosomal recessive inheritance, the laboratory will attempt to ensure 100% coverage of coding sequences either through NGS or Sanger sequencing technologies ("fill-in"). All genes listed were evaluated for large deletions and/or duplications. However, single exon deletions or duplications will not be detected in this assay, nor will copy number alterations in regions of genes with significant pseudogenes. Putative deletions or duplications are analyzed using Fulgent Germline proprietary pipeline for this specimen. Bioinformatics: The Fulgent Germline v2019.2 pipeline was used to analyze this specimen.

LIMITATIONS:

General Limitations

These test results and variant interpretation are based on the proper identification of the submitted specimen, accuracy of any stated familial relationships, and use of the correct human reference sequences at the queried loci. In very rare instances, errors may result due to mix-up or co-mingling of specimens. Positive results do not imply that there are no other contributors, genetic or

15 Crawford St., STE 100
Needham, MA 02494
(p) 626-350-0537
(f) 626-454-1667
Lab Director: Arash Radfar M.D.
CLIA: 22D0957540

otherwise, to future pregnancies, and negative results do not rule out the genetic risk to a pregnancy. Official gene names change over time. Fulgent uses the most up to date gene names based on HUGO Gene Nomenclature Committee (https://www.genenames.org) recommendations. If the gene name on report does not match that of ordered gene, please contact the laboratory and details can be provided. Result interpretation is based on the available clinical and family history information for this individual, collected published information, and Alamut annotation available at the time of reporting. This assay is not designed or validated for the detection of low-level mosaicism or somatic mutations. This assay will not detect certain types of genomic aberrations such as translocations, inversions, or repeat expansions other than specified genes. DNA alterations in regulatory regions or deep intronic regions (greater than 20bp from an exon) may not be detected by this test. Unless otherwise indicated, no additional assays have been performed to evaluate genetic changes in this specimen. There are technical limitations on the ability of DNA sequencing to detect small insertions and deletions. Our laboratory uses a sensitive detection algorithm, however these types of alterations are not detected as reliably as single nucleotide variants. Rarely, due to systematic chemical, computational, or human error, DNA variants may be missed. Although next generation sequencing technologies and our bioinformatics analysis significantly reduce the confounding contribution of pseudogene sequences or other highly-homologous sequences, sometimes these may still interfere with the technical ability of the assay to identify pathogenic alterations in both sequencing and deletion/duplication analyses. Deletion/duplication analysis can identify alterations of genomic regions which include one whole gene (buccal swab specimens and whole blood specimens) and are two or more contiguous exons in size (whole blood specimens only); single exon deletions or duplications may occasionally be identified, but are not routinely detected by this test. When novel DNA duplications are identified, it is not possible to discern the genomic location or orientation of the duplicated segment, hence the effect of the duplication cannot be predicted. Where deletions are detected, it is not always possible to determine whether the predicted product will remain in-frame or not. Unless otherwise indicated, deletion/duplication analysis has not been performed in regions that have been sequenced by Sanger.

Gene Specific Notes and Limitations

ALG1: Due to the interference by highly homologous regions, our current testing method has less sensitivity to detect variants in exons 6-13 of the ALG1 gene (NM_019109.4). *CEP290:* Copy number analysis for exons 8-13 and exons 39-42 may have reduced sensitivity in the CEP290 gene. Confirmation of these exons are limited to individuals with a positive personal history of CEP290-related conditions and/or individuals carrying a pathogenic/likely pathogenic sequence variant. *CFTR:* Analysis of the intron 8 polymorphic region (e.g. IVS8-5T allele) is only performed if the p.Arg117His (R117H) mutation is detected. Single exon deletion/duplication analysis is limited to deletions of previously reported exons: 1, 2, 3, 11, 19, 20, 21. Analysis of the intron 8 polymorphic region (e.g. IVS8-5T allele) is only performed if the p.Arg117His (R117H) mutation is detected. Single exon deletion/duplication analysis is limited to deletions of previously reported exons: 1, 2, 3, 11, 19, 20, 21. CFTR variants primarily associated with CFTR-related isolated congenital bilateral absence of the vas deferens and CFTR-related pancreatitis are not included in this analysis. CFTR variants with insufficient evidence of being cystic fibrosis mutations will not be reported either. *CRYL1:* As mutations in the CRYL1 gene are not known to be associated with any clinical condition, sequence variants in this gene are not analyzed. However, to increase copy number detection sensitivity for large deletions including this gene and a neighboring on gene on the panel (GJB6, also known as connexin 30), this gene was evaluated for copy number variation. *CYP11B1:* The current testing method is not able to reliably detect certain pathogenic variants in this gene due to the interference by highly homologous regions. This analysis is not designed to detect or rule-out copy-neutral chimeric CYP11B1/CYP11B2 gene. *CYP11B2:* The current testing method is not able to reliably detect certain pathogenic variants in this gene due to the interference by highly homologous regions. This analysis is not designed to detect or rule-out copy-neutral chimeric CYP11B1/CYP11B2 gene. *CYP21A2:* Significant pseudogene interference and/or reciprocal exchanges between the CYP21A2 gene and its pseudogene, CYP21A1P, have been known to occur and may impact results. As such, the relevance of variants reported in this gene must be interpreted clinically in the context of the clinical findings, biochemical profile, and family history of each patient. LR-PCR is not routinely ordered for NM_000500.9:c.955C>T (p.Gln319Ter). Individuals with c.955C>T (p.Gln319Ter) will be reported as a Possible Carrier indicating that the precise nature of the variant has not been determined by LR-PCR and that the variant may occur in the CYP21A2 wild-type gene or in the CYP21A1P pseudogene. The confirmation test is recommended if the second reproductive partner is tested positive for variants associated with classic CAH. *DDX11:* Due to the interference by highly homologous regions, our current testing method has less sensitivity to detect variants in the DDX11 gene. *DUOX2:* The current testing method is not able to reliably detect variants in exons 6-8 of the DUOX2 gene (NM_014080.5) due to significant interference by the highly homologous gene, DUOX1. *FANCD2:* Due to pseudogene interference, copy-number-variants within exon 14-17 of the FANCD2 gene (NM _033084.4) are not evaluated and detection of single-nucleotide variants and small insertions/deletions in this region is not guaranteed. *GALT:* In general, the D2 "Duarte" allele is not reported if detected, but can be reported upon request. While this allele can cause positive newborn screening results, it is not known to cause clinical symptoms in any state. See GeneReviews for more information: https://www.ncbi.nlm.nih.gov/books/NBK1518/ *GBA:* Significant pseudogene interference and/or reciprocal exchanges between the GBA gene and its pseudogene, GBAP1, have been known to occur and may impact results. As such, the relevance of variants reported in this gene must be interpreted clinically in the context of this individual's clinical findings, biochemical profile, and family history. The current testing method cannot detect copy-neutral rearrangements between the pseudogene and the functional gene, which have been reported in very rare cases of Gaucher disease (PubMed: 21704274). *HBA1:* Significant interference

15 Crawford St., STE 100
Needham, MA 02494
(p) 626-350-0537
(f) 626-454-1667
Lab Director: Arash Radfar M.D.
CLIA: 22D0957540

from highly homologous regions in exons 1-2 of the HBA1 gene has been recognized to occur, potentially impeding the assay's technical capability to detect pathogenic alterations during sequencing analyses. *HBA2:* Significant interference from highly homologous regions in exons 1-2 of the HBA2 gene has been recognized to occur, potentially impeding the assay's technical capability to detect pathogenic alterations during sequencing analyses. *HSD17B4:* Copy number analysis for exons 4-6 may have reduced sensitivity in the HSD17B4 gene. Confirmation of these exons are limited to individuals with a positive personal history of D-bifunctional protein deficiency and Perrault syndrome and/or individuals carrying a pathogenic/likely pathogenic sequence variant. *LMBRD1:* Copy number analysis for exons 9-12 may have reduced sensitivity in the LMBRD1 gene. Confirmation of these exons are limited to individuals with a positive personal history of combined methylmalonic aciduria and homocystinuria and/or individuals carrying a pathogenic/likely pathogenic sequence variant. *MTHFR:* As recommended by ACMG, the two common polymorphisms in the MTHFR gene - c.1286A>C (p.Glu429Ala, also known as c.1298A>C) and c.665C>T (p.Ala222Val, also known as c.677C>T) - are not reported in this test due to lack of sufficient clinical utility to merit testing (PubMed: 23288205). *NEB:* This gene contains a 32-kb triplicate region (exons 82-105) which is not amenable to sequencing and deletion/duplication analysis. *NPHS2:* If detected, the variant NM_014625.3:c.686G>A (p.Arg229Gln) will not be reported as this variant is not significantly associated with disease when homozygous or in the compound heterozygous state with variants in exons 1-6 of NPHS2. *OTOA:* Due to pseudogene interference, our current testing method is not able to reliably detect variants in exons 20-28 (NM_144672.3) in the OTOA gene. *SMN1:* The current testing method detects sequencing variants in exon 7 and copy number variations in exons 7-8 of the SMN1 gene (NM_022874.2). Sequencing and deletion/duplication analysis are not performed on any other region in this gene. About 5%-8% of the population have two copies of SMN1 on a single chromosome and a deletion on the other chromosome, known as a [2+0] configuration (PubMed: 20301526). The current testing method cannot directly detect carriers with a [2+0] SMN1 configuration but can detect linkage between the silent carrier allele and certain population-specific single nucleotide changes. As a result, a negative result for carrier testing greatly reduces but does not eliminate the chance that a person is a carrier. Only abnormal results will be reported. *TERT:* The TERT promoter region is analyzed for both sequencing and copy number variants. *TYR:* Due to the interference by highly homologous regions, our current testing method has less sensitivity to detect variants in exons 4-5 of the TYR gene (NM_000372.5). *VPS45:* LoF is not a known disease mechanism *WRN:* Due to the interference by highly homologous regions within the WRN gene, our current testing method has less sensitivity to detect variants in exons 10-11 of WRN (NM_000553.6).

SIGNATURE:



Geetu Mendiratta-Vij, PhD, FACMG, CGMBS on **7/17/2024**
Laboratory Director, Fulgent

DISCLAIMER:

This test was developed and its performance characteristics determined by Fulgent Therapeutics LLC CAP #8042697 CLIA #05D2043189; 4399 Santa Anita Ave., El Monte, CA, 91731. It has not been cleared or approved by the FDA. The laboratory is regulated under CLIA as qualified to perform high-complexity testing. This test is used for clinical purposes. It should not be regarded as investigational or for research. Since genetic variation, as well as systematic and technical factors, can affect the accuracy of testing, the results of testing should always be interpreted in the context of clinical and familial data. For assistance with interpretation of these results, healthcare professionals may contact us directly at **626-350-0537** or by email at **info@fulgentgenetics.com**. It is recommended that patients receive appropriate genetic counseling to explain the implications of the test result, including its residual risks, uncertainties and reproductive or medical options.

To view the supplemental table describing the carrier frequencies, detection rates, and residual risks associated with the genes on this test please visit the following link:
Beacon Expanded Carrier Screening Supplemental Table

4399 Santa Anita Ave.
El Monte, CA, 91731
(p) 626-350-0537 (f) 626-454-1667
info@fulgentgenetics.com
www.fulgentgenetics.com

Patient Information:
10912, Donor
DOB: [redacted]
Sex: M
MR#:
Patient#: FT-PT8770582

Partner Information:
Not Tested

Physician:
Kuan, James
ATTN: Kirk, Ashley
Phoenix Sperm Bank
4915 25th Avenue Northeast
Ste 204
Seattle, 98105
Phone: 3039705897

Laboratory:
Fulgent Therapeutics LLC
CAP#: 8042697
CLIA#: 05D2043189
Laboratory Director:
Dr. Amar Jariwala
Report Date: **Jul 21,2026**

Accession:
FT-7017364
Test#: FT-TS15634234
Specimen Type: Blood (EDTA)
Collected: Jun 24,2024

Accession:
N/A

FINAL RESULTS

No carrier mutations identified

TEST PERFORMED

Single Gene Carrier Screening: CSPP1
(1 Gene Panel: *CSPP1*; gene sequencing with deletion and duplication analysis)

INTERPRETATION:

Notes and Recommendations:

- No carrier mutations were identified in the submitted specimen. A negative result does not rule out the possibility of a genetic predisposition nor does it rule out any pathogenic mutations in areas not assessed by this test or in regions that were covered at a level too low to reliably assess. Also, it does not rule out mutations that are of the sort not queried by this test; see Methods and Limitations for more information. A negative result reduces, but does not eliminate, the chance to be a carrier for any condition included in this screen. Please see the supplemental table for details.
- This carrier screening test does not screen for all possible genetic conditions, nor for all possible mutations in every gene tested. This report does not include variants of uncertain significance; only variants classified as pathogenic or likely pathogenic at the time of testing, and considered relevant for reproductive carrier screening, are reported. Please see the gene specific notes for details. Please note that the classification of variants can change over time.
- Patients may wish to discuss any carrier results with blood relatives, as there is an increased chance that they are also carriers. These results should be interpreted in the context of this individual's clinical findings, biochemical profile, and family history.
- Gene specific notes and limitations may be present. See below.
- Genetic counseling is recommended. Available genetic counselors and additional resources can be found at the National Society of Genetic Counselors (NSGC; https://www.nsgc.org).

4399 Santa Anita Ave.
El Monte, CA, 91731
(p) 626-350-0537 (f) 626-454-1667
info@fulgentgenetics.com
www.fulgentgenetics.com

GENES TESTED:

Custom Beacon Carrier Screening Panel - 1 Gene

This analysis was run using the Custom Beacon Carrier Screening Panel gene list. 1 genes were tested with 100.0% of targets sequenced at >20x coverage. For more gene-specific information and assistance with residual risk calculation, see the SUPPLEMENTAL TABLE.

CSPP1

METHODS:

Genomic DNA was isolated from the submitted specimen indicated above (if cellular material was submitted). DNA was barcoded, and enriched for the coding exons of targeted genes using hybrid capture technology. Prepared DNA libraries were then sequenced using a Next Generation Sequencing technology. Following alignment to the human genome reference sequence (assembly GRCh37), variants were detected in regions of at least 10x coverage. For this specimen, 100.00% and 100.00% of coding regions and splicing junctions of genes listed had been sequenced with coverage of at least 10x and 20x, respectively, by NGS or by Sanger sequencing. The remaining regions did not have 10x coverage, and were not evaluated. Variants were interpreted manually using locus specific databases, literature searches, and other molecular biological principles. To minimize false positive results, any variants that do not meet internal quality standards are confirmed by Sanger sequencing. Variants classified as pathogenic, likely pathogenic, or risk allele which are located in the coding regions and nearby intronic regions (+/- 20bp) of the genes listed above are reported. Variants outside these intervals may be reported but are typically not guaranteed. When a single pathogenic or likely pathogenic variant is identified in a clinically relevant gene with autosomal recessive inheritance, the laboratory will attempt to ensure 100% coverage of coding sequences either through NGS or Sanger sequencing technologies ("fill-in"). All genes listed were evaluated for large deletions and/or duplications. However, single exon deletions or duplications will not be detected in this assay, nor will copy number alterations in regions of genes with significant pseudogenes. Putative deletions or duplications are analyzed using Fulgent Germline proprietary pipeline for this specimen. Bioinformatics: The FPLMv2.0 pipeline was used to analyze this specimen.

LIMITATIONS:

General Limitations

These test results and variant interpretation are based on the proper identification of the submitted specimen, accuracy of any stated familial relationships, and use of the correct human reference sequences at the queried loci. In very rare instances, errors may result due to mix-up or co-mingling of specimens. Positive results do not imply that there are no other contributors, genetic or otherwise, to future pregnancies, and negative results do not rule out the genetic risk to a pregnancy. Official gene names change over time. Fulgent uses the most up to date gene names based on HUGO Gene Nomenclature Committee (https://www.genenames.org) recommendations. If the gene name on report does not match that of ordered gene, please contact the laboratory and details can be provided. Result interpretation is based on the available clinical and family history information for this individual, collected published information, and Alamut annotation available at the time of reporting. This assay is not designed or validated for the detection of low-level mosaicism or somatic mutations. This assay will not detect certain types of genomic aberrations such as translocations, inversions, or repeat expansions other than specified genes. DNA alterations in regulatory regions or deep intronic regions (greater than 20bp from an exon) may not be detected by this test. Unless otherwise indicated, no additional assays have been performed to evaluate genetic changes in this specimen. There are technical limitations on the ability of DNA sequencing to detect small insertions and deletions. Our laboratory uses a sensitive detection algorithm, however these types of alterations are not detected as reliably as single nucleotide variants. Rarely, due to systematic chemical, computational, or human error, DNA variants may be missed. Although next generation sequencing technologies and our bioinformatics analysis significantly reduce the confounding contribution of pseudogene sequences or other highly-homologous sequences, sometimes these may still interfere with the technical ability of the assay to identify pathogenic alterations in both sequencing and deletion/duplication analyses. Deletion/duplication analysis can identify alterations of genomic regions which include two or more contiguous exons in size (whole blood specimens only); single exon deletions or duplications may occasionally be identified, but are not routinely detected by this test. Saliva swab samples may have reduced sensitivity for CNVs and a higher CNV failure rate. When novel DNA duplications are identified, it is not possible to discern the genomic location or orientation of the duplicated segment, hence the effect of the duplication cannot be predicted. Where deletions are detected, it is not always possible to determine whether the predicted product will remain in-frame or not. Unless otherwise indicated, deletion/duplication analysis has not been performed in regions that have been sequenced by Sanger.

4399 Santa Anita Ave.
El Monte, CA, 91731
(p) 626-350-0537 (f) 626-454-1667
info@fulgentgenetics.com
www.fulgentgenetics.com

Gene Specific Notes and Limitations

No gene specific limitations apply to the genes on the tested panel.

SIGNATURE:

Dr. Harry Gao, DABMG, FACMG on **07/21/2026**
Laboratory Director, Fulgent

DISCLAIMER:

This test was developed, performed, and its performance characteristics determined by**Fulgent Therapeutics LLC** (CAP# 8042697, CLIA# 05D2043189), 4399 Santa Anita Ave., El Monte, CA 91731. It has not been cleared or approved by the FDA. The laboratory is regulated under CLIA as qualified to perform high-complexity testing. This test is used for clinical purposes. It should not be regarded as investigational or for research. Since genetic variation, as well as systematic and technical factors, can affect the accuracy of testing, the results of testing should always be interpreted in the context of clinical and familial data. For assistance with interpretation of these results, healthcare professionals may contact us directly at **(626) 350-0537** or **info@fulgentgenetics.com**. It is recommended that patients receive appropriate genetic counseling to explain the implications of the test result, including its residual risks, uncertainties and reproductive or medical options.

4399 Santa Anita Ave.
El Monte, CA, 91731
(p) 626-350-0537 (f) 626-454-1667
info@fulgentgenetics.com
www.fulgentgenetics.com

To view the supplemental table describing the carrier frequencies, detection rates, and residual risks associated with the genes tested on any Beacon panel, please visit the following link:
Beacon Expanded Carrier Screening Supplemental Table

Report Status FINAL
Route 2017 Ordered by:
Phoenix Sperm Bank
1492 S Mill Ave
Suite 306
Tempe, AZ 85281

James Kuan, MD

Sonora Quest Laboratories™
A Subsidiary of Laboratory Sciences of Arizona

Patient Information:
10912, DONOR

Order #: 181310000123 / NL101994864

Account: 18131
ID/MR#: 10912
Patient Lab ID:
667aeadb414ea744e6e02d19

Collected: 06/24/2024 11:45 AM
Received: 06/25/2024 08:41 AM
Reported: 07/02/2024 03:14 PM

DOB: [redacted] Age: [redacted]
Sex: M
Patient Phone: 602-888-7255

[redacted] PL

GENETICS

Accession #:
CG240006829
Cell Type/Source:
Blood
Clinician Provided Information:
DONOR TESTING

Chromosome Analysis: Routine Blood

Analysis Details: PV
Metaphases/Cells Counted : 20
Metaphases/Cells Analyzed : 5
Metaphases Karyotyped : 3

Results: PV
NORMAL MALE KARYOTYPE

46,XY

Interpretation: PV
Normal

PV

Normal karyotype at the band level 550 or above as determined by the trypsin-Giemsa method. There was no evidence for a chromosome abnormality within the limits of the band level and technology utilized in this study.

PHA-stimulated lymphocyte chromosome analysis is an accurate technique to detect many constitutional chromosome abnormalities. More extensive investigation may be required to detect mosaicism or subtle structural rearrangement. It also should be noted that this type of testing does not rule out the possibility of mendelian, mitochondrial, multifactorial or environmental etiologies.

Cytogenetics Director: PV

Electronically signed by Liwen Lai R PhD, DABMGG, FACMG R
Verified 07/02/24

Tests Ordered: Chromosome Analysis: Routine Blood

Unless otherwise noted, testing performed by: Sonora Quest Laboratories, 424 S 56th St, Phoenix, AZ 85034 800.766.6721
Testing noted as PV performed by: Genetics/Genomics Div., Sonora Quest Laboratories, 424 S. 56th St, Phoenix, AZ 85034 602.685.5700

End of Report

All Rights Reserved

10912, Donor
Patient ID:
Specimen ID: **176-705-1266-0**

DOB: [redacted]
Age: [redacted]
Sex: **Male**

Patient Report
Account Number: **02003540**
Ordering Physician: **J KUAN**

labcorp

Date Collected: **06/24/2024** Date Received: **06/25/2024** Date Reported: **06/27/2024** Fasting: **Not Given**

Ordered Items: **LP+12AC+CBC/D/Plt+UA+Rh+ABO...**

Date Collected: **06/24/2024**

LP+12AC+CBC/D/Plt+UA+Rh+ABO...

	Test	Current Result and Flag		Previous Result and Date	Units	Reference Interval
	Glucose [01]	92			mg/dL	70-99
	Uric Acid [01]	8.3			mg/dL	3.8-8.4
				Therapeutic target for gout patients: <6.0		
	BUN [01]	12			mg/dL	6-20
▲	**Creatinine** [01]	**1.30**	**High**		mg/dL	0.76-1.27
	eGFR	73			mL/min/1.73	>59
	Calcium [01]	9.3			mg/dL	8.7-10.2
	Protein, Total [01]	8.1			g/dL	6.0-8.5
	Albumin [01]	4.9			g/dL	4.1-5.1
	Bilirubin, Total [01]	0.5			mg/dL	0.0-1.2
	Alkaline Phosphatase [01]	60			IU/L	44-121
	LDH [01]	150			IU/L	121-224
	AST (SGOT) [01]	24			IU/L	0-40
	ALT (SGPT) [01]	23			IU/L	0-44
▲	**Cholesterol, Total** [01]	**214**	**High**		mg/dL	100-199
	Triglycerides [01]	113			mg/dL	0-149
	HDL Cholesterol [01]	50			mg/dL	>39
▲	**LDL Chol Calc (NIH)**	**144**	**High**		mg/dL	0-99
	LDL/HDL Ratio	2.9			ratio	0.0-3.6
	Please Note: [01]					

LDL/HDL Ratio

	Men	Women
1/2 Avg.Risk	1.0	1.5
Avg.Risk	3.6	3.2
2X Avg.Risk	6.2	5.0
3X Avg.Risk	8.0	6.1

	Hgb Fractionation by CE: [02]					
	Hgb F [02]	0.0			%	0.0-2.0
	Hgb A [02]	97.2			%	96.4-98.8
	Hgb A2 [02]	2.8			%	1.8-3.2
	Hgb S [02]	0.0			%	0.0
	Interpretation: [02]					

Normal hemoglobin present; no hemoglobin variant or beta thalassemia identified.
Note: Alpha thalassemia may not be detected by the Hgb Fractionation Cascade panel. If alpha thalassemia is suspected, Labcorp offers Alpha-Thalassemia DNA Analysis (#511172).

	ABO Grouping [01]	A				
	Rh Factor [01]	Positive				

Please note: Prior records for this patient's ABO / Rh type are not available for additional verification.

©2024 Laboratory Corporation of America® Holdings
All Rights Reserved - Enterprise Report Version 2.00

This document contains private and confidential health information protected by state and federal law.
If you have received this document in error please call 800-788-9743

Date Collected: **06/24/2024**

LP+12AC+CBC/D/Plt+UA+Rh+ABO... (Cont.)

.[01]				
CBC, Platelet Ct, and Diff[01]				
WBC[01]	4.9		x10E3/uL	3.4-10.8
RBC[01]	4.75		x10E6/uL	4.14-5.80
Hemoglobin[01]	14.8		g/dL	13.0-17.7
Hematocrit[01]	43.5		%	37.5-51.0
MCV[01]	92		fL	79-97
MCH[01]	31.2		pg	26.6-33.0
MCHC[01]	34.0		g/dL	31.5-35.7
RDW[01]	13.1		%	11.6-15.4
Platelets[01]	281		x10E3/uL	150-450
Neutrophils[01]	47		%	Not Estab.
Lymphs[01]	45		%	Not Estab.
Monocytes[01]	6		%	Not Estab.
Eos[01]	1		%	Not Estab.
Basos[01]	1		%	Not Estab.
Neutrophils (Absolute)[01]	2.3		x10E3/uL	1.4-7.0
Lymphs (Absolute)[01]	2.2		x10E3/uL	0.7-3.1
Monocytes(Absolute)[01]	0.3		x10E3/uL	0.1-0.9
Eos (Absolute)[01]	0.1		x10E3/uL	0.0-0.4
Baso (Absolute)[01]	0.0		x10E3/uL	0.0-0.2
Immature Granulocytes[01]	0		%	Not Estab.
Immature Grans (Abs)[01]	0.0		x10E3/uL	0.0-0.1
.[01]				
Urinalysis Gross Exam[01]				
▶ **Specific Gravity**[01]	**<=1.005**	**Abnormal**		1.005-1.030
pH[01]	6.0			5.0-7.5
Urine-Color[01]	Yellow			Yellow
Appearance[01]	Clear			Clear
WBC Esterase[01]	Negative			Negative
Protein[01]	Negative			Negative/Trace
Glucose[01]	Negative			Negative
Ketones[01]	Negative			Negative
Occult Blood[01]	Negative			Negative
Bilirubin[01]	Negative			Negative
Urobilinogen,Semi-Qn[01]	0.2		mg/dL	0.2-1.0
Nitrite, Urine[01]	Negative			Negative
Microscopic Examination[01]				
	Microscopic not indicated and not performed.			

©2024 Laboratory Corporation of America® Holdings
All Rights Reserved - Enterprise Report Version 2.00

This document contains private and confidential health information protected by state and federal law.
If you have received this document in error please call 800-788-9743

Disclaimer
The Previous Result is listed for the most recent test performed by Labcorp in the past 5 years where there is sufficient patient demographic data to match the result to the patient. Results from certain tests are excluded from the Previous Result display.

Icon Legend
▲ Out of Reference Range ■ Critical or Alert

Performing Labs
01: PDLCA - Labcorp Phoenix, 5005 S 40th Street Ste 1200, Phoenix, AZ 85040-2969 Dir: Earle Collum, MD
02: BN - Labcorp Burlington, 1447 York Court, Burlington, NC 27215-3361 Dir: Sanjai Nagendra, MD
For Inquiries, the physician may contact Branch: 888-522-2677 Lab: 800-788-9743

Patient Details
10912, Donor

Phone:
Date of Birth: [redacted]
Age: [redacted]
Sex: **Male**
Patient ID:
Alternate Patient ID:

Physician Details
J KUAN
Phoenix Sperm Bank
1492 S Mill Ave Suite 306, Tempe, AZ, 85281

Phone: **602-888-7255**
Account Number: **02003540**
Physician ID:
NPI: **1891885745**

Specimen Details
Specimen ID: **176-705-1266-0**
Control ID: **L2404824389**
Alternate Control Number: **L2404824389**
Date Collected: **06/24/2024 1034 Local**
Date Received: **06/25/2024 0000 ET**
Date Entered: **06/24/2024 2258 ET**
Date Reported: **06/27/2024 0909 ET**

©2024 Laboratory Corporation of America® Holdings
All Rights Reserved - Enterprise Report Version 2.00

This document contains private and confidential health information protected by state and federal law.
If you have received this document in error please call 800-788-9743